



US 20160175370A1

(19) **United States**

(12) **Patent Application Publication**
Zhou et al.

(10) **Pub. No.: US 2016/0175370 A1**

(43) **Pub. Date: Jun. 23, 2016**

(54) **TARGETING OF HERPES SIMPLEX VIRUS TO SPECIFIC RECEPTORS**

C12N 7/00 (2006.01)

A61K 47/48 (2006.01)

(71) Applicant: **THE UNIVERSITY OF CHICAGO**,
Chicago, IL (US)

(52) **U.S. Cl.**

CPC *A61K 35/763* (2013.01); *C12N 7/00*

(2013.01); *A61K 47/48269* (2013.01); *A61K*

47/48376 (2013.01); *C07K 14/5437* (2013.01);

C07K 16/18 (2013.01); *C12N 2710/16632*

(2013.01); *C12N 2710/16621* (2013.01); *C12N*

2710/16642 (2013.01); *C07K 2317/622*

(2013.01)

(72) Inventors: **Guoying Zhou**, Madison, WI (US);
Bernard Roizman, Chicago, IL (US)

(21) Appl. No.: **14/589,581**

(22) Filed: **Jan. 5, 2015**

Related U.S. Application Data

(63) Continuation of application No. 12/065,455, filed on Jul. 10, 2008, now Pat. No. 8,927,251, filed as application No. PCT/US06/33865 on Aug. 30, 2006, which is a continuation of application No. 11/215,636, filed on Aug. 30, 2005, now Pat. No. 7,550,148, which is a continuation-in-part of application No. 10/530,774, filed on Nov. 17, 2005, now Pat. No. 7,501,126, filed as application No. PCT/US03/31598 on Oct. 6, 2003.

(60) Provisional application No. 60/416,716, filed on Oct. 7, 2002.

Publication Classification

(51) **Int. Cl.**

A61K 35/763 (2006.01)

C07K 16/18 (2006.01)

C07K 14/54 (2006.01)

(57)

ABSTRACT

The invention relates to engineered Herpes simplex virus (HSV) particles that are targeted to one or more specific binding pair members, such as receptors. Also, recombinant vectors for producing such HSV particles are provided. By reducing the affinity of HSV for its natural receptor(s) and increasing the affinity for a selected receptor, the HSV particles of the invention are useful for targeting cells that express the selected receptor, which itself may be a product of genetic engineering. The ability to selectively target cells renders the HSV particles, particularly useful in selectively diagnosing, treating, and imaging cells bearing the selected binding pair member, such as a receptor. The invention also provides for polynucleotide-based therapy to cells bearing the selected binding pair member such as a receptor.




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 1D

The amino terminal sequence of IL13-gC

cttegtcgggaggcggcatcgaacgcaccccccatccpgtgggtccgtgtggaggtcgttttttcagtgc
 ggtctcgttttggccg99aacgfiaggctcATGGCGCTTTTGTGGACCAAGGTCAATTGCTCTCACCTGGCC
 gC upstream → IL-13*
 GCGGCTTTTGGCTTCCCAAGGCCCTGTGGCTCCCTCTACAGCCCTCAGGTAGCTCATTGAGGAGCTGGTCA
 CATCACCCAGAACTCA GAAGGCTCCGCTCTGCAATGGCAGCATGGTATGGAGCATCAACTGACAGCTGGC
 TGTACTGTGCAGCCCTGGAAATCCCTGATCAACGTGTCAAGCTGCAGTCCCATCGAGAAGACCCAGAGGAT
 CTGAGCGGATTCGTGCCCGCACAAAGGTCTCAGCTGGGCAGTTTTCCAGCTTGCATGTCCGAGACACCAAAA
 CGAGGTGGCCCAAGTTTGTAAAAGATCTGCTCTTACATTTAAAGAAACTTTTTTCGGCAGGGACGGTTgaak
 GCACCCCATGGAGTTCCGCTCCAGATATGGCGTTACTCCATGGGTCCGTCCCCCCCAATGCTCCGGC
 → gC downstream

FIG. 2A

The sequence of the gB_{Δpoly(K)} domain

GGGTCTTGGTGGCGTCCGGGGCTCCGAGTTCCCTCCGGCAGCCCTGGGCTCGCGGGCCGCAACCCAGGCGGC
 GAAAGGGGGACCTGCCACTTCGGCCCGCCCGCCCTGGGCCCCGCCCCCAACGGGGGAATCCGAAAACCGAAG
 AAGAACAGAAACCGAAACCCCCAAAGCGCCCGCCCGCCGGGACAAACCGGACCCCTCGCCGGGGCCA
 CGCCACCTTGGCCGAGCACCTGGGGACATCAAGGGGAGAACCCGATGCAAACTTTTACGTGTGCCCA
 CCCCCACGGGGGCCACGGTGGTGCAGTTCGAGCAGCCCGCCGCTGCCCCGACCCCGGCCGAGGGTCCAGA

FIG. 2B

The amino terminal sequence of IL13-gD

ATGGGGGGGGCTGCCGCCAGGTTGGGGGGCCGTGATTTTGTGTTGTCGTCAATAGCTGGGCTTC
 Signal peptide of gD →
 CATGGGGTCCGGGGCAAAATATGCCCTGGCGGATGCCCTCTCTCAAAGCTGGCCGACCCCAAT
 ←
 CGCTTTCGCCGCAAAAGACCTTCCGGTCCctcgag*ATGGCGCTTTTGTGGACCAAGGTCAAT
 24AA XhoI IL13 →
 GCTCTCACTTGCCTTGGCGGCTTTGCCCTCCCAAGGCCCTGTGCCTCCCTCTACAGCCCTC
 AAGGAGCTCATTTGAGGAGCTGGTCAACATCACCCAGAACAGAGGCTCCGCTCTGCAAT
 GGCAGCATGGTTTGGAGCATCAACCTGACAGCTGGCATGTA CTGTGCAGCCCTGGAAATCC
 CTGATCAACGTGTGAGGCTGCAGTGCATCGAGAAGACCCAGAGGATGCTGGGGCGGATTC
 TGGCCCGCACAAAGGTCTCAGCTGGGCAGTTTTCCAGCTTGCATGTCCGAGACACCAAAATC
 GAGGTGGCCCAAGTTTGTAAAAGGACCTGCTCTTACATTTAAAGAAACTTTTTTCGGCAGGGGA
 CGGTTCAACTGAAAC*ggtagccCTGGACCAGCTGACCCGACCCCTCCGGGGGTCCGGCCCGTG
 ← IL13 KpnI 25AA
 TACCACATCCAGGGGGGCTACCGGACCCGTTCCAGCCCCCAGCCCTCCCGATC

FIG. 2C




FIG. 3A




FIG. 3B




FIG. 4A

FIG. 4B

FIG. 4C





FIG. 5



gD- in pcDNA3.1(-) with CMV promoter collapsed by NruI/NheI digestion. The fragment containing gD upstream at NotI/BamHI, and gD downstream at XhoI/KpnI in pBluescript II SK was excised by NotI/KpnI and ligated into pcDNA3.1(-) in which the CMV promoter had been collapsed. N, NotI, B, BamHI, P, PstI, E, EcoRI, H, HindIII, C, ClaI, X, XhoI, and K, KpnI.

FIG. 6




FIG. 7

TARGETING OF HERPES SIMPLEX VIRUS TO SPECIFIC RECEPTORS

[0001] This application is a continuation-in-part application of U.S. Ser. No. 10/530,774, which is the US national phase of PCT/US03/31598 filed Oct. 6, 2003, which claims the priority benefit of U.S. Ser. No. 60/416,716, filed Oct. 7, 2002.

GOVERNMENT INTERESTS

[0002] The U.S. Government owns rights in the invention pursuant to National Cancer Institute grant number 1PO1 CA71933.

BACKGROUND OF THE INVENTION

[0003] A steady rate of healthcare advances has led to continuing improvement in the health and quality of life for humans and animals. Nevertheless, a variety of diseases, disorders, and conditions have largely eluded the best efforts at prevention or treatment. Chief among these maladies is the loss of cell-cycle control that frequently results in the undesirable cell proliferation characteristic of cancer in its many forms, such as malignant glioma. Malignant gliomas are devastating brain tumors that afflict animals such as humans. The average life span after diagnosis is less than one year and few patients have been reported to survive five years. Furthermore, none of the conventional anti-cancer therapies has been successful in significantly prolonging the lifespan of patients with this disease. In recent years there have been numerous attempts to use genetically engineered herpes simplex viruses (HSV) as oncolytic agents to treat malignant gliomas. Because wild-type viruses are highly virulent, the viruses used in preclinical evaluations and in phase-1 clinical studies have been thoroughly attenuated. While several deletion mutants have been tested, the mutants that reached clinical trials lacked the $\gamma_134.5$ gene encoding infected cell protein 34.5 (ICP34.5) and optionally, the U_L39 gene encoding the large subunit of ribonucleotide reductase.

[0004] These attenuated HSV viruses, however, have been imperfectly engineered as oncolytic agents. One advantage of these mutant viruses is that they have a significantly reduced capacity to replicate in normal, non-dividing cells *in vivo*. Viral ribonucleotide reductase is an essential gene for viral replication in resting cells and, hence, the U_L39 mutant virus is dysfunctional in the normal environment of the central nervous system (Simard et al. 1995). The major function of ICP34.5 is to preclude the shutoff of protein synthesis caused by activation of protein kinase R in infected cells. Once activated, this enzyme phosphorylates the a subunit of translation initiation factor 2 (eIF2 α), resulting in complete cessation of translation. Mutants lacking the $\gamma_134.5$ genes are highly attenuated because the lytic life cycle is completely blocked in an interferon⁺ cellular background. In contrast, $\gamma_134.5$ mutants are nearly as virulent as wild-type virus in mice lacking interferon receptor. Although mutants deleted in both $\gamma_134.5$ and U_L39 are not significantly more attenuated than those lacking the $\gamma_134.5$ genes, such mutants do provide added insurance in the form of a reduced risk of reversion.

[0005] A significant disadvantage of these mutant HSV viruses is their poor replication, even in dividing cells. In experimental animal systems, the mutant viruses do not exhibit sustained lytic life cycles, with the loss of a potentially amplified response to a given therapeutic dose of the virus that would be expected upon re-infection of tumor cells by the

multiplied viral progeny. Consequently, maximum killing of tumors cells requires high doses of virus. Given the poor growth of these mutant HSV viruses, even in dividing cells, production of virus pools large enough to yield efficacious inocula of $>10^9$ plaque forming units (PFU) has remained a major obstacle. Moreover, indiscriminate binding of virus to non-tumor cells further diminishes the effectiveness of HSV virus dosages because mis-targeted viral particles do not contribute to the desired beneficial therapeutic effect of tumor cell destruction. One approach to overcoming these obstacles is to achieve a more thorough understanding of the HSV lytic life cycle and thereby facilitate the development of HSV mutants tailored for use as targeted therapeutic agents, such as targeted oncolytic agents.

[0006] HSV enters host cells using a two-step mechanism. The first step of entry is HSV attachment to the cell surface. This step is initiated by glycoproteins B and C (gB and gC), which project from the viral envelope, attaching to heparan sulfate proteoglycans on host cell surfaces. The gB and gC domains interacting with heparan sulfate have been mapped at the sequence level (Laquerre et al. 1998). Following this initial attachment, viral glycoprotein D (gD) interacts with one of several receptors. Of these gD receptors, two are particularly important for entry (Spear et al. 2000). One receptor, designated HveA (formerly, HveM), is a member of the family of receptors for tumor necrosis proteins. A second receptor, designated Nectin-1 (HveC), is a member of the nectin family of proteins, structurally related to the immunoglobulin superfamily, which serve as intercellular connectors (Campadelli-Fiume et al. 2000). The second step of HSV entry into a cell is fusion of the viral envelope with the plasma membrane of the cell. To effect fusion, gD, when bound to its receptor, recruits glycoproteins B, H and L, which results in fusion of the envelope with the plasma membrane.

[0007] Additional understanding of HSV infection has come from recent studies that have lent significance to an old observation that gD interacts with the cation-independent mannose 6 phosphate receptor, contributing to the accumulation of HSV in endosomes. Endocytosis of viral particles results in particle degradation by lysosomal enzymes, but the cells succumb as a consequence of the degradation of cellular DNA by lysosomal DNase. HSV gD blocks this apoptotic pathway to cell death through its interaction with the mannose 6 phosphate receptor. Thus, gD interacts with HveA, nectins, the mannose 6 phosphate receptor, and at least one of the complex of viral glycoproteins involved in the fusion of HSV with the plasma membrane.

[0008] In an attempt to target HSV-1 infection to specific cells, a recombinant HSV having a chimeric protein comprising gC and erythropoietin (EPO) on its surface was constructed. Although the recombinant virus bound to cells expressing EPO receptor and endocytosis of the virus occurred, successful infection of these EPO-receptor expressing cells did not occur.

[0009] Accordingly, a need continues to exist in the art for viral therapeutic agents exhibiting improved targeting capacities while retaining sufficient capacity to infect to be therapeutically useful. Ideally, suitable viruses would be therapeutic agents, such as oncolytic agents, themselves as well as providing a targeting vehicle or vector for the controlled delivery of polynucleotide coding regions useful as therapeutic agents. Another need in the art is for targeted agents useful in diagnostic applications as, e.g., imaging agents or targeted vehicles for imaging agents.

SUMMARY

[0010] The invention satisfies the aforementioned need in the art by providing viral forms suitable for use as therapeutic and diagnostic agents themselves, as well as providing a ready vehicle for the delivery of therapeutic or diagnostic polynucleotide coding regions to cells. These viral forms are modified viruses of the Herpesviridae family of viruses, and are preferably derived from herpes simplex virus type 1 or type 2. The invention provides a method of making virus particles with a novel ligand (or binding pair member), and making said particles totally dependent on a receptor of the ligand (or binding pair member) for entry into targeted cells.

[0011] Disclosed herein are methods to modify the surface of, e.g., an HSV virus particle in a manner that targets the virus to a specific receptor present on the surface of a cell of choice, typically a cell in need of therapy or a cell whose presence provides information of diagnostic value. The invention provides viral particles, e.g., HSV particles, having a reduced affinity for their natural cell-surface receptor(s), and methods for producing and using such particles, which minimizes or eliminates the problem of reduced efficiency associated with the mis-targeting of therapeutic and diagnostic viruses. Additionally, the invention provides viral particles, e.g., HSV particles, that exhibit specific affinity for a cell surface component that is not a natural viral receptor and that is present solely or predominantly on a given target cell, as well as methods for producing and using such viruses. Modified viral particles (e.g., HSV) having increased affinity for a cell surface component associated with one or more target cells exhibit improved targeting capabilities relative to known viral particles. The modified HSV particles have reduced indiscriminate binding, thereby minimizing sequestration of viral dosages away from the target cells. The invention further provides modified viral particles, such as modified HSV particles, that have both a reduced affinity for natural viral receptors and an increased affinity for a cell surface component associated with a particular target cell(s), with the modified viral particle effectively recruiting that cell surface component for use as a viral receptor. Other benefits of the modified viruses are described herein and will be apparent to those of skill in the art upon review of this disclosure.

[0012] In one aspect, the invention provides a recombinant herpes simplex virus (HSV) particle having at least one protein on its surface, comprising: (a) an altered gD, wherein the alteration reduces binding of gD to one or more of its cellular receptors, said alteration comprising (i) a heterologous peptide ligand on the surface of the recombinant HSV particle forming a fusion protein with the altered gD; and (ii) an amino acid alteration; wherein said recombinant HSV particle preferentially binds to cells expressing a binding partner to said heterologous peptide ligand. In some embodiments, these particles preferentially bind to target cells (cell expressing a binding partner) in whole or part due to the greater frequency of the binding partner on the surface of the cell relative to any natural HSV binding proteins on the surface of that cell. In some embodiments, the recombinant HSV particle further comprises an altered viral surface protein, wherein the alteration reduces binding of the viral surface protein to a sulfated proteoglycan. Such recombinant herpes simplex virus (HSV) particles comprise a virus surface protein altered to reduce the wild-type level of binding of that protein to a sulfated proteoglycan on the surface of a cell and an altered gD. The altered gD exhibits a reduced binding to one or more of the natural cellular receptors for gD; the altered gD is also fused

to a heterologous peptide ligand (or binding pair member) having a binding partner, e.g., a peptide ligand receptor, found on the surface of a cell. Stated in the alternative, this aspect of the invention provides a recombinant herpes simplex virus (HSV) particle having at least one protein on its surface, comprising: (a) an altered viral surface protein, wherein the alteration reduces binding of the viral surface protein to a sulfated proteoglycan; and (b) an altered gD, wherein the alteration reduces binding of gD to one or more of its cellular receptors, the alteration comprising (i) a heterologous peptide ligand (or binding pair member) on the surface of the recombinant HSV particle forming a fusion protein with the altered gD; and (ii) an amino acid alteration; wherein the recombinant HSV particle preferentially binds to cells expressing a binding partner to the heterologous peptide ligand (or binding pair member).

[0013] The invention comprehends a recombinant HSV particle wherein the amino acid alteration is selected from the group consisting of an amino acid deletion, an amino acid substitution and an amino acid insertion. A preferred site for the amino acid alteration is amino acid position 34 of gD. Exemplary recombinant HSV particles according to the invention include HSV R5141 and HSV R5161, each described below.

[0014] Contemplated amino acid alterations include insertions or deletions of 1-10 amino acids, such as insertions or deletions of 1-5 amino acids. Exemplary insertions occur immediately upstream (N-terminal) or downstream (C-terminal) to amino acid position 34 of gD. Exemplary deletions include amino acid position 34 of gD. For alterations comprising amino acid substitutions, 1-10 amino acids are substituted, such as substitutions of 1-5 amino acids. Non-contiguous (dispersed) or contiguous amino acid substitutions are contemplated. In some embodiments, conservative amino acids known in the art are substituted. Exemplary amino acid substitutions include single amino acid substitutions for the Serine at position 34 of gD. Preferably, substitutions for Ser34 will be conservative as, for example, an amino acid substitution of V34S in gD.

[0015] The altered gD, moreover, reduces binding of the recombinant HSV particle to at least one HSV entry mediator (Hve) cell-surface protein, such as an Hve selected from the group consisting of HveA (formerly, HveM) and Nectin-1 (HveC). Further, the recombinant HSV particles of the invention include particles wherein the altered viral surface protein is selected from the group consisting of gB and gC. In some embodiments, the altered viral surface protein, preferably selected from the group of gB and gC, forms a fusion protein with a heterologous peptide ligand. In some embodiments, the binding partner is a cell surface receptor for the heterologous peptide ligand.

[0016] The preferential binding of the recombinant HSV particles of the invention results in a detectable variation in effective binding of the particle to the cells being compared. By "effective binding" is meant either sufficiently stable binding to permit detection of binding or binding sufficient to result in productive infection of the cell. In preferred embodiments, the preferential binding is such that the recombinant HSV particles bind only to one of the cell types being compared (e.g., cancer cells compared to healthy cells of the same type). Suitable cells include any hyperproliferative cell type, such as a cancer cell. A cancer cell, in turn, includes a tumor cell, e.g., a malignant glioma cell.

[0017] The invention comprehends recombinant HSV particles wherein the heterologous peptide ligand (or binding pair member) is any ligand (or binding pair member) for which a cell surface binding partner exists. Preferably, heterologous peptide ligands have specific cell surface binding partners, e.g., ligand receptors, that are preferentially exhibited on the surface of a target cell. More preferably, the cell surface binding partner is only exhibited on the surface of a target cell, when compared to the cells in an organism containing the target cell. Exemplary heterologous peptide ligands that include cytokines, such as IL13, and fragments, variants and derivatives thereof, provided that the ligand retains the capacity of binding to a cell-surface binding partner. An exemplary binding pair member contemplated as suitable for each aspect of the invention is a single-chain antibody, for which a binding partner would include an antigen thereof, or a fragment, derivative or variant thereof that retains the capacity to bind to the single-chain antibody.

[0018] Another aspect of the invention is drawn to the recombinant HSV particle described above, wherein a polynucleotide encoding the fusion protein is joined to a heterologous coding region for a leader sequence. In this context, "heterologous" means that the leader sequence is not found naturally associated with the upstream or 5' coding region participating in the fusion. Exemplary leader sequences include HSV leader sequences, e.g., an HSV gD leader sequence. In a related aspect, the invention provides the recombinant HSV particle described above, wherein a polynucleotide encoding the fusion protein is joined to a heterologous expression control element, such as a heterologous promoter (a promoter not naturally found in association with the polynucleotide coding region fused upstream or 5' in the fusion), a heterologous enhancer, or expression factor binding site known in the art.

[0019] Another aspect of the invention provides a pharmaceutical composition comprising a recombinant HSV particle described above and a pharmaceutically acceptable carrier, diluent, or excipient. Any pharmaceutical carrier, diluent or excipient known in the art is contemplated. A related aspect of the invention provides a kit comprising the pharmaceutical composition and a set of instructions for administering the composition to a subject in need. In each of these aspects of the invention, i.e., the pharmaceutical compositions and the kits, the heterologous peptide ligands (or binding pair members) and cell-surface binding partners described in the context of describing the recombinant HSV particles are contemplated.

[0020] Yet another aspect of the invention provides a method of targeting a recombinant HSV particle to a cell comprising (a) identifying a binding pair member, such as a ligand for a ligand binding partner, exhibited on the surface of a target cell; and (b) creating an HSV particle as described herein, wherein the ligand or, more generally, the binding pair member, binds to the binding partner exhibited on the surface of the target cell. In some preferred embodiments of this aspect of the invention, the altered viral surface protein is selected from the group consisting of gB and gC. In some embodiments, the alteration to gD reduces binding of gD to at least one cellular receptor for gD selected from the group consisting of HveA and Nectin-1. In preferred embodiments, the altered gD has a conservative substitution at position 34 of gD, such as a V34S substitution. A second fusion protein, joining the ligand (or binding pair member) and either of gB or gC, is also contemplated in some embodiments. Suitable

cells for targeting include any hyperproliferative cell, such as a cancer cell, including tumor cells (e.g., malignant glioma cells). Any of the heterologous peptide ligands (or binding pair members) and cell-surface binding partners described above in the context of describing the recombinant HSV particles is suitable for use in the method.

[0021] Another aspect of the invention is drawn to a method of imaging a cell comprising: (a) contacting the cell with a recombinant HSV particle as described above, the recombinant HSV particle further comprising a coding region for a marker protein; and (b) detecting the presence of the marker protein. Any type of cell exhibiting a cell-surface binding partner for a ligand (or binding pair member) fusible to HSV gD is suitable for use in this aspect of the invention, such as a cancer cell. Using cancer cells as an example, the method is useful provided that the binding partner is present at a higher number on the cancer cell as compared to a non-cancerous cell of the same type. Any known marker protein is useful in this aspect of the invention, e.g., a marker protein selected from the group consisting of thymidine kinase, green fluorescent protein, and luciferase. In preferred embodiments, the altered gD exhibits an amino acid substitution of V34S. Any of the heterologous peptide ligands (or binding pair members) and cell-surface binding partners described above in the context of describing the recombinant HSV particles is suitable for use in the method.

[0022] Another aspect of the invention provides a method of treating a cell-based disease comprising delivering a therapeutically effective amount of a recombinant HSV particle as described herein to a subject in need. A related aspect is the use of a recombinant HSV particle as described above in the preparation of a medicament for the treatment of a cell-based disease. A therapeutically effective amount of a recombinant HSV particle is that amount that produces the desired therapeutic effect, as would be understood and readily determinable by those of skill in the art. Any cell-based disease known or reasonably suspected to be amenable to treatment with a specifically targeted HSV is contemplated, e.g., a cell hyperproliferation disease such as cancer. Any of the heterologous peptide ligands (or binding pair members) and cell-surface binding partners described above in the context of describing the recombinant HSV particles is suitable for use in the method.

[0023] In a related aspect, the invention provides a method of ameliorating a symptom associated with a disease comprising administering a therapeutically effective amount of a recombinant HSV particle described above to a subject in need. Another aspect is drawn to the use of a recombinant HSV particle as described above in the preparation of a medicament for ameliorating a symptom associated with a disease in a subject in need. Again, any disease known or reasonably suspected to have a symptom amenable to application of a specifically targeted HSV is contemplated, including any disease characterized by hyperproliferative cells, such as cancer. Any of the heterologous peptide ligands (or binding pair members) and cell-surface binding partners described above in the context of describing the recombinant HSV particles is suitable for use in the method.

[0024] Another aspect of the invention is directed to a method of delivering a therapeutically useful peptide to a cell comprising: (a) inserting a therapeutically useful polynucleotide, such as an expression control element, an rDNA, or a coding region for a therapeutically useful peptide, into the DNA of a recombinant HSV particle as described above,

thereby producing a recombinant HSV clone; and (b) delivering a therapeutically effective amount of the recombinant HSV clone to the cell. In a related aspect, the invention provides for the use of a recombinant HSV clone comprising a recombinant HSV particle according to claim 1 in the preparation of a medicament for delivering a therapeutically useful peptide to a cell comprising inserting a coding region for a therapeutically useful peptide into the DNA of the recombinant HSV particle, thereby producing the recombinant HSV clone. Each of the method and use comprehends delivery of the recombinant HSV clone in vivo, ex vivo, or in vitro. Any of the heterologous peptide ligands (or binding pair members) and cell-surface binding partners described above in the context of describing the recombinant HSV particles is suitable for use in the method or use.

[0025] Another aspect of the invention provides a method of killing a target cell, comprising contacting the target cell with a recombinant HSV particle as described above. A related aspect is the use of a recombinant HSV particle as described above in the preparation of a medicament for killing a target cell by contacting the target cell with the recombinant HSV particle. In preferred embodiments of either the method or the use, the recombinant HSV particle has an altered gD in which the V34S substitution is found. Any of the heterologous peptide ligands (or binding pair members) and cell-surface binding partners described above in the context of describing the recombinant HSV particles is suitable for use in the method or use.

[0026] In each of the above-described aspects of the invention, it is preferred that gD, or a portion thereof, maintains its membrane fusion properties, but has reduced capacity to bind HveA and/or Nectin-1.

[0027] Other features and advantages of the invention will be better understood by reference to the brief description of the drawing and the description of the illustrative embodiments that follow.

BRIEF DESCRIPTION OF THE DRAWING

[0028] FIG. 1. Schematic representation of the HSV-1 (F) genome and gene manipulations in glycoprotein C (gC) (FIG. 1A), glycoprotein B (gB) (FIG. 1B), and glycoprotein D (gD) (FIG. 1C). Line 1, sequence arrangement of the HSV-1 genome. The rectangular boxes represent the inverted repeat sequences ab and b' a' flanking the unique long (U_L) sequence, and inverted repeat c'a' and ca flanking the unique short (U_S) sequence. Line 2, sequence arrangement of domains of the glycoprotein C; the signal peptide (SP) domain and heparan sulfate (HS)-binding domain of gC are highlighted. Line 3, human IL13 with signal peptide that replaced the N-terminal segment of 148 amino acids of gC. Line 4, sequence arrangement of the poly-lysine domain of gB. Line 5, schematic representation of a recombinant HSV-1(F) genome, in which the N-terminal domain of gC was replaced with IL13 and the polylysine domain (from codon 68 to codon 77) of gB was deleted. Line 6, sequence arrangement of glycoprotein J (gJ), glycoprotein D (gD), and glycoprotein I (gI) in U_S . Line 7, replacement of gD with the immediate early promoter of CMV in order to enable the expression of gI. Line 8, schematic representation of recombinant HSV-1(F) genome, in which the N-terminal domain of gC was replaced with IL13, the poly-lysine domain of gB was deleted, and IL13 was inserted after amino acid 24 of gD (FIG. 1D). Line 9, a

polylinker XhoI-BglIII-EcoRI-KpnI was inserted after amino acid 24 of gD, with IL13 inserted into the XhoI and KpnI sites of gD.

[0029] FIG. 2. Amino acid sequence alignment of IL13-gC, IL13-gD junction sequence, and the HS binding domain of gB. FIG. 2A. The amino-terminal sequence of IL13-gC chimeric protein (SEQ ID NO:22). The sequences upstream and downstream of the HS binding site portion are shown. IL13 was inserted between the two restriction enzyme sites that are underlined. FIG. 2B. The domain of the gB open reading frame (i.e., ORF) from which the poly lysine [poly(K)] sequence was deleted (SEQ ID NO:23). The underlined sequences (codons 68-77) were not present in gB amplified from R5107. FIG. 2C. The amino-terminal sequence of IL13-gD (SEQ ID NO:24). The first underlined sequence identifies the gD signal peptide. IL13 (bracketed by arrows) was inserted between residues 24 and 25 (underlined) of gD, between the XhoI and KpnI restriction enzyme sites.

[0030] FIG. 3. Verification of R5111 viral DNA by PCR. Photographs of electrophoretically separated PCR products amplified directly from the plaques picked from Vero (FIG. 3A) and HEp-2 (FIG. 3B) cells. Viral DNAs were extracted as described in Example 1 and subjected to PCR with "IL13" primers from the IL13 ORF and IL13-gD primers, which bracketed IL13 and the gD ectodomain.

[0031] FIG. 4. Photograph of electrophoretically separated proteins from lysates of cells infected with R5111 reacted with antibody to gC, gD or IL13. HEp-2 cells grown in 25 cm² flasks were exposed to 10 PFU of HSV-1 or R5111 per cell. The cells were harvested 24 hours after infection, solubilized, subjected to electrophoresis in 10% denaturing polyacrylamide gels, electrically transferred onto a nitrocellulose sheet, and exposed to a monoclonal antibody against gD (FIG. 4A), gC (FIG. 4B) or IL13 (FIG. 4C), respectively. The protein bands corresponding to the gC, IL13-gC fusion protein, gD and the IL13-gD fusion protein are indicated. IL13-gC was the same size as native gC, as expected.

[0032] FIG. 5. HA-tagged IL13R α 2 expression from individual clones of stable transfectants of the J1.1 cell line. The individual clones were amplified as described in Example 1. Cells were harvested from 25 cm² flasks, solubilized, and subjected to electrophoresis in 12% denaturing polyacrylamide gels, electrically transferred onto a nitrocellulose sheet, and exposed to a polyclonal antibody to HA tag.

[0033] FIG. 6. Diagram of the pgD-vector.

[0034] FIG. 7. Schematic representation of the HSV-1 (F) genome and genetic structure of R5141 and R5144.

DETAILED DESCRIPTION

[0035] The invention provides benefits that will improve the health and well-being of animals such as man by providing a targeted approach to the treatment of a variety of conditions and diseases that currently impair health, resulting in significant economic burdens using conventional treatments. In providing modified viral particles having controllable targeting capacities, the diagnostic and therapeutic benefit of the viruses themselves can be delivered with greater precision to particular cells. Additionally, these viral particles can be used as targeting vehicles for the delivery of a wide variety of therapeutic and diagnostic biomolecules, such as polynucleotides encoding therapeutic or diagnostic peptides.

[0036] Beyond the modified viral particles, the invention provides methods for making such therapeutic and diagnostic agents as well as methods for using the agents to diagnose or

treat a variety of diseases and conditions, such as tumorigenic disease (e.g., gliomas). To facilitate an understanding of the invention and all of its aspects, illustrative embodiments are described below. The descriptions of these illustrative embodiments are not meant to limit the invention to the embodiments disclosed herein. In light of the description, one of skill in the art will understand that many changes and modifications can be made to the illustrative embodiments and still remain within the invention. The illustrative embodiments are disclosed using as an exemplary virus a member of the Herpesviridae family of viruses, herpes simplex virus (HSV).

[0037] As noted above, HSV-1 and HSV-2 are members of the family of viruses known as the Herpesviridae, whose structures are well known in the art. The targeting methods of the invention are applicable to any member of the Herpesviridae and are not limited to the exemplary embodiments described in the examples. Furthermore, a large number of recombinant HSV viruses are known in the art. Such viruses may contain one or more heterologous genes. Also, such viruses may contain one or more mutated HSV genes, for example, mutations that render the virus replication-deficient or affect the virulence of the virus in one or more cell types.

[0038] Examples of recombinant HSV containing a heterologous gene and methods of making and using such viruses are described in U.S. Pat. No. 5,599,691 (incorporated herein by reference in its entirety). Preferred heterologous genes include genes encoding marker proteins. Marker proteins, such as green fluorescent protein, luciferase, and beta-galactosidase, allow detection of cells expressing the protein. In other embodiments, the heterologous gene encodes an enzyme that activates a prodrug thereby killing adjacent uninfected cells. In yet other embodiments, the heterologous gene encodes a protein that affects the immune response, such as interleukin 12 (IL-12). Such proteins that activate the immune response against a tumor are particularly useful.

[0039] In one aspect, the invention relates to altering the surface of an HSV particle to target the virus to a specific receptor. By creating a fusion protein comprising a portion of gD and a ligand (or binding pair member), the virus is targeted to a cell having a cell surface receptor that binds the ligand (or binding pair member). In preferred embodiments, one or more HSV surface proteins, such as gB (SEQ ID NOs:27 and 28), gC (SEQ ID NOs:29 and 30), or gD (SEQ ID NOs:25 and 26), are altered to reduce binding to natural HSV receptors.

[0040] "Alterations" of the surface of an HSV particle or HSV surface protein include insertions, deletions, and/or substitutions of one or more amino acid residues. One type of alteration is an insertion, which involves the incorporation of one or more amino acids into a known peptide, polypeptide or protein structure. For ease of exposition, alterations will be further described using a protein exemplar. Insertions may be located at either or both termini of the protein, or may be positioned within internal regions of known proteins, which yield proteins such as fusion proteins and proteins having amino acid tags or labels.

[0041] Another type of alteration is a deletion, wherein one or more amino acid residues in a protein are removed. Deletions can be effected at one or both termini of the protein, or with removal of one or more residues within the amino acid sequence. Deletion alterations, therefore, include all fragments of a protein described herein.

[0042] Yet another type of alteration is a substitution, which includes proteins wherein one or more amino acid

residues are removed and replaced with alternative residues. In one aspect, the substitutions are conservative in nature; however, the invention embraces substitutions that are also non-conservative. Conservative substitutions for this purpose may be defined as set out in Tables A or B, below.

[0043] Amino acids can be classified according to physical properties and contribution to secondary and tertiary protein structure. A conservative substitution is recognized in the art as a substitution of one amino acid for another amino acid that has similar properties. Exemplary conservative substitutions are set out in Table A as described in Lehninger, [Biochemistry, 2nd Edition; Worth Publishers, Inc. New York (1975), pp. 71-77] and set out immediately below.

TABLE A

Conservative Substitutions I	
SIDE CHAIN CHARACTERISTIC	AMINO ACID
Non-polar (hydrophobic):	
A. Aliphatic	A L I V P
B. Aromatic	F W
C. Sulfur-containing	M
D. Borderline	G
Uncharged-polar:	
A. Hydroxyl	S T Y
B. Amides	N Q
C. Sulfhydryl	C
D. Borderline	G
Positively charged (basic)	K R H
Negatively charged (acidic)	D E

[0044] Alternative, exemplary conservative substitutions are set out in Table B, immediately below.

TABLE B

Conservative Substitutions II	
ORIGINAL RESIDUE	EXEMPLARY SUBSTITUTION
Ala (A)	Val, Leu, Ile
Arg (R)	Lys, Gln, Asn
Asn (N)	Gln, His, Lys, Arg
Asp (D)	Glu
Cys (C)	Ser
Gln (Q)	Asn
Glu (E)	Asp
His (H)	Asn, Gln, Lys, Arg
Ile (I)	Leu, Val, Met, Ala, Phe,
Leu (L)	Ile, Val, Met, Ala, Phe
Lys (K)	Arg, Gln, Asn
Met (M)	Leu, Phe, Ile
Phe (F)	Leu, Val, Ile, Ala
Pro (P)	thy
Ser (S)	Thr
Thr (T)	Ser
Trp (W)	Tyr
Tyr (Y)	Trp, Phe, Thr, Ser
Val (V)	Ile, Leu, Met, Phe, Ala

[0045] The binding site of HveA has been reported to be at the amino terminal domain of gD (Carfi A., et al., 2001) The precise binding sites of gD for Nectin 1 are not known, although it has previously been reported to involve gD amino acids 38 and 221 (Manoj S., et al., 2004; Zago A., et al., 2004; Connolly S A., 2005). Accordingly, in one aspect the invention relates to amino acid alterations in the N-terminal region of gD such that the ability of gD to bind HveA or Nectin1 is

reduced or eliminated A “natural receptor” as used herein is a cell surface molecule that interacts with wild-type HSV in the absence of human intervention. For example, gB and gC of HSV-1 interact with heparan sulfate proteoglycans in a natural infection. In preferred embodiments, gB and/or gC are altered to reduce or eliminate binding to heparan sulfate proteoglycans. As another example, gD is known to bind to several receptors, including HveA and Nectin-1, in a natural infection. In preferred embodiments, gD is altered to reduce or eliminate binding to HveA and/or Nectin-1.

[0046] Receptor-Ligands

[0047] As used herein, “receptor” and “ligand” refer to two members of a specific binding pair and, hence, are binding partners. A receptor is that member of the pair that is found localized on the surface of the cell; the ligand is the member of the pair that is found on the surface of HSV. Thus, in certain embodiments, the “ligand” may actually be what the art recognizes as a receptor outside the context of the invention and the “receptor” may be its respective ligand. More generally, the invention comprehends an HSV exhibiting a member of a binding pair, or a fragment thereof that retains the capacity to specifically bind the other member of the binding pair, on its surface and the other member of that binding pair, or a fragment thereof that retains the capacity to specifically bind its partner, is present on the surface of a target cell.

[0048] One advantage of the invention is the ability to tailor HSV to target a specific receptor while maintaining infectivity of the virus. In an exemplary embodiment, an HSV particle contains a fusion protein comprising a portion of gD and the cytokine IL-13. Such a virus is able to infect cells expressing the receptor IL-13R α 2. Because IL-13R α 2 is expressed on the surface of cells of malignant gliomas, HSV containing the gD/IL-13 fusion protein are effectively targeted to such cells. Ligands that bind to receptors which are overexpressed or differentially expressed on either tumor cells or cells associated with tumor growth (e.g., neovasculature) are particularly preferred. Examples include the $\alpha_v\beta_3$ - $\alpha_v\beta_5$ integrins, which are overexpressed in tumor neovasculature; epidermal growth factor receptor (EGFR), which is overexpressed in head, neck, lung, colon, breast, and brain cancer cells; HER-2/Neu, which is overexpressed in breast cancer cells; MUC-1, which is overexpressed in breast, lung, and pancreas cancer cells; and prostate-specific membrane antigen, which is overexpressed in prostate cancer cells. In certain embodiments, the ligand is a single-chain antibody that binds to its cognate specific binding pair member, herein referred to as a receptor. Single-chain antibodies have been shown to be effective in targeting applications, as evidenced by their ability to target retroviruses to specific receptors.

[0049] Essentially any two binding pair members or partners may be used as receptor-ligands in the invention. However, it is contemplated that certain factors, such as the distance from the binding site on the receptor to the membrane, or the conformation of the ligand when fused to gD, may affect the efficiency of recombinant HSV fusion to the cell membrane. Therefore, screens for effective receptor-ligand pairs are contemplated, using no more than routine procedures known in the art. Additional screens, conventional in nature, may be used to optimize constructs. One routine method of screening is to follow the protocol provided in the example for candidate receptor/ligand pairs, using IL-13R/IL-13 as a control receptor/ligand pair.

[0050] Alternatively, one may use a membrane fusion assay as described in Turner et al., 1998, incorporated herein by

reference in its entirety. In the Turner assay, cells transfected with construct(s) encoding gB, gH, gL, and the gD/ligand fusion protein, and cells expressing the receptor, are co-cultured and the cells are examined for membrane fusion. Membrane fusion between gD/ligand-expressing cells and receptor-expressing cells indicates that the candidate receptor-ligand pair (the ligand being a gD/ligand fusion protein) is functional. Constructs encoding functional gD/ligand proteins can then be used to create recombinant HSV that are targeted to cells expressing the receptor.

[0051] Cell Targeting

[0052] Evident from the preceding discussion, another aspect of the invention is the targeting of a recombinant HSV to a cell having a specific receptor on its surface. In preferred embodiments, a recombinant HSV is designed to comprise a ligand that interacts with a receptor known to be expressed on a cell of interest. The cell of interest is then infected with recombinant HSV. Such targeting methods may be used for a variety of purposes.

[0053] In one aspect, a recombinant HSV is used to introduce a heterologous gene into a cell that expresses the receptor. In preferred embodiments, the cell is not infected by, or is poorly infected by, wild-type HSV. Thus, in certain embodiments, the invention provides a vector for transforming a cell of interest with a heterologous gene.

[0054] Further, a cell can be rendered a target of a recombinant HSV of the invention. The cell can be rendered a target by transforming the cell to express one member of a binding pair, e.g., a receptor capable of specifically binding a ligand expressed on a recombinant HSV. For example, as described in Example 2, the J1.1 cell line, which was resistant to infection by a recombinant HSV expressing an IL-13 ligand, was rendered susceptible to infection by transforming the cell line with a vector encoding IL12R α 2 to produce the cell line J13R.

[0055] Generally, the targeted HSV according to the invention exhibit one member of a binding pair, with the other member of that pair found on the surface of a target cell. In some embodiments of the invention, targeting is achieved with a ligand-receptor binding pair, with the ligand exhibited on the targeted HSV and the cognate receptor found on the surface of the target cell, as described above. Although the invention comprehends embodiments involving binding pairs that do not exhibit a ligand-receptor relationship (e.g., biotin-avidin) and embodiments in which the receptor is exhibited by the targeted HSV (the “receptor” defined above as a “ligand” using an alternative definition of “ligand”) while the cognate ligand is found on the target cell (the “ligand” defined above as a “receptor” using an alternative definition of “receptor”), embodiments in which the targeted HSV exhibits a ligand and the target cell presents the cognate receptor on its surface is used as an illustrative embodiment to reveal the versatility of the invention and to disclose the full scope thereof. For example, several ligands have been used for receptor-mediated polynucleotide transfer. Some ligands that have been characterized are asialoorosomuroid (ASOR) and transferrin (Wagner et al., Proc. Natl. Acad. Sci. USA, 87(9): 3410-3414, 1990). A synthetic neoglycoprotein, which recognizes the same receptor as ASOR, has also been used in a polynucleotide delivery vehicle (Ferkol et al., FASEB J., 7:1081-1091, 1993; Perales et al., Proc. Natl. Acad. Sci., USA 91:4086-4090, 1994) and epidermal growth factor (EGF) has further been used to deliver polynucleotides to squamous carcinoma cells (Myers, EPO 0273085). Each of these spe-

cific approaches, and other approaches known in the art to achieve some selectivity in DNA delivery, or targeting, are amenable to use in the compositions and methods of the invention and are contemplated as embodiments of the invention.

[0056] For embodiments in which a targeted HSV harboring a coding region, e.g., a therapeutic coding region or gene, is delivered to a target cell, the nucleic acid encoding the therapeutic gene product may ultimately be positioned and expressed at different sites. In certain embodiments, the nucleic acid encoding the therapeutic polynucleotide may be stably integrated into the genome of the cell. This integration may place the gene in its native location and orientation via homologous recombination (gene replacement) or it may be integrated in a random, non-specific location (gene augmentation). In yet further embodiments, the nucleic acid may be stably maintained in the cell as a separate, episomal segment of DNA. Such nucleic acid segments or episomes encode functions sufficient to permit maintenance and replication independent of or in synchronization with the host cell cycle. How the expression construct is delivered to a cell and where in the cell the nucleic acid remains is dependent on the type of expression construct employed, as would be understood in the art.

[0057] It is envisioned that promoters subject to cell cycle regulation will be useful in the present invention. For example, in a bicistronic HSV vector designed to treat a disease, disorder or condition by killing a target cell, use of a strong CMV promoter to drive expression of a first gene, such as p16, that arrests a cell in the G1 phase is accompanied by expression of a second gene, such as p53, under the control of a promoter that is active in the G1 phase of the cell cycle, thus providing a dual-gene approach to ensure that the target cell undergoes apoptosis. Other promoters, such as those of various cyclins, PCNA, galectin-3, E2F1, p53, BRCA1, and, indeed, any suitable promoter or expression element known in the art, could be used.

[0058] In embodiments of the invention designed to treat diseases, disorders, or conditions associated with unwanted or excessive cell proliferation, such as cancer or restenosis, HSV is targeted to proliferating cells thereby killing the cells. Because HSV is lethal to infected cells, expression of a heterologous gene is not required. However, in embodiments wherein the lethality of HSV is attenuated, an HSV harboring a gene that is lethal to the infected cell or that prevents proliferation of the infected cell may be used to target a cell.

[0059] Alternatively, HSV targeted to specific surface markers can be used to visualize the distribution of tumor cells in tissues. This diagnostic tool had been unavailable because of the indiscriminate binding of HSV to cells. Modification of HSV by eliminating (ablating) or reducing the indiscriminate binding of HSV to heparan sulfate proteoglycans without deleteriously affecting the capacity of such HSV to replicate in both dividing and non-dividing cells makes possible the use of these modified viral forms to visualize the distribution of tumor cells.

[0060] In one preferred method for visualizing the distribution of tumor cells, radioactive visualization is achieved by viral thymidine kinase (TK)-dependent incorporation of a radioactive precursor. Methods of molecular imaging of gene expression are well known in the art. Methods often use highly sensitive detection techniques such as positron emission tomography (PET) or single-photon emission-computed tomography (SPECT). In one embodiment, TK expression is

measured using a gancyclovir analog, such as 9-(3-[¹⁸F] fluoro-1-hydroxy-2-propoxy)methyl)guanine, as the tracer or marker (Vries et al., 2002). For a review of imaging TK gene expression using PET or SPECT, see Sharma et al., 2002 or Vries et al., 2002.

[0061] A second preferred imaging method is to fuse a non-critical tegument protein (e.g. U_S11, which is present in nearly 2000 copies per virus particle) to a marker protein, such as green fluorescent protein, which is capable of being visualized *in vivo*. Alternatively, a non-critical protein can be fused to a luciferase and the presence of the luciferase visualized with a luminescent or chromatic luciferase substrate. Although a marker protein can be fused to essentially any viral structural protein, preferred viral proteins include gC, gE, gI, gG, gJ, gK, gN, U_L11, U_L13, U_L14, U_L21, U_L41, U_L35, U_L45, U_L46, U_L47, U_L51, U_L55, U_L56, U_S10, and U_S11. The marker protein also may be fused to thymidine kinase (Soling et al., 2002).

[0062] Library Screening

[0063] As noted above, HSV comprising a gD/ligand fusion protein can bind and infect cells expressing a receptor to the ligand. In one embodiment, a cell line expressing a receptor is used in screening for ligands of the receptor. cDNA from a cDNA library is cloned into a vector encoding a portion of gD to produce a gD/cDNA-encoded fusion protein. The resulting vectors are then screened for membrane fusion using the assay of Turner et al. described above or by creating recombinant HSV expressing the gD/cDNA-encoded fusion protein and screening the viruses for the ability to infect receptor-expressing cells. Such methods may be used, e.g., to identify a ligand to an orphan receptor.

[0064] In other embodiments, mutations in, or variants of, the receptor or ligand are screened to determine whether the mutants or variants maintain the ability to interact with the respective partner. Such methods may be useful in determining the specific residues important in receptor-ligand interaction.

[0065] Therapeutic Methods

[0066] Another aspect of the invention is the use of the targeted HSV in therapeutic methods. By altering the cell-binding and infectivity properties of the virus, many routes and methods of administration become viable. For example, non-targeted HSV will bind indiscriminately to a variety of cells. Because of this property, large virus numbers are used and intravenous administration is generally not effective. However, by targeting the virus, one may lower the viral load (i.e., quantity of virus), yet maintain or increase efficacy. Furthermore, the targeted HSV can be administered intravenously and produce therapeutic effects.

[0067] Therapeutic methods of the invention include those methods wherein an HSV is targeted to a receptor of a cell that contributes to, or is the basis of, a disease or disorder. These targeted HSV can either exploit the therapeutic properties of HSV itself (e.g., the lethality of HSV to infected cells) or the targeted HSV can serve as a vector for the targeted delivery of at least one therapeutic polynucleotide, such as an expressible polynucleotide comprising a coding region. For example, in methods wherein the targeted HSV contains one or more gene products that render the virus toxic to the cell or that prevent or inhibit cell proliferation, a preferred receptor is overexpressed or selectively expressed on harmful or undesirable cells, such as cancer cells. In other embodiments, the targeted HSV encodes a gene product that provides a desired function

or activity in the targeted cell, e.g., when a cell has one or more genetic defects preventing the cell from functioning properly.

[0068] Additionally, it is contemplated that a therapeutic polynucleotide (e.g., gene or coding region) of a targeted HSV may be engineered to be under the expression control of a cell- or tissue-specific expression control element, e.g., a promoter. In such embodiments, the targeted HSV provide a further enhancement to the selective treatment of a suitable disorder, disease or condition. The targeted HSV is specific for a binding partner located on the surface of those cells for which treatment is intended, and expression of the therapeutic coding region or gene borne by the targeted HSV is limited to particular cells or tissues.

[0069] As HSV has been engineered to overcome the barriers to vector-based therapies, the choice of recombinant polynucleotide to be inserted into the vector has widened to the point where a wide variety of diseases, disorders and conditions are amenable to treatment with targeted HSV. A number of diseases are amenable to polynucleotide-based therapy using HSV (see, e.g., Kennedy, et al., *Brain*, 120, 1245-1259, 1997, incorporated by reference herein in its entirety). Though most attention has focused on cancers, there has been success in treating Parkinson's disease by expressing tyrosine hydroxylase in striatal cells, thus restoring L-dopa-induced nerve repair following axotomy of the superior cervical ganglion. Injection of a vector expressing nerve growth factor resulted in restored levels of tyrosine hydroxylase. More generally, HSV can now be used in polynucleotide-based therapy to replace missing or defective coding regions in the target cells. In the event of an inherited single-gene disorder (such as Lesch-Nyhan syndrome) where the complete DNA sequence, cause, and effect of the disorder are known, a single polynucleotide replacement mediated by targeted HSV is appropriate and contemplated. Another strategy amenable to the use of targeted HSV is the enhancement of endogenous expression levels of a gene product, e.g., a growth factor or enzyme. Yet another strategy for using targeted HSV is HSV-directed enzyme pro-drug therapy. The delivery of a drug-sensitivity gene would be beneficial in the treatment of, e.g., a malignant brain tumor, making the tumor more susceptible to conventional anti-cancer agents.

[0070] In other embodiments, the targeted HSV of the invention provide for vector-mediated delivery of anti-sense oligodeoxyribonucleotides (oligonucleotides). The oligonucleotides, short segments of DNA (e.g., 2-100 nucleotides in length), are delivered to target cells and therein bind to complementary mRNA, thus blocking the expression of specific genes within the target cells. The encoded protein fail to be synthesized, as the mRNA is not recognized by the translational components of the cell. In preferred embodiments, a deleterious gene is targeted.

[0071] In yet other embodiments, targeted HSV are used to deliver polynucleotides, e.g., DNAs encoding gene products, that can recruit or enhance an immune system response, thereby bringing a subject's or patient's own immune system to bear in the treatment of a disease, disorder or condition known in the art to be amenable to immune system activity. For example, an increase in cellular antigen expression of tumor cells, mediated by delivery of an expressible coding region for the antigen by a targeted HSV, would enhance the immune response and increase the susceptibility of such tumor cells to host cytotoxic immunity.

[0072] In some embodiments, a targeted HSV composition of the invention is delivered to a patient at or around the site of a tumor, which is a very efficient method for counteracting clinical disease. Alternatively, systemic delivery of targeted HSV compositions may be appropriate in other circumstances, for example, where extensive metastasis has occurred, or where inaccessible tumors are encountered.

[0073] It is contemplated that in certain embodiments of the invention a protein that acts as an angiogenesis inhibitor is targeted to a tumor. Also, an angiogenesis inhibitor agent may be administered in combination with a targeted HSV of the invention. These agents include, for example, Marimastat (British Biotech, Annapolis MD; indicated for non-small cell lung, small cell lung and breast cancers); AG3340 (Agouron, LaJolla, Calif.; for glioblastoma multiforme); COL-3 (Collagenex, Newtown Pa.; for brain tumors); Neovastat (Aeterna, Quebec, Canada; for kidney and non-small cell lung cancer) BMS-275291 (Bristol-Myers Squibb, Wallingford Conn.; for metastatic non-small cell lung cancer); Thalidomide (Celgen; for melanoma, head and neck cancer, ovarian, and metastatic prostate cancers; Kaposi's sarcoma; recurrent or metastatic colorectal cancer (with adjuvants); gynecologic sarcomas, liver cancer; multiple myeloma; CLL, recurrent or progressive brain cancer, multiple myeloma, and non-small cell lung, nonmetastatic prostate, refractory multiple myeloma, and renal cancer); Squalamine (Magainin Pharmaceuticals Plymouth Meeting, PA; non-small cell lung cancer and ovarian cancer); Endostatin (EntreMED, Rockville, Md.; for solid tumors); SU5416 (Sugen, San Francisco, Calif.; recurrent head and neck, advanced solid tumors, stage IIIB or IV breast cancer; recurrent or progressive brain (pediatric) cancer; ovarian cancer, AML (acute myeloid leukemia); glioma, advanced malignancies, advanced colorectal cancer, von-Hippel Lindau disease, advanced soft tissue cancer; prostate cancer, colorectal cancer, metastatic melanoma, multiple myeloma, malignant mesothelioma; metastatic renal, advanced or recurrent head and neck cancer, metastatic colorectal cancer); SU6668 (Sugen San Francisco, Calif.; advanced tumors); interferon- α ; Anti-VEGF antibody (National Cancer Institute, Bethesda Md.; Genentech San Francisco, Calif., for refractory solid tumors; metastatic renal cell cancer, in untreated advanced colorectal cancer; EMD121974 (Merck KGaA, Darmstadt, Germany, for HIV-related Kaposi's sarcoma, and progressive or recurrent Anaplastic Glioma); Interleukin 12 (Genetics Institute, Cambridge, Mass., for Kaposi's sarcoma) and IM862 (Cytran, Kirkland, Wash., for ovarian cancer, untreated metastatic cancers of colon and rectal origin, and Kaposi's sarcoma). The parenthetical information following the agents indicates the cancers against which the agents are being used in these trials. It is contemplated that any of these disorders may be treated with the targeted HSV compositions of the invention, either alone or in combination with the agents listed.

[0074] In order to prepare a therapeutic composition for clinical use, it will be necessary to prepare the therapeutic composition as a pharmaceutical composition, i.e., in a form appropriate for in vivo applications. Generally, this will entail preparing compositions that are essentially free of pyrogens, as well as other impurities that could be harmful to humans or other vertebrates.

[0075] Generally, appropriate salts and buffers are included to render delivery vectors stable and to allow for uptake by target cells. Aqueous compositions of the invention comprise an effective amount of the targeted HSV, dissolved or dis-

persed in a pharmaceutically acceptable carrier or aqueous medium. Such compositions also are referred to as inocula. The phrase “pharmaceutically acceptable” or “pharmacologically acceptable” refer to molecular entities and compositions that do not produce adverse, allergic, or other untoward reactions when administered to an animal or a human. As used herein, “pharmaceutically acceptable carriers” includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Unless a conventional medium or agent is incompatible with either the vectors of the invention or the intended subject receiving treatment, its use in therapeutic compositions is contemplated. Supplementary active or inert ingredients also can be incorporated into the compositions.

[0076] The active compositions of the invention include standard pharmaceutical preparations. Administration of these compositions according to the invention is by any known route, provided that the target tissue is accessible via that route. The pharmaceutical compositions may be introduced into the subject by any conventional method, e.g., by intravenous, intradermal, intramuscular, intramammary, intraperitoneal, intrathecal, retrobulbar, intravesicular, intrapulmonary (e.g., term release); sublingual, nasal, anal, vaginal, or transdermal delivery, or by surgical implantation at a particular site. The treatment may consist of a single dose or a plurality of doses over a period of time.

[0077] Upon formulation, solutions are administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. Appropriate dosages may be ascertained through the use of established routine assays. As studies are conducted, further information will emerge regarding optimal dosage levels and duration of treatment for specific diseases, disorders, and conditions.

[0078] In preferred embodiments, the unit dose may be calculated in terms of the dose of viral particles being administered. Viral doses are defined as a particular number of virus particles or plaque forming units (pfu). Particular unit doses include 10^3 , 10^4 , 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} , 10^{11} , 10^{12} , 10^{13} or 10^{14} pfu. Particle doses may be somewhat higher (10- to 100-fold) due to the presence of infection-defective particles, which is determinable by routine assays known in the art.

[0079] The pharmaceutical compositions and treatment methods of the invention are useful in the fields of human medicine and veterinary medicine. Thus, the subject to be treated may be a vertebrate, e.g., a mammal, preferably human. For veterinary purposes, subjects include, for example, farm animals such as cows, sheep, pigs, horses and goats, companion animals such as dogs and cats, exotic and/or zoo animals, laboratory animals including mice, rats, rabbits, guinea pigs and hamsters; and poultry such as chickens, turkey, ducks and geese.

[0080] In some embodiments of the invention, it is contemplated that the targeted HSV is administered in conjunction with chemo- or radiotherapeutic intervention, immunotherapy, or with any other therapy conventionally employed in the treatment of cancer.

[0081] To kill cells, inhibit cell growth, inhibit metastasis, inhibit angiogenesis or otherwise reverse or reduce malignant phenotypes using the methods and compositions of the invention, one contacts a “target” cell, a tumor, or its vasculature with a targeted HSV composition and at least one other agent. The components of these compositions are provided in a

combined amount effective to kill or inhibit proliferation of cancer cells. This process may involve contacting the cells with the targeted HSV composition and the agent(s) or factor (s) at the same time. This may be achieved by contacting the subject organism, or cell of interest, with a single composition or pharmacological formulation that includes both agents, or by contacting the cell with two distinct compositions or formulations, at the same or different times, wherein one composition includes a targeted HSV composition of the invention and the other composition includes the second agent.

[0082] Another aspect of the invention provides diagnostic methods that involve imaging a tumor or diseased tissue using a targeted HSV. Such methods are useful in diagnosing a patient with a disease, disorder, or condition that is indicated by the presence of a receptor on the surface of a cell. Diagnostic imaging methods are discussed above.

[0083] Kits

[0084] Kits according to the invention may include recombinant viruses of the invention or may include vectors for producing such recombinant viruses. A vector for producing a recombinant virus of the invention may encode the gD/ligand fusion protein or may be designed to facilitate cloning of a ligand to produce a gD/ligand fusion protein (e.g., a vector containing a multiple cloning site within the gD coding region that facilitates in-frame insertions).

[0085] Other components that can be included in a kit of the invention include a receptor-expressing cell line (useful as a control), a nucleic acid molecule for expressing the receptor in a particular cell type, and instructions for using the kit to effect diagnostic analyses or therapeutic treatments. In certain embodiments, a therapeutic kit will further contain a component for bringing about a therapeutic effect, such as a prodrug or a toxic compound. In other embodiments, a diagnostic kit will contain a compound useful in imaging methods, such as a chromophore or fluorophore, or an antibody for detecting infected cells.

[0086] Having provided a general description of the various aspects of the invention, the following disclosure provides examples illustrative of the invention, wherein Example 1 describes construction of a targeted HSV, Example 2 illustrates the construction of a cell line expressing a targeted HSV, and Example 3 describes the controlled infection of a desired cell by a targeted HSV.

Example 1

Construction of HSV Targeting Vector R5111

[0087] A targeted HSV was constructed to target the recombinant virus to cells of malignant gliomas. The target for entry of the virus into such cells is the IL13R α 2 receptor known to be present in malignant gliomas. Unlike the more prevalent IL13R α 1 receptor, the IL13R α 2 receptor has a shorter cytoplasmic domain and does not interact with IL4, of which IL13 is a close relative. In general, the construction of the targeted HSV involved mutagenizing gB and gC to preclude their interaction with heparan sulfate. Also, IL13 was inserted into gD at amino acid 24 thereby disrupting the gD binding site for HveA. The resulting IL13-gD chimeric virus can use IL13R α 2 for entry into cells carrying that receptor.

[0088] More specifically, the targeted HSV R5111 was constructed in several steps depicted in the four panels of FIG. 1 and detailed below.

[0089] (i) Substitution of Amino Terminal Domain of gC with IL13 Fused to the Signal Sequence of gC.

[0090] FIG. 1A, lines 1-3 schematically depicts a cDNA consisting of the IL13 coding sequence fused at its amino terminus to its signal sequence. The complete cDNA of IL 13, with the N-terminal signal peptide coding region, was amplified using the PCR primer elongation method. The primers were as follows:

pIL13F1,

(SEQ ID NO: 1)
CATTGCTCTCACTTGCCTTGGCGGCTTGCCTCCCCAGGCCCTGTGC
CTCCCTCTACAGC;

pIL13F2,

(SEQ ID NO: 2)
GCAGCTAGCCTCATGGCGCTTTTGTGACCACGGTCATTGCTCTCAC
TTGCCTTGGCGGC;

and pIL13REcoRI,

(SEQ ID NO: 3)
GAGCTCGGATCCTGAATTCAACCGTCCCTC.

[0091] First-round PCR used pIL13F1 and pIL13REcoRI as primers, with pRB5830 (containing the IL13 coding region) as the template. The PCR reaction mixture was then diluted 10-fold and 1 μ l of the diluted reaction mixture was used as template for the second round of PCR amplifications with pIL13F2 and pIL13REcoRI as the primer set. The PCR product was gel-purified, digested with NheI/EcoRI, and ligated into pBluescript II KS(+) at XbaI/EcoRI sites to generate pRB5832. To construct the transfer plasmid pRB5835, a 4.8-kbp HindIII/SacI fragment containing the HSV-1 gC coding region was released from cosmid pBC1007 and inserted into pBluescript II KS(+) to generate pRB5833. pRB5833 was cleaved with NheI and EcoRI and the N-terminal 148 residues of gC were replaced with the gC-signal/IL13 chimeric sequence (pRB5834). The insert in pRB5834 was released by XhoI/SacI digestion and ligated into pKO5Y at the same sites to generate pRB5835.

[0092] The recombinant virus R5107 (FIG. 1A, line 1) carrying the IL13-gC chimera was generated with the aid of the BAC-HSV system. RR1 competent cells that harbored bacterial artificial chromosome (BAC)-HSV bacmids were transformed with the transfer plasmid pRB5835 by electroporation. After incubation for 1 hour at 30° C. in LB broth, the transformed bacteria were plated on pre-warmed Zeocine (Zeo) plus chloramphenicol (Cm) (20 μ g/ml of each) plates and incubated overnight at 43° C. for integration. The next day, six colonies were picked and each was separately diluted in 1 ml LB. Five pi of the diluted bacteria were then plated on Cm/10% sucrose (Suc) plates, and incubated at 30° C. overnight. To further confirm the loss of the replacement vector, 24 Cm/Suc-resistant colonies (four colonies from each plate) were restreaked in duplicate on Cm LB and Zeo LB plates, respectively. The Suc^r/Cm^r/Zeo^r colonies were further screened by PCR (95° C., 4 minutes for cycle 1; then 35 cycles of 94° C., 1 minute; 60° C., 1 minute; and 72° C., 1 minute). The primers were:

[0093] pgC-F,

(SEQ ID NO: 4)
GACACGGGCTACCCCTCACTATCGAGGGC;

from nt 96158 to 96185 in HSV-1 strain 17),

[0094] and pgC-R,

(SEQ ID NO: 5)
GGTGATGTTTCGTACGACCTCCTCTAGGTC;

from nt 96859 to 96830 in HSV-1 strain 17).

[0095] The DNA fragment amplified from PCR-positive clones (FIG. 2B) was sequenced to further confirm the integration of IL13 in the correct open reading frame (ORF) of gC. To verify the viability of the recombinant (R5607), the recombinant BAC-HSV DNA was prepared as described elsewhere (Ye et al., 2000) and transfected into rabbit skin cells by Lipofectamine reagent (Life Technologies, Grand Island, N.Y.). The resultant virus, R5607, was stored at -80° C.

[0096] (ii) Deletion of the Polylysine Track from gB, FIG. 1 Panel B.

[0097] To make a transfer plasmid for the deletion of the gB heparan sulfate binding domain (polylysine), a 4.76 kbp BstEII fragment (from nt 53164 to 57923 of HSV-1) containing the U_L27 (gB) ORF released from cosmid BC1014 was blunt-ended and cloned into pBluescript II KS (+) at a SmaI site to generate pRB5846. To construct pRB5847, from which the 10-amino-acid polylysine domain of gB was deleted, two fragments flanking the polylysine domain were amplified by PCR from pRB5846. The primer sets were:

pgB1BamHI:

(SEQ ID NO: 6)
GTTCTTCTTCGGTTTCGGATCCCCCG;

pgB2BspEI:

(SEQ ID NO: 7)
CGGCATTTCCGGAATAACGCCCACTC;

and pgB3BamHI:

(SEQ ID NO: 8)
CAGAAAACCGGATCCCCCAAAGCCGCC;

pgB4BsiWI:

(SEQ ID NO: 9)
GCCAACACAAACTCGTCGTACGGGTAC.

[0098] PCR amplified fragments were then cut with BspEI/BamHI, or BsiWI/BamHI and ligated into pRB5846, which had the 1.2 kbp BsiWI/BspEI fragment already deleted. To generate the transfer plasmid pRB5848, the 4.76 kbp insert in pRB5847 was released by XbaI/EcoRV digestion and ligated into pKO5Y at the sites of XbaI and Scat Recombinant HSV-1 virus R5108 is based on R5107 with the additional deletion of the gB heparan sulfate binding domain. It was made by the same procedure as BAC-R5607, except that the transfer plasmid pRB5848 was used instead of BAC-HSV wild-type and pRB5835. The sequence of the mutant gB was verified by sequencing the entire ORF.

[0099] (iii) Deletion of gD (FIG. 1 Panel C, Lines 6 and 7).

[0100] The coding sequence of gD was replaced with the human cytomegalovirus immediate early promoter to enable the expression of glycoprotein I. A 0.65 kbp fragment containing the promoter was released from pRB5836 by ClaI digestion and inserted into pgD⁻ (FIG. 6), a plasmid obtained from G. Campadelli-Fiume. This plasmid, containing the flanking sequences of gD but lacking the gD ORF, had been cut with ClaI to generate pRB5849. pRB5849 was then cut with NotI and PmeI and ligated into pKO5Y at the NotI and ScaI sites to generate the transfer plasmid pRB5850. Recombinant HSV-1 virus R5110 is based on R5608 with the additional deletion of gD. It was made by the same procedure as BAC-R5607 except that transfer plasmid pRB5850 was used instead of BAC-HSV wild-type and pRB5835. The recombinant BAC-HSV DNA was prepared as described in (Ye et al., 2000). The mutant virus was designated R5110.

[0101] (iv) Construction of the R5111 Mutant Carrying the IL-13-gD Chimeric Gene (FIG. 1 Panel D).

[0102] Plasmid pRB123 carries a 6,584 by BamHI J fragment containing the gD coding region and flanking sequences in the BamHI site of pBR322. To construct the IL 13-gD chimeric plasmid, pRB123 was digested with AfIII and HpaI to release two fragments of 7.6 kb and 3.2 kb. The 3.2 kb fragment was further digested with FspI to release 2.5 kb and 0.7 kb fragments that contain the amino-terminal 661 bp of the gD ORF. A polylinker sequence containing the restriction sites XhoI-BglIII-EcoRI-KpnI was inserted into the 0.7 kb fragment downstream of the 24th codon of gD by two PCR reactions using a first forward primer,

(SEQ ID NO: 10)
5' - CAGTTATCCTTAAGGTCTCTTTTGTGTGGTG - 3' ,

and a first reverse primer,

(SEQ ID NO: 11)
5' - CCGGAATTCGGAGATCTTCCCTCGAGGACCGAAGGTCT
TTGCCGCGAAAG - 3' ,

and a second forward primer,

(SEQ ID NO: 12)
5' CCGGAATTCGGGGTACCCTGGACCAGCTGACCGACCTC
CGG - 3' ,

and a second reverse primer,

(SEQ ID NO: 13)
5' - CGGGGGATGCGCAGCGGGGCGTACTTAC - 3' ,

[0103] respectively. After digestion of the two PCR products by EcoRI, they were ligated and amplified by PCR again to obtain the desired DNA fragment containing the polylinker insertion.

[0104] IL13 was amplified by PCR with the forward primer,

(SEQ ID NO: 14)
5' - CCGCTCGAGATGGCGCTTTTGTGACCACGG - 3' ,

and the reverse primer,

(SEQ ID NO: 15)
5' - GGGGTACCGTTGAACCGTCCCTCGCGAAA - 3' ,

[0105] and then inserted into the XhoI and KpnI sites of the 0.7 kb fragment described above. This new fragment with the IL13 insertion was then ligated with the 2.5 kb and 7.6 kb fragments (see above) to generate the IL13-gD chimeric transfer plasmid, pRB13-24.

[0106] R5111 was generated by co-transfection of transfer plasmid pRB13-24 and the R5110 viral DNA into U87 glioma cells. The progeny of the transfection was plated at a high dilution on Vero and HEp-2 cell cultures to yield individual, well-spaced plaques. From each of the infected cell cultures, six single plaques were picked, frozen-thawed, sonicated, and then replated on fresh cultures of Vero or HEp-2 cells (depending on the origin of the plaque) for preparation of virus stocks and to prepare viral DNA for sequencing.

[0107] Viral DNA Extraction.

[0108] Infected cells were removed from each of the 25 cm² flasks exposed to individual plaque isolates, rinsed, and resuspended in 500 μ l of Lyse-O-Lot (150 mM NaCl, 10 mM Tris, 1.5 mM MgCl₂ in the presence of 0.1% of NP40). Nuclei were removed by low-speed centrifugation. To the supernatant fluid were added sodium dodecyl sulfate (SDS) to 0.2%, EDTA to 5 mM and β -ME to 50 mM. The solution was then extracted twice with phenol/chloroform. Viral DNA was finally precipitated by ethanol, resuspended, and the IL13 ORF and IL13-gD chimeric reading frame were amplified by PCR with two sets of primers. The first set, designed to amplify IL13, consisted of: a forward primer,

(SEQ ID NO: 16)
5' - CCGCTCGAGATGGCGCTTTTGTGACCACGG - 3' ,

and a reverse primer,

(SEQ ID NO: 17)
5' - GGGGTACCGTTGAACCGTCCCTCGCGAAA - 3' ,

which will amplify the IL13 ORF. The second set, designed to amplify the IL13-gD junction, consisted of a forward junction primer,

(SEQ ID NO: 18)
5' - CCGCTCGAGATGGCGCTTTTGTGACCACGG - 3' ,

and a reverse junction primer,

(SEQ ID NO: 19)
5' - AACTGCAGTTGTTCGGGGTGGCCGGGG - 3' .

All 12 IL13-gD PCR products were sequenced to determine whether the gD sequence contained deletions or substitutions.

[0109] Verification of the structure of R5111

[0110] The construction of the R5111 virus is depicted in FIG. 1. The design involved replacement of the HveA binding site with the IL13 ligand to enable the recombinant virus to bind the IL13 α 2 receptor on cell surfaces and to delete the sequences reported to bind to heparan sulfate. Verification of the structure of R5111 was done as follows:

[0111] (i) The replacement of the amino-terminal domain of gC with IL13 and the disruption of the hepa-

ran sulfate binding site was initially verified by sequencing gC from recombinant R5107 (FIG. 2 A).

[0112] (ii) The deletion of codons 68-77 of gB was verified by sequencing the gB ORF amplified by PCR from recombinant R5108 (FIG. 2 B). The nucleotide and amino acid sequences of gB with the polylysine track deleted are set out in SEQ ID NOs.:35 and 36, respectively.

[0113] (iii) The presence of chimeric IL13-gD in R5111 was verified by PCR, as illustrated in FIG. 3, and by sequencing the entire IL13-gD coding region, amplified by PCR, as shown in FIG. 2. The nucleotide and amino acid sequences of gD with the IL13 integration are set out in SEQ ID NOs.:39 and 40, respectively. The R5111 recombinant was initially isolated from transfected U87 cells and grown in parallel in Vero cells and HEp-2 cells. To determine whether the virus grown in Vero cells or HEp-2 cells differed with respect to amino acid sequence, six plaques each from Vero or HEp-2 cultures containing well-separated plaques were picked. In this series of verification experiments, two sets of primers were used to confirm the presence of the IL13 insert in gD and to verify the presence of a junction between IL13 and gD. In a second round of verifications, the 12 clones of gD were sequenced to determine whether the isolates obtained from the viruses passaged in Vero cells or in HEp-2 cells differed in amino acid sequence. No differences were found. Furthermore, except for the inserted IL13 sequence, no differences were found between the sequence of HSV-1(F) gD and those of the cloned IL-13-gD chimeric genes (FIG. 2 C).

[0114] (iv) In denaturing polyacrylamide gels, IL13 migrated as a protein with an apparent Mr of 15-17,000. In the recombinant R5111, IL13 replaced 148 amino acids of gC. FIG. 4B shows an immunoblot of electrophoretically separated proteins from a lysate of R5111 mutant-infected cells exposed to an antibody to gC. As illustrated in that figure, the anti-gC antibody reacted with proteins present in lysates of HSV-1(F) and with proteins from R5111 lysates, exhibiting similar electrophoretic mobilities. In contrast, an antibody to IL13 reacted with a band of similar mobility in R5111 lysates (FIG. 4C, lane 3) but not in lysates of HSV-1(F) (FIG. 4 C, lane 2). The IL13-gD fusion protein in the R5111 mutant virus was verified by reacting the cell lysates with gD and IL13 antibody. Comparison of wild-type gD and the chimeric IL13-gD chimeric protein (FIG. 4 A, lane 3), showed that, as expected, IL13-gD migrated more slowly than the wild-type gD (FIG. 4 A, lane 2). The faster migrating band of gD did not react with the antibody to IL13 (FIG. 4C, lane 2).

Example 2

Construction of a Cell Line Expressing the IL13 Receptor (IL13R α 2)

[0115] A rigorous test of the ability of R5111 to utilize the IL13R α 2 protein as a receptor for entry required construction of a cell line expressing IL13R α 2 (nucleotides 126-1265 of SEQ ID NO.:33; SEQ ID NO:34) in the absence of other HSV-1 entry receptors. The J1.1 cell line was selected for this construction. In essence, this cell line lacks the receptors necessary for the entry of virus into cells and the cell line is not susceptible to infection by wild-type virus. The construc-

tion of a plasmid encoding a IL13R α 2 protein fused at its carboxyl terminus to a HA tag, transfection of J1.1 cells with the plasmid encoding the tagged IL13R α 2 protein, and selection of the cell line expressing the protein is described below.

[0116] To test for the production of IL13R α 2 protein, five clones of the selected cells were harvested, solubilized, subjected to electrophoresis in denaturing polyacrylamide gels and tested for expression of the protein.

[0117] Construction of J13R, a Cell Line Stably Expressing IL13R α 2 Receptor.

[0118] The IL13 α 2 coding region was tagged with an HA tag at its 3' end (the carboxyl terminus of the encoded polypeptide) by PCR with forward primer,

(SEQ ID NO: 20)
5' - AAGATTTGGGC - TAGCATGGCTTTCGTTTGC - 3' ,

and reverse primer,

(SEQ ID NO: 21)
5' - TCCCTCGAAGCTTCAAGCATAATCTGGCACATCATATGTA
TCACAGAA - AAA - 3' .

NheI and HindIII restriction digests were used to create compatible ends. The DNA fragment was then inserted into pcDNA 3.1 (zeo) vector (Invitrogen; Carlsbad, Calif.) to generate transfer plasmid pRB13-R2. All of the constructs were sequenced to insure fidelity.

[0119] J1.1, a derivative of BHK thymidine kinase⁻ cells which lack both HveA and nectin 1 receptors, was obtained from Dr. G. Campadelli-Fiume, University of Bologna, Italy. J1.1 cells, stably transfected with pRB 13-R2 using a Lipofectamine kit (Gibco-BRL), were selected on the basis of their resistance to zeocin (Invitrogen). Zeocin-resistant clones were amplified and screened for IL13R α 2 expression by immunoblotting with anti-HA polyclonal antibody. Lysates of parental and transformed cells formed by solubilized in SDS were each electrophoretically separated in a denaturing gel (50 μ g/lane), transferred to a nitrocellulose sheet, and probed with antibody against HA (Santa Cruz Biotechnology). The protein bands were visualized by an enhanced chemiluminescent detection (ECL) system (Pierce, Rockford, Ill.) according to the instructions of the manufacturer. One (J13R-2) was selected for testing the ability of R5111 to use the IL13R α 2 receptor.

[0120] As shown in FIG. 5, all clones expressed a protein band reactive with the anti-HA antibody. The apparent size of the protein was consistent with the reported size of IL13R α 2. Of 5 J13R-positive clones, J13R-2 (FIG. 5, lane 3) was selected and designated J13R.

Example 3

Infection by the HSV Targeting Vector R5111

[0121] SK-N-SH, HEp-2, Vero, and U87 cells were obtained from American Type Culture Collection (Rockville, Md.) and maintained in Dulbecco's modification of Eagle's Minimal Essential Medium (DMEM) supplemented with 10% fetal bovine serum. Replicate cultures of SK-N-SH, HEp-2, Vero, U87, J1.1, and J13R were exposed to 0.01 PFU of R5111 virus per cell. After 24 hours of incubation, the cells were harvested and viral yields were titered on Vero cells.

[0122] Immunoblotting Electrophoretically Separated Proteins.

[0123] The indicated cells were mock-infected or exposed to 10 PFU of recombinant or wild-type HSV-1(F) per cell. The cells were harvested at 24 hours after infection, disrupted in SDS disruption buffer, boiled, cleared by centrifugation and electrophoretically separated on a 10% denaturing polyacrylamide gel. After transfer to a nitrocellulose membrane, the isolated proteins were reacted with antibodies as indicated using known and conventional techniques. Monoclonal antibodies against gD- (clone H170), gC- and HA-specific polyclonal antisera were purchased from the Goodwin Institute, Plantation, Fla. Polyclonal antibodies against IL13 were purchased from Santa Cruz Biotechnology.

[0124] The results shown in Table 1 were as follows: R5111 replicated to within a 10-fold range in HEp-2, Vero, U87, and J13R cells. The titer obtained from J1.1 cells was approximately 10^5 -fold lower than that obtained from all other cell lines. To test whether the J13R cell line had acquired a receptor for wild-type HSV-1 (HSV-1(F)), J1.1 and J13R cells were also exposed to the wild-type virus. The results, also shown in Table 1, indicate that the cells remain resistant to the wild-type virus. It was known that HEp-2 cells express the nectin receptor but not the HveA receptor. The results show that the targeted HSV containing an IL13-gD fusion can target (i.e., bind and infect) cells expressing a particular receptor (IL13R α 2) approximately as well as wild-type HSV targets cells expressing the HveA receptor. The results indicate that R5111 can use IL13R α 2 as a receptor for entry in a cell line lacking all other HSV-1 receptors.

TABLE 1

Replication of R5111 in various cell lines		
Virus	Cell Line*	Yield**
R5111	Vero	11×10^7
	HEp-2	1.2×10^7
	SK-N-SH	17×10^7
	U87	27×10^7
	J1.1	2×10^2
	J13R	11×10^7
HSV-1(F)	J1.1	6×10^3
	J13R	4×10^3

*cell lines derived from human brain tumors.

**The cells were exposed to 0.01 PFU of R5111 or HSV-1(F) per cell and harvested 24 hours after infection. Progeny virus were titered on Vero cells.

[0125] This disclosure contains an exemplary description of the construction and properties of a recombinant HSV virus, R5111. In R5111, the heparan sulfate binding sites on the surface of the viral particle were ablated to preclude or at least reduce the attachment of virus to non-targeted cells. Attachment even in the absence of fusogenic activity may lead to endocytosis, degradation of the virus particle, and to potential damage to the cell by lysosomal enzymes (Zhou et al. 2002; Zhou et al. 2000). At the same time, a copy of IL13 was inserted into gC to enhance binding of virus particles to the IL13R α 2 receptor. The major restructuring of the viral genome, consisted of insertion of IL13 at amino acid 24 of gD. Available data indicate that this modification ablates the gD binding site for the HveA receptor (Carfi et al. 2001). The data obtained using R5111 indicate that the virus retains the capacity to interact with the Nectin receptor. Nonetheless, the R5111-targeted HSV was able to infect and replicate in J13R cells but not in the parental, J1.1, cells.

Example 4

Construction of HSV Targeting Vector R5141 and R5144

[0126] A therapeutic herpes simplex virus 1 (HSV-1) capable of infecting and replicating solely in cells harboring the IL13R α 2 receptor was constructed using recombinant DNA techniques. As disclosed above, construction of R5111, which expresses IL13 on its surface and lacks the binding sites for heparin sulfate, allowed R5111 to infect 3-13R cells as well as cells exhibiting the natural receptors for HSV-1. Thus, the involvement of fusogenic glycoproteins of R5111—a key step in viral entry—is independent of the receptor with which gD interacts.

[0127] The binding site of HveA has been reported to be at the amino terminal domain of gD (Carfi A., et al., 2001). The precise binding sites of gD for Nectin 1 are not known, although it has previously been reported to involve gD amino acids 38 and 221 (Manoj S., et al., 2004; Zago A., et al., 2004; Connolly S.A., 2005). The general assumption within the field is that the HveA and Nectin 1 sites do not overlap and that each independently promotes the same structural alteration of gD to enable entry of the virus into cells. However, it is possible that the surface structure of HveA and Nectin 1 at their binding sites may be similar even though the Hvel and Nectin1 amino acid sequences are not identical.

[0128] Based in part on the foregoing information, viruses capable of productive replication solely in targeted cells were designed as shown in FIG. 7. Using standard molecular biological cloning techniques known in the art, recombinant virus R5141 was constructed by inserting IL13 in the place of gD residues 1-32. In addition, the valine residue at position 34 was substituted with serine (“V34S”) (SEQ ID NOs.:41 and 42, respectively). Similarly, recombinant virus R5144 was constructed by inserting IL13 in the place of gD residues 1-32, and the valine at position 37 was substituted with serine (“V37S”) (SEQ ID NOs.:43 and 44, respectively).

[0129] One of skill would appreciate that a variety of re-targeted SHV, both HSV-1 and HSV-2, could be constructed and assessed using routine techniques in view of the disclosures herein. In particular, substituting a binding domain of a binding partner for the N-terminal region of gD, e.g., for amino acids 1-32 of gD, would be within the skill in the art. Further alteration of the gD fusion, e.g., by amino acid substitution, whether conservative substitution or not, would also be within the skill in the art. Of course, additions or deletions to gD fusions would also be within the skill in the art and assessing the targeting capacities of such constructs would involve routine experimentation in view of the teachings herein.

Example 5

Infection by the HSV Targeting Vectors R5141 and R5144

[0130] The capacities of the recombinant viruses described in Example 4 to productively replicate solely in targeted cells were assessed using the cell lines which express either HveA (J-HveA) alone (relative to the group of HveA, Nectin1, and IL13R α 2), Nectin1 (J-Nectin1) alone, or IL13R α 2 (J-13R) alone. For cell infection, the procedure set forth in Example 3 was repeated using recombinant virus R5141 and recombinant virus R5144.

[0131] The replication of R5141 and R5144 in J-Nectin-, J-HveA-, and J-13R-specific cells are summarized in Table 2. R5141 and R5144 do not productively interact with either native gD receptors, HveA or Nectin1. Significantly, however, R5141 interacts with, and replicates in IL13R α 2 for productive entry into cells.

TABLE 2

Replication of R5141 and R5144 in J-Nectin, J-HveA and J-13R cells.			
	HSV-1 (F)	R5141	R5144
J-Nectin	4×10^8	7×10^1	5×10^1
J-HveA	3×10^8	4×10^1	3×10^2
J-13R	3×10^1	5×10^6	7×10^2

[0132] Thus, recombinant virus R5141 is capable of productive replication solely in targeted cells and this result opens the way for development of therapeutic viruses targeting cells exhibiting the IL13R α 2 receptor, such as malignant gliomas and other human tumors exhibiting IL13R α 2. It is expected that other mutations (i.e., those that abolish binding of Nectin and those that have a similar effect on HveA) will yield viruses that enter solely via non-natural HSV receptors.

Example 6

HSV Targeting Vector R5161

[0133] An HSV targeting vector designated HSV R5161 has a structure analogous to the structure of HSV R5141, and HSV R5161 was constructed in the manner described in Example 4, above, with the exception that HSV R5161 contains the sequence encoding the HSV gD leader sequence, whereas HSV R5141 contains the sequence encoding the IL-13 leader sequence. In particular, HSV R5161 encodes an IL-13-gD fusion protein in which IL-13 sequence replaces the sequence encoding gD amino acids 1-32, with a V34S substitution in the gD moiety of the fusion protein, as described above in the context of describing HSV R5141.

[0134] The relative capacities of recombinant viruses HSV R5141 and HSV R5161 to productively replicate in targeted cells were measured in the J-13R cell line, which expresses IL13R α 2, but not HveA or Nectin1. HSV R5161 was expressed at a level 10-fold higher than the expression level of HSV R5141 in J-13R cells.

[0135] It is expected that in the majority of embodiments of the invention, relatively high levels of expression will be advantageous in ensuring that targeted cells are efficiently contacted by the re-targeted HSV. In re-targeting the HSV, the invention provides an approach to controlling the virulence of the virus in a manner that minimizes undesirable pathogenicity, i.e., pathogenicity towards non-targeted cells. In addition, the virulence of any re-targeted HSV can be further attenuated using known approaches to virulence control that do not interfere with the re-targeting, such as by mutating the γ_1 34.5 gene(s). In those embodiments in which it is desirable to have relatively high expression levels of the re-targeted HSV, it is preferred that the leader sequence of gD be used. Alternative leader sequences, such as leaders from other HSV genes, are contemplated. Moreover, expression control elements (e.g., promoters, enhancers, expression factor binding sites) can be

engineered to achieve desired expression levels of the fusion protein using ordinary levels of skill and techniques known in the art.

REFERENCES

- [0136]** Davis F G, Freels S, Grutsch J, Barlas S, Brem S. (1998) *J. Neurosurg.* 88:1-10.
- [0137]** Pyles R B, Warrick R E, Chalk C L, Szanti B E, Parysek L M. (1997) *Hum Gene Ther.* 8(5):533-44.
- [0138]** Rampling R, Cruickshank G, Papanastassiou V, Nicoll J, Hadley D, Brennan D, Petty R, MacLean A, Harland J, McKie E, Mabbs R, Brown M. (2000) *Gene Ther.* 7(10):859-66.
- [0139]** McKie E A, Brown S M, MacLean A R, Graham D I. (1998) *Neuropathol. Appl Neurobiol.* 24(5):367-72.
- [0140]** Markert J M, Medlock M D, Rabkin S D, Gillespie G Y, Todo T, Hunter W D, Palmer C A, Feigenbaum F, Tornatore C, Tufaro F, Martuza R L. (2000) *Gene Ther.* 7(10):867-74.
- [0141]** Mineta T, Rabkin S D, Yazaki T, Hunter W D, Martuza R L. (1995) *Nat Med.* 1(9):938-43.
- [0142]** Simard C, Langlois I, Styger D, Vögt B, Vlcek C, Chalifour A, Trudel M, Schwyer M. (1995) *Virology* 212 (2):734-40.
- [0143]** Chou J, Chen J J, Gross M, Roizman B. (1995) *Proc Natl Acad Sci USA.* 92(23):10516-20.
- [0144]** He B, Chou J, Brandimarti R, Mohr I, Gluzman Y, Roizman B. (1997) *J Virol.* 71(8):6049-54
- [0145]** Cassidy K A, Gross M, Roizman B. (1998) *J Virol.* 72(9):7005-11.
- [0146]** Leib, D. A., Harrison, T. E., Laslo, K. M., Machalek, M. A., Moorman N. J. and Virgin, H. A W. (1999) *J. Exp. Med.* 189:663-672.
- [0147]** Laquerre S, Argnani R, Anderson D B, Zucchini S, Manservigi R, Glorioso J C. (1998). *J Virol.* 72(7):6119-30.
- [0148]** Spear, P. G., R. J. Eisenberg, and G. H. Cohen. (2000) *Virology* 275: 1-9.
- [0149]** Montgomery, R. L., M. S. Warner, B. J. Lum, and P. G. Spear. (1996) *Cell* 87:427-436.
- [0150]** Campadelli-Fiume, G., F. Cocchi, L. Menotti, and M. Lopez. (2000) *Reviews in Medical Virology.* 10:305-319.
- [0151]** Zhou G, Roizman B. (2002) *J Virol.* 76(12):6197-204.
- [0152]** Debinski W, Gibo D M, Hulet S W, Connor J R, Gillespie G Y. (1999) *Cancer Res.* 5:985-990.
- [0153]** Mintz A, Gibo D M, Slagle-Webb B, Christensen N D, Debinski W. (2002) *Neoplasia* 4:388-399.
- [0154]** Debinski W. (1998) *Crit. Rev. Oncogen.* 9:255-268.
- [0155]** Debinski W, Gibo D M. (2000) *Mol. Med.* 6:440-449.
- [0156]** Zhou G, Roizman B. (2001) *J Virol.* 75(13):6166-72.
- [0157]** Arsenakis M, Tomasi L F, Speziali V, Roizman B, Campadelli-Fiume G. (1986) *J Virol.* 58(2):367-76.
- [0158]** Ye G J, Roizman B. (2000) *Proc Natl Acad Sci USA.* 97(20):11002-7.
- [0159]** Zhou G, Galvan V, Campadelli-Fiume G, Roizman B. (2000) *J Virol* 74(24):11782-91.
- [0160]** Carfi A, Willis S H, Whitbeck J C, Krummenacher C, Cohen G H, Eisenberg R J, Wiley D C. (2001) *Mol Cell.* 8(1):169-79.

- [0161] Cocchi, F., Menotti, L., Mirandola, P and Campadelli-Ofiume, G. (1998) *J. Virol.* 72:9992-10002.
- [0162] Debinski W, Thompson J.P. 1999. *Clin Cancer Res.* 5(10 Suppl):3143s-3147s.
- [0163] Brooks, P. C., Clark, R. A. F., and Cheresch, D. A. Requirement of vascular integrin $\alpha_v\beta_3$ for angiogenesis. *Science* 264:569-571, 1994.
- [0164] Brooks, P. C., Montgomery, A. M. P., Rosenfeld, M., Reisfeld, R. A., H U, T., Klier, G., and Cheresch, D. A. Integrin $\alpha_v\beta_3$ antagonists promote tumor regression by inducing apoptosis of angiogenic blood vessels. *Cell* 79:1157-1164, 1994.
- [0165] Burger, M. J., Tebay, M. A., Keith, P. A., Samarantunga, H. M., Clements, J., Lavin, M. F., and Gardiner, R. A. Expression analysis of 6-Catenin and prostate-specific membrane antigen: Their potential as diagnostic markers for prostate cancer. *Int. J. Cancer* 100:228-237, 2002.
- [0166] Ellerman, T. C., Domagala, T., McKem N. T. et al, Identification of the determinant of Epidermal growth factor receptor ligand-binding specificity using truncated, high affinity form of the ectodomain. 2001 *Biochemistry* 40 8930-8939.
- [0167] Genbitsky, D. S., Bozso, Z., O'Flaharty, M. et al., 2001 A specific binding site for a fragment of the B-loop of epidermal growth factor and related peptides. *Peptides* 23:97-102 A.
- [0168] Urbanelli, L., Ronchini, C., Fontana, L. et al., Targeted gene transduction of mammalian cells expressing the HER2/neu receptor by filamentous phage. *J Mol Biol.* 2001 Nov. 9; 313(5):965-76.
- [0169] Hayashi, T., Takahashi, T., Motoya, S., et al. MUC1 Mucin core protein binds to the dfomasin of ICAM-1 2001 *Digestion* 63:87-92.
- [0170] Fracasso, G., Bellisola, G., Cingarlini, S., Castelletti, D., Prayer-Galletti, T., Pagano, F., Tridente, G., and Colombatti, M. Anti-tumor effects of toxins targeted to the prostate specific membrane antigen. *Prostate* 53:9-23, 2002.
- [0171] Mabjeesh, N. J., Zhong, H., and Simons, J. W. Gene therapy of prostate cancer: current and future directions. *Endo. Related Cancer* 9:115-139, 2002.
- [0172] Ross, S., Spencer, S. D., Holcomb, I., Tan, C., Hongo, J., Devaux, B., Rangell, L., Keller, G. A., Schow, P., Steeves, R. M., Lutz, R. J., Frantz, G., Hillan, K., Peale, F., Tobin, P., Eberhard, D., Rubin, M. A., Lasky, L. A., and Koeppen, H. Prostate stem cell antigen as therapy target: Tissue expression and in Vivo efficacy of an immunocjugate. *Cancer Res.* 62:2546-2553, 2002.
- [0173] Ruoslahti, E. RGD and other recognition sequences for integrins. *Annu. Rev. Cell Dev. Biol.* 12:697-715, 1996.
- [0174] Thomas, J., Gupta, M., Grasso, Y., Reddy, C. A., Heston, W. D., Zippe, C., Dreicer, R., Kupelian, P. A., Brainard, J., Levin, H. S., and Klein, E. A. Preoperative combined nested reverse transcriptase polymerase chain reaction for prostate-specific antigen and prostate-specific membrane antigen does not correlate with pathologic stage or biochemical failure in patients with localized prostate cancer undergoing radical prostatectomy. *J. Clin. Oncol.* 20:3213-3218, 2002.
- [0175] Lorimer and Lavictoire, Targeting retrovirus to cancer cells expressing a mutant EGF receptor by insertion of a single chain antibody variable domain in the envelope glycoprotein receptor binding lobe, *J Immunol Methods* 237(1-2):147-57, 2000.
- [0176] Turner et al., Glycoproteins gB, gD, and gHgL of Herpes Simplex Virus Type 1 are Necessary and Sufficient t Mediate membrane fusion in a Cos cell transfection system, *J of Virol.* 72(1): 873-75, 1998.
- [0177] Brunetti et al., Herpes Simplex Virus gD and Virions Accumulate in Endosomes by Mannose 6-Phosphate-Dependent and -Independent Mechanisms, *J of Virol.* 72(4): 3330-3339, 1998.
- [0178] Sharma et al., Molecular imaging of gene expression and protein function in vivo with PET and SPECT, *J Magn Reson Imaging*, 16(4):336-51, 2002.
- [0179] Vries et al., Scintigraphic Imaging of HSVtk Gene Therapy, *Curr Pharm Des*, 8(16):1435-50, 2002.
- [0180] Vries et al., Positron emission tomography: measurement of transgene expression, *Methods*, 27(3):234, 2002.
- [0181] Soling et al., Intracellular localization of Herpes simplex virus type 1 thymidine kinase fused to different fluorescent proteins depends on choice of fluorescent tag, *FEBS Lett*, 527(1-3):153, 2002.
- [0182] Zhou G., and Roizman B., *J Virol.* 9:5272-5277 (2005)
- [0183] Manoj S., et al., *Proc. Natl. Acad. Sci. USA.*, 101: 12414-12421 (2004);
- [0184] Zago A., et al., *Proc. Natl. Acad. Sci. USA.*, 101: 17498-17503 (2004);
- [0185] Connolly S A., *J. Virol.*, 79:1282-1295 (2005)

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 44

<210> SEQ ID NO 1
 <211> LENGTH: 60
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 1

gattgctctc acttgcccttg gcggettgtgc ctccccaggg cctgtgcctc cctctacagc 60

<210> SEQ ID NO 2
 <211> LENGTH: 60
 <212> TYPE: DNA

-continued

<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 2

gcagctagcc tcatggcgct tttgttgacc acggtcattg ctctcacttg ccttggcggc 60

<210> SEQ ID NO 3
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 3

gagctcggat cctgaattca accgtccctc 30

<210> SEQ ID NO 4
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 4

gacacgggct acctcacta tcgagggc 28

<210> SEQ ID NO 5
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 5

ggtgatgttc gtcaggacct cctctaggtc 30

<210> SEQ ID NO 6
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 6

gtttttcttc ggtttcggat cccccg 26

<210> SEQ ID NO 7
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 7

cggcatttcc ggaataacgc ccaactc 26

<210> SEQ ID NO 8
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

-continued

<400> SEQUENCE: 8
cagaaaaccg gatcccccaa agccgcc 27

<210> SEQ ID NO 9
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 9
gccaacacaa actcgtcgta cgggtac 27

<210> SEQ ID NO 10
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 10
cagttatcct taaggtctct tttgtgtggt g 31

<210> SEQ ID NO 11
<211> LENGTH: 52
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 11
ccggaattcc ggagatcttc cctcgaggac cggaaggtct ttgccgcgaa ag 52

<210> SEQ ID NO 12
<211> LENGTH: 43
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 12
ccggaattcc ggggtaccct ggaccagctg accgaccctc cgg 43

<210> SEQ ID NO 13
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 13
cgggggatg cgcagcggga gggcgactt ac 32

<210> SEQ ID NO 14
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 14
ccgctcgaga tggcgctttt gttgaccag g 31

-continued

<210> SEQ ID NO 15
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 15

ggggtaccgt tgaaccgtcc ctgcgaaa 29

<210> SEQ ID NO 16
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 16

ccgctcgaga tggcgctttt gttgaccacg g 31

<210> SEQ ID NO 17
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 17

ggggtaccgt tgaaccgtcc ctgcgaaa 29

<210> SEQ ID NO 18
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 18

ccgctcgaga tggcgctttt gttgaccacg g 31

<210> SEQ ID NO 19
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 19

aactgcaggt tgttcggggt ggccggggg 29

<210> SEQ ID NO 20
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 20

aagatttggg ctagcatggc tttcgtttgc 30

<210> SEQ ID NO 21
<211> LENGTH: 51
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic primer

 <400> SEQUENCE: 21

 tccctcgaag cttcaagcat aatctggcac atcatatgta tcacagaaaa a 51

<210> SEQ ID NO 22
 <211> LENGTH: 568
 <212> TYPE: DNA
 <213> ORGANISM: Herpes Simplex Virus-1

 <400> SEQUENCE: 22

 gcttggtcgg gaggcgcgat cgaacgcaca cccccatecg gtggtccgtg tggaggctgt 60
 ttttcagtgc cgggtctcgc tttgcogga acgctagcct catggcgctt ttgttgacca 120
 cggtcattgc tetcaattgc cttggcggtc ttgcctcccc aggccctgtg cctcccctca 180
 cagccctcag gtacctcatt gaggagctgg tcaacatcac ccagaaccag aaggctccgc 240
 tctgcaatgg cagcatggta tggagcatca acctgacagc tggcatgtac tgtgcagccc 300
 tggaatccct gatcaacgtg tcaggctgca gtgccatcga gaagaccagc aggatgctga 360
 gcggattctg cccgcacaag gtctcagctg ggcagtttcc cagcttgcct gtccgagaca 420
 ccaaaatcga ggtggcccag tttgtaaaag atctgctctt acatttaaag aaacttttcc 480
 gcgaggggacg gttgaattcc acccgcatgg agttccgect ccagatatgg cgttactcca 540
 tgggtccgctc ccccccaatc gctccggc 568

<210> SEQ ID NO 23
 <211> LENGTH: 350
 <212> TYPE: DNA
 <213> ORGANISM: Herpes Simplex Virus-1

 <400> SEQUENCE: 23

 gggctcctggt ggcgtcggcg gctccgagtt cccccggcac gcctgggggc gcggcccgca 60
 cccaggcggc gaacggggga cctgccactc cggcgccgcc cgcacctggc cccgcccaca 120
 cgggggatcc gaaaccgaag aagaacagaa aaccgaaacc cccaaagcgc cgcgccccgc 180
 cggcgacaac ggcacctg cgcggggcca cgcaccctg cgcgagcacc tgcgggacat 240
 caagcgggag aacaccgatg caaactttta cgtgtgcccc cccccacgg gcgcccaggt 300
 ggtgcagttc gacgagccgc gccgctgccc gaccggccc gagggtcaga 350

<210> SEQ ID NO 24
 <211> LENGTH: 655
 <212> TYPE: DNA
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Amino-terminal sequence of IL13-gD

 <400> SEQUENCE: 24

 atgggggggg ctgccccag gttgggggcc gtgattttgt ttgtcgtcat agtcggcctc 60
 catgggggtcc gcggcaaata tgccctggcg gatgcctctc tcaagctggc cgaccccaat 120
 cgctttcgcc gaaaagacct tccggtcctc gagatggcgc ttttgttgac cacggtcatt 180
 gctctcactt gccttgggcg ctttgctccc ccaggccctg tgcctcccct ctacagccct 240
 caggagctc attgaggagc tggtaaacat caccagaac cagaaggctc cgctctgcaa 300

-continued

```

tggcagcatg gtttgagca tcaacctgac agctggcatg tactgtgcag ccttggaaac 360
cctgatcaac gtgtcaggct gcagtgccat cgagaagacc cagaggatgc tgggcccatt 420
ctgcccgcac aaggtctcag ctgggcagtt ttccagcttg catgtccgag acacccaaat 480
cgaggtggcc cagttttaa aggacctgct cttacattta aagaactttt ttcgcgaggg 540
acggttcaac tgaacggta ccttggacca gctgaccgac cctccggggg tccggcgcgt 600
gtaccacatc caggcgggcc taccggacc gttccagccc cccagcctcc cgatc 655

```

```

<210> SEQ ID NO 25
<211> LENGTH: 1185
<212> TYPE: DNA
<213> ORGANISM: Herpes Simplex Virus-1

```

```

<400> SEQUENCE: 25

```

```

atgggggggg ctgccccag gttgggggcc gtgattttgt ttgtcgtcat agtgggcctc 60
catgggggtcc gcggcaata tgccttggcg gatgcctctc tcaagatggc cgacccaat 120
cgctttcgcg gaaaagacct tccggtcccg gaccggctga cgcacctcc gggggtccgg 180
cgcgtgtacc acatccaggc gggcctaccg gaccggtcc agccccccag cctcccgatc 240
acggtttact acgccgtgtt ggagcgcgcc tgccgcagcg tgcctctaaa cgcaccgtcg 300
gaggcccccc agattgtccg cggggcctcc gaagacgtcc ggaacaacc ctacaacctg 360
accatcgctt ggtttcggat gggaggcaac tgtgctatcc ccatcacggt catggagtac 420
accgaatgct cctacaacaa gtctctgggg gctgtccca tccgaacgca gccccgctgg 480
aactactatg acagcttcag cgccgtcagc gaggataacc tggggttcct gatgcacgcc 540
ccccgctttg agaccgcccg cacgtacctg cggctcgtga agataaacga ctggacggag 600
attacacagt ttatcctgga gcaccgagcc aagggtcctt gtaagtacgc ccttccgctg 660
cgcatcccc cgtcagcctg cctctcccc caggcctacc agcagggggg gacgggtggac 720
agcatcggga tgcgtcccc ctatcccc gagaaccagc gcaccgtcgc cgtatacagc 780
ttgaagatcg ccgggtggca cgggcccagg gcccataca cgagcacctt gctgcccccg 840
gagctgtccg agacccccaa cgccaacgag ccagaactcg ccccgaaga ccccgaggat 900
tcggccctct tggaggacc cgtggggacg gtggtgccc aaatcccacc aaactggcac 960
atacctgca tccaggacgc cgcgacgctt taccatcccc cggccacccc gaacaacatg 1020
ggcctgatcg ccggcgcggg gggcggcagt ctctggttag ccttggatcat ttgcggaatt 1080
gtgtactgga tgcgcccgcg cactcaaaaa gcccacaaagc gcatacgcct cccccacatc 1140
cgggaagacg accagccgct ctcgcaccag cccttgtttt actag 1185

```

```

<210> SEQ ID NO 26
<211> LENGTH: 394
<212> TYPE: PRT
<213> ORGANISM: Herpes Simplex Virus-1

```

```

<400> SEQUENCE: 26

```

```

Met Gly Gly Ala Ala Ala Arg Leu Gly Ala Val Ile Leu Phe Val Val
1           5           10          15

Ile Val Gly Leu His Gly Val Arg Gly Lys Tyr Ala Leu Ala Asp Ala
20          25          30

Ser Leu Lys Met Ala Asp Pro Asn Arg Phe Arg Gly Lys Asp Leu Pro
35          40          45

```

-continued

Val Pro Asp Arg Leu Thr Asp Pro Pro Gly Val Arg Arg Val Tyr His
 50 55 60
 Ile Gln Ala Gly Leu Pro Asp Pro Phe Gln Pro Pro Ser Leu Pro Ile
 65 70 75 80
 Thr Val Tyr Tyr Ala Val Leu Glu Arg Ala Cys Arg Ser Val Leu Leu
 85 90 95
 Asn Ala Pro Ser Glu Ala Pro Gln Ile Val Arg Gly Ala Ser Glu Asp
 100 105 110
 Val Arg Lys Gln Pro Tyr Asn Leu Thr Ile Ala Trp Phe Arg Met Gly
 115 120 125
 Gly Asn Cys Ala Ile Pro Ile Thr Val Met Glu Tyr Thr Glu Cys Ser
 130 135 140
 Tyr Asn Lys Ser Leu Gly Ala Cys Pro Ile Arg Thr Gln Pro Arg Trp
 145 150 155 160
 Asn Tyr Tyr Asp Ser Phe Ser Ala Val Ser Glu Asp Asn Leu Gly Phe
 165 170 175
 Leu Met His Ala Pro Ala Phe Glu Thr Ala Gly Thr Tyr Leu Arg Leu
 180 185 190
 Val Lys Ile Asn Asp Trp Thr Glu Ile Thr Gln Phe Ile Leu Glu His
 195 200 205
 Arg Ala Lys Gly Ser Cys Lys Tyr Ala Leu Pro Leu Arg Ile Pro Pro
 210 215 220
 Ser Ala Cys Leu Ser Pro Gln Ala Tyr Gln Gln Gly Val Thr Val Asp
 225 230 235 240
 Ser Ile Gly Met Leu Pro Arg Phe Ile Pro Glu Asn Gln Arg Thr Val
 245 250 255
 Ala Val Tyr Ser Leu Lys Ile Ala Gly Trp His Gly Pro Lys Ala Pro
 260 265 270
 Tyr Thr Ser Thr Leu Leu Pro Pro Glu Leu Ser Glu Thr Pro Asn Ala
 275 280 285
 Thr Gln Pro Glu Leu Ala Pro Glu Asp Pro Glu Asp Ser Ala Leu Leu
 290 295 300
 Glu Asp Pro Val Gly Thr Val Val Pro Gln Ile Pro Pro Asn Trp His
 305 310 315 320
 Ile Pro Ser Ile Gln Asp Ala Ala Thr Pro Tyr His Pro Pro Ala Thr
 325 330 335
 Pro Asn Asn Met Gly Leu Ile Ala Gly Ala Val Gly Gly Ser Leu Leu
 340 345 350
 Val Ala Leu Val Ile Cys Gly Ile Val Tyr Trp Met Arg Arg Arg Thr
 355 360 365
 Gln Lys Ala Pro Lys Arg Ile Arg Leu Pro His Ile Arg Glu Asp Asp
 370 375 380
 Gln Pro Ser Ser His Gln Pro Leu Phe Tyr
 385 390

<210> SEQ ID NO 27

<211> LENGTH: 2715

<212> TYPE: DNA

<213> ORGANISM: Herpes Simplex Virus-1

<400> SEQUENCE: 27

atgcccagg ggcccccgc gcggggggcgc cggtggttcg tcgtatgggc gctcttgggg 60

-continued

ttgacgctgg	gggtcctggt	ggcgtcggcg	gctccgagtt	cccccgcaac	gectggggte	120
gcggccgcga	cccaggcggc	gaacgggggc	cctgccactc	cggcgcgcgc	cgccccctggc	180
gcccccccaa	cgggggacc	gaaacggaag	aagaacagaa	aaccgaaacc	cccaaagccg	240
ccgcgccccg	ccggcgacaa	cgcgaccgtc	gccgcggggc	acgccaccct	gcgcgagcac	300
ctgcgggaca	tcaaggcggg	gaacaccgat	gaaaactttt	acgtgtgccc	acccccacg	360
ggcgccacgg	tgggtgcagtt	cgagcagccg	cgccgctgcc	cgaccggcc	cgagggtcag	420
aactacacgg	agggcatcgc	ggtggtcttc	aaggagaaca	tcgccccgta	caagttcaag	480
gccaccatgt	actacaaaga	cgtaaccgtt	tcgcaggtgt	ggttcggcca	ccgctactcc	540
cagtttatgg	ggatccttga	ggaccgcgcc	cccgtcccct	tcgaggaggt	gatcgacaag	600
atcaacgcc	aggggctctg	tcggtcacg	gccaagtacg	tcgcaacaa	cctggagacc	660
accggtttc	accgggacga	ccacgagacc	gacatggagc	tgaaacggc	caacgcccgc	720
accgcacga	gcccgggctg	gcacaccacc	gacctcaagt	acaaccctc	gcgggtggag	780
gcgttcacc	ggtacgggac	gacggtaaac	tgcatcgtcg	aggaggtgga	cgcgcgctcg	840
gtgtaccgt	acgacgagtt	tgtgtggcg	actggcgact	ttgtgtacat	gtccccgttt	900
tacggctacc	gggaggggtc	gcacaccgaa	cacaccagct	acgccgcga	ccgcttcaag	960
caggctcagc	gcttctacgc	gcgcgacctc	accaccaagg	cccgggccac	ggcgccgacc	1020
accggaacc	tgctcacgac	ccccaaagtc	accgtggcct	gggactgggt	gccaagcgc	1080
ccgtcggct	gcaccatgac	caagtggcag	gaggtggacg	agatgctgcg	ctccgagtac	1140
ggcgctcct	tcgattctc	ttccgacgcc	atatccacca	ccttcaccac	caacctgacc	1200
gagtaccgc	tctcgcgct	ggacctgggg	gactgcatcg	gcaaggacgc	ccgcgacgcc	1260
atggaccgca	tcttcgcccc	caggtacaac	gcgacgcaca	tcaaggtggg	ccagccgcag	1320
tactacctgg	ccaatggggg	ctttctgac	gcgtaccagc	cccttctcag	caaacgctc	1380
gcggagctgt	acgtgcggga	acacctccgc	gagcagagcc	gcaagcccc	aaacccacg	1440
cccccgccg	ccggggccag	cgccaacgcg	tcctgggagc	gcatcaagac	caactcctcc	1500
atcgagtctg	ccaggctgca	gtttacgtac	aaccacatac	agcgcctatg	caacgatatg	1560
ttgggcccgc	ttgccatcgc	gtggtgcgag	ctgcagaatc	acgagctgac	cctgtggaac	1620
gaggcccgca	agctgaaccc	caacgccatc	gcctcggcca	ccgtgggccc	gcgggtgagc	1680
gcgaggatgc	tcggcgacgt	gatggcgcgc	tcacagtgcg	tcgcggtcgc	cgcggaacaac	1740
gtgatcgtcc	aaaactcgat	gcgcatcagc	tcgcgccccg	ggcctgcta	cagccgcccc	1800
ctggtcagct	ttcggtagca	agaccagggc	ccgttggtcg	aggggcagct	gggggagaac	1860
aacgagctgc	ggctgacgcg	cgatgcgatc	gagccgtgca	ccgtgggaca	ccggcgctac	1920
ttcaccttcg	gtgggggcta	cgtgtacttc	gaggagtacg	cgtactcca	ccagctgagc	1980
cgcgccgaca	tcaccaccgt	cagcaacctc	atcgacctca	acatcaccat	gctggaggat	2040
cacgagtctg	tccccctgga	ggtgtacacc	cgccacgaga	tcaaggacag	cgccctgctg	2100
gactacacgg	aggtccagcg	ccgcaaccag	ctgcacgacc	tgcgcttcgc	cgacatcgac	2160
acggteatcc	acgccagcgc	caacgcccgc	atgtttgceg	gcctgggccc	gctcttcgag	2220
gggatgggcg	acctggggcg	cgcggtcggc	aaggtggtga	tgggcatcgt	ggcgccgctg	2280
gatatggccc	tgctggggct	gtcctccttc	atgtccaacc	cctttggggc	gctggccgtg	2340

-continued

```

ggctctgttg tcttgccgg cctggcggcg gccttcttcg cctttcgcta cgtcatgceg 2400
ctgcagagca accccatgaa ggcctgttac cgcctaacca ccaaggagct caagaacccc 2460
accaacccgg acgcgtccgg ggaggcggag gagggcggcg actttgacga ggccaagcta 2520
gccgaggccc gggagatgat acggtacatg gccctggtgt ctgccatgga ggcacacggaa 2580
cacaaggcca agaagaaggg cacgagcgcg ctgctcagcg ccaaggtcac cgacatggtc 2640
atgcgcaagc gccgcaacac caactacacc caagtccca acaagacgg tgacgccgac 2700
gaggacgacc tgtga 2715

```

```

<210> SEQ ID NO 28
<211> LENGTH: 904
<212> TYPE: PRT
<213> ORGANISM: Herpes Simplex Virus-1

```

```

<400> SEQUENCE: 28

```

```

Met Arg Gln Gly Ala Pro Ala Arg Gly Arg Arg Trp Phe Val Val Trp
1           5           10           15
Ala Leu Leu Gly Leu Thr Leu Gly Val Leu Val Ala Ser Ala Ala Pro
20           25           30
Ser Ser Pro Gly Thr Pro Gly Val Ala Ala Ala Thr Gln Ala Ala Asn
35           40           45
Gly Gly Pro Ala Thr Pro Ala Pro Pro Ala Pro Gly Ala Pro Pro Thr
50           55           60
Gly Asp Pro Lys Pro Lys Lys Asn Arg Lys Pro Lys Pro Pro Lys Pro
65           70           75           80
Pro Arg Pro Ala Gly Asp Asn Ala Thr Val Ala Ala Gly His Ala Thr
85           90           95
Leu Arg Glu His Leu Arg Asp Ile Lys Ala Glu Asn Thr Asp Ala Asn
100          105          110
Phe Tyr Val Cys Pro Pro Pro Thr Gly Ala Thr Val Val Gln Phe Glu
115          120          125
Gln Pro Arg Arg Cys Pro Thr Arg Pro Glu Gly Gln Asn Tyr Thr Glu
130          135          140
Gly Ile Ala Val Val Phe Lys Glu Asn Ile Ala Pro Tyr Lys Phe Lys
145          150          155          160
Ala Thr Met Tyr Tyr Lys Asp Val Thr Val Ser Gln Val Trp Phe Gly
165          170          175
His Arg Tyr Ser Gln Phe Met Gly Ile Phe Glu Asp Arg Ala Pro Val
180          185          190
Pro Phe Glu Glu Val Ile Asp Lys Ile Asn Ala Lys Gly Val Cys Arg
195          200          205
Ser Thr Ala Lys Tyr Val Arg Asn Asn Leu Glu Thr Thr Ala Phe His
210          215          220
Arg Asp Asp His Glu Thr Asp Met Glu Leu Lys Pro Ala Asn Ala Ala
225          230          235          240
Thr Arg Thr Ser Arg Gly Trp His Thr Thr Asp Leu Lys Tyr Asn Pro
245          250          255
Ser Arg Val Glu Ala Phe His Arg Tyr Gly Thr Thr Val Asn Cys Ile
260          265          270
Val Glu Glu Val Asp Ala Arg Ser Val Tyr Pro Tyr Asp Glu Phe Val
275          280          285

```

-continued

Leu Ala Thr Gly Asp Phe Val Tyr Met Ser Pro Phe Tyr Gly Tyr Arg
 290 295 300

Glu Gly Ser His Thr Glu His Thr Ser Tyr Ala Ala Asp Arg Phe Lys
 305 310 315 320

Gln Val Asp Gly Phe Tyr Ala Arg Asp Leu Thr Thr Lys Ala Arg Ala
 325 330 335

Thr Ala Pro Thr Thr Arg Asn Leu Leu Thr Thr Pro Lys Phe Thr Val
 340 345 350

Ala Trp Asp Trp Val Pro Lys Arg Pro Ser Val Cys Thr Met Thr Lys
 355 360 365

Trp Gln Glu Val Asp Glu Met Leu Arg Ser Glu Tyr Gly Gly Ser Phe
 370 375 380

Arg Phe Ser Ser Asp Ala Ile Ser Thr Thr Phe Thr Thr Asn Leu Thr
 385 390 395 400

Glu Tyr Pro Leu Ser Arg Val Asp Leu Gly Asp Cys Ile Gly Lys Asp
 405 410 415

Ala Arg Asp Ala Met Asp Arg Ile Phe Ala Arg Arg Tyr Asn Ala Thr
 420 425 430

His Ile Lys Val Gly Gln Pro Gln Tyr Tyr Leu Ala Asn Gly Gly Phe
 435 440 445

Leu Ile Ala Tyr Gln Pro Leu Leu Ser Asn Thr Leu Ala Glu Leu Tyr
 450 455 460

Val Arg Glu His Leu Arg Glu Gln Ser Arg Lys Pro Pro Asn Pro Thr
 465 470 475 480

Pro Pro Pro Pro Gly Ala Ser Ala Asn Ala Ser Val Glu Arg Ile Lys
 485 490 495

Thr Thr Ser Ser Ile Glu Phe Ala Arg Leu Gln Phe Thr Tyr Asn His
 500 505 510

Ile Gln Arg His Val Asn Asp Met Leu Gly Arg Val Ala Ile Ala Trp
 515 520 525

Cys Glu Leu Gln Asn His Glu Leu Thr Leu Trp Asn Glu Ala Arg Lys
 530 535 540

Leu Asn Pro Asn Ala Ile Ala Ser Ala Thr Val Gly Arg Arg Val Ser
 545 550 555 560

Ala Arg Met Leu Gly Asp Val Met Ala Val Ser Thr Cys Val Pro Val
 565 570 575

Ala Ala Asp Asn Val Ile Val Gln Asn Ser Met Arg Ile Ser Ser Arg
 580 585 590

Pro Gly Ala Cys Tyr Ser Arg Pro Leu Val Ser Phe Arg Tyr Glu Asp
 595 600 605

Gln Gly Pro Leu Val Glu Gly Gln Leu Gly Glu Asn Asn Glu Leu Arg
 610 615 620

Leu Thr Arg Asp Ala Ile Glu Pro Cys Thr Val Gly His Arg Arg Tyr
 625 630 635 640

Phe Thr Phe Gly Gly Gly Tyr Val Tyr Phe Glu Glu Tyr Ala Tyr Ser
 645 650 655

His Gln Leu Ser Arg Ala Asp Ile Thr Thr Val Ser Thr Phe Ile Asp
 660 665 670

Leu Asn Ile Thr Met Leu Glu Asp His Glu Phe Val Pro Leu Glu Val
 675 680 685

-continued

Tyr Thr Arg His Glu Ile Lys Asp Ser Gly Leu Leu Asp Tyr Thr Glu
 690 695 700
 Val Gln Arg Arg Asn Gln Leu His Asp Leu Arg Phe Ala Asp Ile Asp
 705 710 715 720
 Thr Val Ile His Ala Asp Ala Asn Ala Ala Met Phe Ala Gly Leu Gly
 725 730 735
 Ala Phe Phe Glu Gly Met Gly Asp Leu Gly Arg Ala Val Gly Lys Val
 740 745 750
 Val Met Gly Ile Val Gly Gly Val Val Ser Ala Val Ser Gly Val Ser
 755 760 765
 Ser Phe Met Ser Asn Pro Phe Gly Ala Leu Ala Val Gly Leu Leu Val
 770 775 780
 Leu Ala Gly Leu Ala Ala Ala Phe Phe Ala Phe Arg Tyr Val Met Arg
 785 790 795 800
 Leu Gln Ser Asn Pro Met Lys Ala Leu Tyr Pro Leu Thr Thr Lys Glu
 805 810 815
 Leu Lys Asn Pro Thr Asn Pro Asp Ala Ser Gly Glu Gly Glu Glu Gly
 820 825 830
 Gly Asp Phe Asp Glu Ala Lys Leu Ala Glu Ala Arg Glu Met Ile Arg
 835 840 845
 Tyr Met Ala Leu Val Ser Ala Met Glu Arg Thr Glu His Lys Ala Lys
 850 855 860
 Lys Lys Gly Thr Ser Ala Leu Leu Ser Ala Lys Val Thr Asp Met Val
 865 870 875 880
 Met Arg Lys Arg Arg Asn Thr Asn Tyr Thr Gln Val Pro Asn Lys Asp
 885 890 895
 Gly Asp Ala Asp Glu Asp Asp Leu
 900

<210> SEQ ID NO 29
 <211> LENGTH: 1800
 <212> TYPE: DNA
 <213> ORGANISM: Herpes Simplex Virus-1

<400> SEQUENCE: 29

```

aggcgcttg gtcgggaggc cgcacggaac gcacaccccc atccggtggt ccgtgtggag      60
gtcgttttca tgccccgtct cgctttgtcg ggaacgctag ccggtccctc gcgaggggga    120
ggcgtcgggc atggccccgg ggcgggtggg ccttgccgtg gtcctgtgga gcctgttgtg    180
gctcggggcg ggggtggcgg ggggctcgga aactgcctcc accgggcccc cgatcaccgc    240
gggagcgggt acgaacgcga gcgagggccc cacatcgggg tcccccggt cagccgccag    300
cccggaagtc acccccacat cgaccccaaa cccaacaat gtcacacaaa aaaaaaccac    360
ccccaccgag ccggccagcc cccaacaac cccaagccc acctccacgc caaaagccc    420
ccccacgtcc acccccgacc ccaaacccaa gaacaacacc acccccgcca agtcgggccc    480
ccccactaaa cccccgggc ccgtgtggtg cgaccgccc gaccattgg ccgggtacgg    540
ctcgtgggtg cagatccgat gccggttttg gaattccacc cgcattggagt tccgcctcca    600
gatatggcgt tactccatgg gtccgtcccc cccaatcgt ccggtcccc acctagagga    660
ggtcctgacg aacatcaccg ccccacccgg gggactcctg gtgtacgaca gcgcccccaa    720
cctgaecggc ccccacgtgc tctgggcgga gggggccggc ccgggcgccg acctccgtt    780
    
```

-continued

```

gtattctgtc accgggcccg tgcgaccca gggctgatt atcggcgagg tgacgcccgc 840
gaccagggga atgtattact tggcctgggg ccgatggac agcccgcacg agtacgggac 900
gtgggtgcgc gtccgcatgt tccgcccccc gtctctgacc ctccagcccc acgcggtgat 960
ggaggggtcag ccgttcaagg cgacgtgcac ggccgcccgc tactaccgcg gtaaccccgt 1020
ggagtttgtc tggttcgagg acgacgcgca ggtgtttaac ccgggcccaga tcgacacgca 1080
gacgcacgag cacccegcag ggttcaccac agtctctacc gtgacctccg aggctgtcgg 1140
cggccaggtc cccccggga ccttcacctg ccagatgacg tggcaccgcg actccgtgac 1200
gttctcgcca cgcaatgcca ccgggctggc cctggtgctg ccgcgcccaa ccatcaccat 1260
ggaatttggg gtccggcatg tggctgcac ggccggctgc gtccccgagg gcgtagcgtt 1320
tgcctggttc ctgggggacg acccctcacc ggccgctaag tcggccgtta cggcccagga 1380
gtcgtgcgac cacccegggc tggctacggt ccggtccacc ctgcccattt cgtacgacta 1440
cagcgagtac atctgtcggg tgaccggata tccggccggg attcccgttc tagagacca 1500
cggcagtcac cagccccac ccagggacc caccgagcgg caggtgatcg aggcgatcga 1560
gtgggtgggg attggaatcg gggttctcgc ggcgggggtc ctggtcgtaa cggcaatcgt 1620
gtacgtcgtc cgcacatcac agtcgcgcca gcgtcatcgg cgtaacgcg agaccccccc 1680
gttacctttt taatatctat atagtttggg cccccctcta tcccgccac cgctggggcg 1740
tataaagccg ccaccctctc tccctcagg tcctccttgg tcgatcccga acgacacacg 1800

```

<210> SEQ ID NO 30

<211> LENGTH: 511

<212> TYPE: PRT

<213> ORGANISM: Herpes Simplex Virus-1

<400> SEQUENCE: 30

```

Met Ala Pro Gly Arg Val Gly Leu Ala Val Val Leu Trp Gly Leu Leu
1          5          10          15
Trp Leu Gly Ala Gly Val Ala Gly Gly Ser Glu Thr Ala Ser Thr Gly
20          25          30
Pro Thr Ile Thr Ala Gly Ala Val Thr Asn Ala Ser Glu Ala Pro Thr
35          40          45
Ser Gly Ser Pro Gly Ser Ala Ala Ser Pro Glu Val Thr Pro Thr Ser
50          55          60
Thr Pro Asn Pro Asn Asn Val Thr Gln Asn Lys Thr Thr Pro Thr Glu
65          70          75          80
Pro Ala Ser Pro Pro Thr Thr Pro Lys Pro Thr Ser Thr Pro Lys Ser
85          90          95
Pro Pro Thr Ser Thr Pro Asp Pro Lys Pro Lys Asn Asn Thr Thr Pro
100         105         110
Ala Lys Ser Gly Arg Pro Thr Lys Pro Pro Gly Pro Val Trp Cys Asp
115         120         125
Arg Arg Asp Pro Leu Ala Arg Tyr Gly Ser Arg Val Gln Ile Arg Cys
130         135         140
Arg Phe Arg Asn Ser Thr Arg Met Glu Phe Arg Leu Gln Ile Trp Arg
145         150         155         160
Tyr Ser Met Gly Pro Ser Pro Pro Ile Ala Pro Ala Pro Asp Leu Glu
165         170         175
Glu Val Leu Thr Asn Ile Thr Ala Pro Pro Gly Gly Leu Leu Val Tyr

```

-continued

180			185			190									
Asp	Ser	Ala	Pro	Asn	Leu	Thr	Asp	Pro	His	Val	Leu	Trp	Ala	Glu	Gly
	195						200					205			
Ala	Gly	Pro	Gly	Ala	Asp	Pro	Pro	Leu	Tyr	Ser	Val	Thr	Gly	Pro	Leu
	210						215					220			
Pro	Thr	Gln	Arg	Leu	Ile	Ile	Gly	Glu	Val	Thr	Pro	Ala	Thr	Gln	Gly
	225				230					235					240
Met	Tyr	Tyr	Leu	Ala	Trp	Gly	Arg	Met	Asp	Ser	Pro	His	Glu	Tyr	Gly
			245						250					255	
Thr	Trp	Val	Arg	Val	Arg	Met	Phe	Arg	Pro	Pro	Ser	Leu	Thr	Leu	Gln
			260					265					270		
Pro	His	Ala	Val	Met	Glu	Gly	Gln	Pro	Phe	Lys	Ala	Thr	Cys	Thr	Ala
		275					280							285	
Ala	Ala	Tyr	Tyr	Pro	Arg	Asn	Pro	Val	Glu	Phe	Asp	Trp	Phe	Glu	Asp
	290					295					300				
Asp	Arg	Gln	Val	Phe	Asn	Pro	Gly	Gln	Ile	Asp	Thr	Gln	Thr	His	Glu
	305				310					315					320
His	Pro	Asp	Gly	Phe	Thr	Thr	Val	Ser	Thr	Val	Thr	Ser	Glu	Ala	Val
				325					330					335	
Gly	Gly	Gln	Val	Pro	Pro	Arg	Thr	Phe	Thr	Cys	Gln	Met	Thr	Trp	His
			340					345					350		
Arg	Asp	Ser	Val	Thr	Phe	Ser	Arg	Arg	Asn	Ala	Thr	Gly	Leu	Ala	Leu
	355						360					365			
Val	Leu	Pro	Arg	Pro	Thr	Ile	Thr	Met	Glu	Phe	Gly	Val	Arg	His	Val
	370					375					380				
Val	Cys	Thr	Ala	Gly	Cys	Val	Pro	Glu	Gly	Val	Thr	Phe	Ala	Trp	Phe
	385				390					395					400
Leu	Gly	Asp	Asp	Pro	Ser	Pro	Ala	Ala	Lys	Ser	Ala	Val	Thr	Ala	Gln
			405						410					415	
Glu	Ser	Cys	Asp	His	Pro	Gly	Leu	Ala	Thr	Val	Arg	Ser	Thr	Leu	Pro
			420					425					430		
Ile	Ser	Tyr	Asp	Tyr	Ser	Glu	Tyr	Ile	Cys	Arg	Leu	Thr	Gly	Tyr	Pro
	435						440						445		
Ala	Gly	Ile	Pro	Val	Leu	Glu	His	His	Gly	Ser	His	Gln	Pro	Pro	Pro
	450					455					460				
Arg	Asp	Pro	Thr	Glu	Arg	Gln	Val	Ile	Glu	Ala	Ile	Glu	Trp	Val	Gly
	465				470					475					480
Ile	Gly	Ile	Gly	Val	Leu	Ala	Ala	Gly	Val	Leu	Val	Val	Thr	Ala	Ile
				485					490					495	
Val	Tyr	Val	Val	Arg	Thr	Ser	Gln	Ser	Arg	Gln	Arg	His	Arg	Arg	
			500					505					510		

<210> SEQ ID NO 31
 <211> LENGTH: 1282
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 31

aagccacca gcctatgcat ccgctcctca atcctctcct gttggcactg ggctcatgg 60
 cgcttttggt gaccacggtc attgctctca cttgcttgg cggtttgcc tccccaggcc 120
 ctgtgctcc ctctacagcc ctcagggagc tcattgagga gctggtaaac atcaccagga 180

-continued

```

accagaaggc tccgctctgc aatggcagca tggatggag catcaacctg acagctggca 240
tgtactgtgc agccctggaa tccctgatca acgtgtcagg ctgcagtgcc atcgagaaga 300
cccagaggat gctgagcgga ttctgcccgc acaaggtctc agctgggcag tttccagct 360
tgcagtgtcc agacacacaa atcgaggtgg cccagtttgt aaaggacctg ctcttacatt 420
taaagaaact ttttcgagag ggacagttca actgaaactt cgaagcatc attatttgca 480
gagacaggac ctgactattg aagttgcaga ttcatttttc tttctgatgt caaaaatgtc 540
ttgggtaggc ggggaaggagg gttagggagg ggtaaaattc cttagcttag acctcagcct 600
gtgctgcccc tcttcagcct agccgacctc agccttcccc tgcccaggg ctcagcctgg 660
tgggcctcct ctgtccaggg cctgagctc ggtggacca gggatgacat gtcctacac 720
ccctccctg ccctagagca cactgtagca ttacagtggg tgccccctt gccagacatg 780
tggtaggaca gggaccact tcacacacag gcaactgagg cagacagcag ctcaggcaca 840
cttctctctg gtcttattta ttattgtgtg ttatttaaat gagtgtgttt gtcaccgtt 900
gggattgggg aagactgtgg ctgctagcac ttggagcaa gggttcagag actcagggcc 960
ccagcactaa agcagtggac accaggagtc cctggtaata agtactgtgt acagaattct 1020
gtacacctac tggggctcctg gggcctcgga gcctcctcg aggcagggtc aggagagggg 1080
cagaacagcc gctcctgtct gccagccagc agccagctct cagccaacga gtaatttatt 1140
gttttccctt gtatttaaat attaaatag ttagcaaaga gttaatatat agaagggtac 1200
cttgaacct gggggagggg acattgaaca agttgtttca ttgactatca aactgaagcc 1260
agaaataaag ttggtgacag at 1282

```

<210> SEQ ID NO 32

<211> LENGTH: 146

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 32

```

Met His Pro Leu Leu Asn Pro Leu Leu Leu Ala Leu Gly Leu Met Ala
1           5           10          15
Leu Leu Leu Thr Thr Val Ile Ala Leu Thr Cys Leu Gly Gly Phe Ala
20          25          30
Ser Pro Gly Pro Val Pro Pro Ser Thr Ala Leu Arg Glu Leu Ile Glu
35          40          45
Glu Leu Val Asn Ile Thr Gln Asn Gln Lys Ala Pro Leu Cys Asn Gly
50          55          60
Ser Met Val Trp Ser Ile Asn Leu Thr Ala Gly Met Tyr Cys Ala Ala
65          70          75          80
Leu Glu Ser Leu Ile Asn Val Ser Gly Cys Ser Ala Ile Glu Lys Thr
85          90          95
Gln Arg Met Leu Ser Gly Phe Cys Pro His Lys Val Ser Ala Gly Gln
100         105         110
Phe Ser Ser Leu His Val Arg Asp Thr Lys Ile Glu Val Ala Gln Phe
115         120         125
Val Lys Asp Leu Leu Leu His Leu Lys Lys Leu Phe Arg Glu Gly Gln
130         135         140
Phe Asn
145

```

-continued

<210> SEQ ID NO 33
 <211> LENGTH: 1376
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 33

```

gtaagaacac tctcgtgagt ctaacgggtct tccggatgaa ggctatttga agtcgccata      60
acctggtcag aagtgtgcct gtcggcgggg agagaggcaa tatcaagggt ttaaatctcg      120
gagaaatggc tttcgtttgc ttggctatcg gatgcttata tacctttctg ataagcacia      180
catttggtcgt tacttcatct tcagacaccg agataaaaagt taaccctcct caggattttg      240
agatagtgga tccccgatac ttaggttata tctatttga atggcaaccc ccaactgtctc      300
tggatcattt taaggaatgc acagtggaat atgaactaaa ataccgaaac attggtagtg      360
aaacatggaa gaccatcatt actaagaatc tacattacaa agatgggttt gatcttaaca      420
agggcattga agcgaagata cacacgcttt taccatggca atgcacaaat ggatcagaag      480
ttcaaagtcc ctgggcagaa actacttatt ggatatcacc acaaggaatt ccagaaacta      540
aagttcagga tatggattgc gtatattaca attggcaata tttactctgt tcttggaaac      600
ctggcatagg tgtacttctt gataccaatt acaacttgtt ttactggtat gagggcttgg      660
atcatgcatt acagtgtggt gattacatca aggctgatgg acaaaatata ggatgcagat      720
ttccctattt ggagcatca gactataaag atttctatat ttgtgttaat ggatcatcag      780
agaacaagcc tatcagatcc agttatttca cttttcagct tcaaaatata gttaaaacct      840
tgccgccagt ctatcttact tttactcggg agagttcatg tgaattaag ctgaaatgga      900
gcataccttt gggacatatt ccagcaaggt gttttgatta tgaattgag atcagagaag      960
atgatactac cttggtgact gctacagttg aaaatgaaac atacaccttg aaaacaacia     1020
atgaaacccg acaattatgc tttgtagtaa gaagcaaagt gaatatttat tgctcagatg     1080
acggaatttg gagtgagtgg agtgataaac aatgctggga aggtgaagac ctatcgaaga     1140
aaactttgct acgtttctgg ctaccatttg gtttcactct aatattagtt atatttgtaa     1200
ccggtctgct tttgcgtaag ccaaacacct acccaaaaat gattccagaa tttttctgtg     1260
atacatgaag actttccata tcaagagaca tggatttgac tcaacagttt ccagtcgatg     1320
ccaaatgttc aatatgagtc tcaataaaact gaatttttct tgcgaatggt gaaaaa      1376

```

<210> SEQ ID NO 34
 <211> LENGTH: 380
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 34

```

Met Ala Phe Val Cys Leu Ala Ile Gly Cys Leu Tyr Thr Phe Leu Ile
1           5           10           15
Ser Thr Thr Phe Gly Cys Thr Ser Ser Ser Asp Thr Glu Ile Lys Val
20          25          30
Asn Pro Pro Gln Asp Phe Glu Ile Val Asp Pro Gly Tyr Leu Gly Tyr
35          40          45
Leu Tyr Leu Gln Trp Gln Pro Pro Leu Ser Leu Asp His Phe Lys Glu
50          55          60
Cys Thr Val Glu Tyr Glu Leu Lys Tyr Arg Asn Ile Gly Ser Glu Thr
65          70          75          80

```


-continued

gcaaactttt acgtgtgccc acccccacg ggcgccacgg tggatgcagtt cgagcagccg	360
cgccgctgcc cgaccggccc cgagggtcag aactacacgg agggcatcgc ggtggtcttc	420
aaggagaaca tcgccccgta caagttcaag gccaccatgt actacaaaga cgtcaccggt	480
tcgcagggtg ggttcggcca ccgctactcc cagtttatgg ggatcttga ggaccgccc	540
cccgccccct tcgaggaggt gatcgacaag atcaacgcca agggggtctg tcggtccacg	600
gccaaagtaag tcgcaacaaa cctggagacc acccggttc accgggacga ccacgagacc	660
gacatggagc taaaaccggc caaccccgcg acccgcacga gccggggctg gcacaccacc	720
gacctcaagt acaaccctc gcgggtggag gcgttccacc ggtacgggac gacggtaaac	780
tgatcgtcg aggaggtgga cgcgcgctcg gtgtaccctg acgacgagtt tgtgttgccg	840
actggcgact ttgtgtacat gtccccgttt tacggctacc gggaggggtc gcacaccgaa	900
cacaccagct acgcccgcga ccgcttcaag caggtcgacg gcttctacgc gcgacacctc	960
accaccaagg cccggggccc ggcgcgac acccggaacc tgctcacgac ccccaagttc	1020
accgtggcct gggactgggt gccaaagcgc ccgtcggctc gcaccatgac caagtggcag	1080
gaggtggagc agatgctgcg ctccgagtac ggcggctcct tccgattctc ttcgacgcc	1140
atatccacca ccttcaccac caacctgacc gagtaccgc tctcgcgctg ggacctgggg	1200
gactgcatcg gcaaggacgc ccgcgacgcc atggaccgca tcttcgcccg caggtacaac	1260
gcgacgcaca tcaaggtggg ccagcgcgag tactacctgg ccaatggggg ctttctgatc	1320
gcgtaccagc cccttctcag caacaogctc gcggagctgt acgtgcggga acacctccgc	1380
gagcagagcc gcaagcccc aaacccccag cccccgcgc ccggggccag cgccaacgcg	1440
tccgtggagc gcatcaagac cacctcctcc atcgagttcg ccaggctgca gtttacgtac	1500
aaccacatac agcgcctatg caacgatatg ttgggcccgc ttgccatcgc gtggtgcgag	1560
ctgcagaatc acgagctgac cctgtggaac gaggcccga agctgaacc caacgccatc	1620
gcctcgcca ccgtgggccc gcgggtgagc gcgcggatgc tcggcgacgt gatggccgctc	1680
tccacgtgag tcgggtgccc cgcggacaac gtgatcgtcc aaaactcagc gcgcatcagc	1740
tcgggcccc gggcctgcta cagccgcccc ctggtcagct ttcggtaaga agaccagggc	1800
ccgttggtcg aggggagct gggggagaac aacgagctgc ggctgacgcg cgatgcgac	1860
gagccgtgca ccgtgggaca ccggcgctac ttcacctcgc gtgggggcta cgtgtacttc	1920
gaggagtacg cgtactccca ccagctgagc cgcgccgaca tcaccacctg cagcaccttc	1980
atcgacctca acatcaccat gctggaggat cacgagtttg tccccctgga ggtgtacacc	2040
cgccacgaga tcaaggacag ccggcctgctg gactacacgg aggtccagcg ccgcaaccag	2100
ctgcacgacc tcgcttccgc cgacatcgac acggtcatcc acgccgacgc caacgccgcc	2160
atgtttgccc gcctgggggc gttcttcgag gggatgggccc acctggggcg ccgggtcggc	2220
aagggtggtg tgggcatcgt gggcgccgctg gtatcggccc tgctgggctg gtcctccttc	2280
atgtccaacc cctttggggc gctggccgctg ggtctgttgg tcctggcccg cctggcggcg	2340
gccttcttcg cctttcgtca cgtcatgccc ctgcagagca accccatgaa ggcctgtac	2400
ccgtaacca ccaaggagct caagaacccc accaaccggc acgcgtccgg ggaggcgag	2460
gaggcgccgc actttgacga ggccaagcta gccgagccc gggagatgat acggtacatg	2520
gccttggtgt ctgcatgga gcgcacggaa cacaaggcca agaagaaggg cacgagcgcg	2580

-continued

```
ctgctcagcg ccaaggtcac cgacatggtc atgcgcaagc gccgcaacac caactacacc 2640
caagttccca acaaagacgg tgacgccgac gaggacgacc tgtga 2685
```

```
<210> SEQ ID NO 36
<211> LENGTH: 894
<212> TYPE: PRT
<213> ORGANISM: Herpes Simplex Virus-1
```

```
<400> SEQUENCE: 36
```

```
Met Arg Gln Gly Ala Pro Ala Arg Gly Arg Arg Trp Phe Val Val Trp
 1          5          10          15
Ala Leu Leu Gly Leu Thr Leu Gly Val Leu Val Ala Ser Ala Ala Pro
 20          25          30
Ser Ser Pro Gly Thr Pro Gly Val Ala Ala Ala Thr Gln Ala Ala Asn
 35          40          45
Gly Gly Pro Ala Thr Pro Ala Pro Pro Ala Pro Gly Ala Pro Pro Thr
 50          55          60
Gly Asp Pro Pro Lys Pro Pro Arg Pro Ala Gly Asp Asn Ala Thr Val
 65          70          75          80
Ala Ala Gly His Ala Thr Leu Arg Glu His Leu Arg Asp Ile Lys Ala
 85          90          95
Glu Asn Thr Asp Ala Asn Phe Tyr Val Cys Pro Pro Pro Thr Gly Ala
 100         105         110
Thr Val Val Gln Phe Glu Gln Pro Arg Arg Cys Pro Thr Arg Pro Glu
 115         120         125
Gly Gln Asn Tyr Thr Glu Gly Ile Ala Val Val Phe Lys Glu Asn Ile
 130         135         140
Ala Pro Tyr Lys Phe Lys Ala Thr Met Tyr Tyr Lys Asp Val Thr Val
 145         150         155         160
Ser Gln Val Trp Phe Gly His Arg Tyr Ser Gln Phe Met Gly Ile Phe
 165         170         175
Glu Asp Arg Ala Pro Val Pro Phe Glu Glu Val Ile Asp Lys Ile Asn
 180         185         190
Ala Lys Gly Val Cys Arg Ser Thr Ala Lys Tyr Val Arg Asn Asn Leu
 195         200         205
Glu Thr Thr Ala Phe His Arg Asp Asp His Glu Thr Asp Met Glu Leu
 210         215         220
Lys Pro Ala Asn Ala Ala Thr Arg Thr Ser Arg Gly Trp His Thr Thr
 225         230         235         240
Asp Leu Lys Tyr Asn Pro Ser Arg Val Glu Ala Phe His Arg Tyr Gly
 245         250         255
Thr Thr Val Asn Cys Ile Val Glu Glu Val Asp Ala Arg Ser Val Tyr
 260         265         270
Pro Tyr Asp Glu Phe Val Leu Ala Thr Gly Asp Phe Val Tyr Met Ser
 275         280         285
Pro Phe Tyr Gly Tyr Arg Glu Gly Ser His Thr Glu His Thr Ser Tyr
 290         295         300
Ala Ala Asp Arg Phe Lys Gln Val Asp Gly Phe Tyr Ala Arg Asp Leu
 305         310         315         320
Thr Thr Lys Ala Arg Ala Thr Ala Pro Thr Thr Arg Asn Leu Leu Thr
 325         330         335
Thr Pro Lys Phe Thr Val Ala Trp Asp Trp Val Pro Lys Arg Pro Ser
```

-continued

340				345				350							
Val	Cys	Thr	Met	Thr	Lys	Trp	Gln	Glu	Val	Asp	Glu	Met	Leu	Arg	Ser
	355						360					365			
Glu	Tyr	Gly	Gly	Ser	Phe	Arg	Phe	Ser	Ser	Asp	Ala	Ile	Ser	Thr	Thr
	370					375					380				
Phe	Thr	Thr	Asn	Leu	Thr	Glu	Tyr	Pro	Leu	Ser	Arg	Val	Asp	Leu	Gly
385					390					395					400
Asp	Cys	Ile	Gly	Lys	Asp	Ala	Arg	Asp	Ala	Met	Asp	Arg	Ile	Phe	Ala
			405						410					415	
Arg	Arg	Tyr	Asn	Ala	Thr	His	Ile	Lys	Val	Gly	Gln	Pro	Gln	Tyr	Tyr
		420						425					430		
Leu	Ala	Asn	Gly	Gly	Phe	Leu	Ile	Ala	Tyr	Gln	Pro	Leu	Leu	Ser	Asn
	435					440						445			
Thr	Leu	Ala	Glu	Leu	Tyr	Val	Arg	Glu	His	Leu	Arg	Glu	Gln	Ser	Arg
	450					455					460				
Lys	Pro	Pro	Asn	Pro	Thr	Pro	Pro	Pro	Pro	Gly	Ala	Ser	Ala	Asn	Ala
465					470					475					480
Ser	Val	Glu	Arg	Ile	Lys	Thr	Thr	Ser	Ser	Ile	Glu	Phe	Ala	Arg	Leu
			485						490					495	
Gln	Phe	Thr	Tyr	Asn	His	Ile	Gln	Arg	His	Val	Asn	Asp	Met	Leu	Gly
		500						505					510		
Arg	Val	Ala	Ile	Ala	Trp	Cys	Glu	Leu	Gln	Asn	His	Glu	Leu	Thr	Leu
		515					520					525			
Trp	Asn	Glu	Ala	Arg	Lys	Leu	Asn	Pro	Asn	Ala	Ile	Ala	Ser	Ala	Thr
	530					535					540				
Val	Gly	Arg	Arg	Val	Ser	Ala	Arg	Met	Leu	Gly	Asp	Val	Met	Ala	Val
545					550					555					560
Ser	Thr	Cys	Val	Pro	Val	Ala	Ala	Asp	Asn	Val	Ile	Val	Gln	Asn	Ser
			565						570					575	
Met	Arg	Ile	Ser	Ser	Arg	Pro	Gly	Ala	Cys	Tyr	Ser	Arg	Pro	Leu	Val
		580						585					590		
Ser	Phe	Arg	Tyr	Glu	Asp	Gln	Gly	Pro	Leu	Val	Glu	Gly	Gln	Leu	Gly
		595					600					605			
Glu	Asn	Asn	Glu	Leu	Arg	Leu	Thr	Arg	Asp	Ala	Ile	Glu	Pro	Cys	Thr
	610					615						620			
Val	Gly	His	Arg	Arg	Tyr	Phe	Thr	Phe	Gly	Gly	Gly	Tyr	Val	Tyr	Phe
625					630					635					640
Glu	Glu	Tyr	Ala	Tyr	Ser	His	Gln	Leu	Ser	Arg	Ala	Asp	Ile	Thr	Thr
			645						650					655	
Val	Ser	Thr	Phe	Ile	Asp	Leu	Asn	Ile	Thr	Met	Leu	Glu	Asp	His	Glu
		660						665					670		
Phe	Val	Pro	Leu	Glu	Val	Tyr	Thr	Arg	His	Glu	Ile	Lys	Asp	Ser	Gly
		675					680					685			
Leu	Leu	Asp	Tyr	Thr	Glu	Val	Gln	Arg	Arg	Asn	Gln	Leu	His	Asp	Leu
	690					695					700				
Arg	Phe	Ala	Asp	Ile	Asp	Thr	Val	Ile	His	Ala	Asp	Ala	Asn	Ala	Ala
705					710					715					720
Met	Phe	Ala	Gly	Leu	Gly	Ala	Phe	Phe	Glu	Gly	Met	Gly	Asp	Leu	Gly
			725						730					735	
Arg	Ala	Val	Gly	Lys	Val	Val	Met	Gly	Ile	Val	Gly	Gly	Val	Val	Ser
			740					745					750		

-continued

Ala Val Ser Gly Val Ser Ser Phe Met Ser Asn Pro Phe Gly Ala Leu
755 760 765

Ala Val Gly Leu Leu Val Leu Ala Gly Leu Ala Ala Ala Phe Phe Ala
770 775 780

Phe Arg Tyr Val Met Arg Leu Gln Ser Asn Pro Met Lys Ala Leu Tyr
785 790 795 800

Pro Leu Thr Thr Lys Glu Leu Lys Asn Pro Thr Asn Pro Asp Ala Ser
805 810 815

Gly Glu Gly Glu Glu Gly Gly Asp Phe Asp Glu Ala Lys Leu Ala Glu
820 825 830

Ala Arg Glu Met Ile Arg Tyr Met Ala Leu Val Ser Ala Met Glu Arg
835 840 845

Thr Glu His Lys Ala Lys Lys Lys Gly Thr Ser Ala Leu Leu Ser Ala
850 855 860

Lys Val Thr Asp Met Val Met Arg Lys Arg Arg Asn Thr Asn Tyr Thr
865 870 875 880

Gln Val Pro Asn Lys Asp Gly Asp Ala Asp Glu Asp Asp Leu
885 890

<210> SEQ ID NO 37

<211> LENGTH: 1626

<212> TYPE: DNA

<213> ORGANISM: Herpes Simplex Virus-1

<400> SEQUENCE: 37

```

atggcgcttt tgttgaccac ggtcattgct ctcaactgcc ttggcggtt tgcctcccca    60
ggccctgtgc ctcccctac agccctcagg gagctcattg aggagctggt caacatcacc    120
cagaaccaga aggtccgct ctgcaatggc agcatggtat ggagcatcaa cctgacagct    180
ggcatgtact gtgcagccct ggaatccctg atcaacgtgt caggctgcag tgccatcgag    240
aagaccaga g gatgctgag cggattctgc cgcacaagg tctcagctgg gcagttttcc    300
agcttgcagt tccgagacac caaaatcgag gtggcccagt ttgtaaagga cctgctctta    360
catttaaga aactttttcg cgagggacag ttcaacgaat tccaccgca tggagtccg    420
ctccagata tggcgtaact ccattgggtcc gtcccccca atcgtccgg ctcccgaact    480
agaggaggtc ctgacgaaca tcaccgccc acccggggga ctctggtgt acgacagcg    540
ccccacctg acggaccccc acgtgctctg ggccggagggg gccggcccgg gcgccgacc    600
tccgttgat tctgtaccg gcccgctgcc gaccagcgg ctgattatcg gcgagtgac    660
gcccgcgacc cagggaatgt attacttggc ctggggccgg atggacagcc cgcacgagta    720
cgggacgtgg gtgcgcgtcc gcatgttccg cccccgtct ctgaccctcc agccccacgc    780
ggtgatggag ggtcacccgt tcaaggcgac gtgcacggcc gccgcctact acccgctaa    840
ccccgtggag tttgtctggt tcgaggacga ccgccaggtg ttaaccgg gccagatcga    900
cacgcagacg cacgagcacc ccgacgggtt caccacagtc tctaccgtga cctccgaggc    960
tgtcggcggc caggtccccc cgcggactt cacctgccag atgacgtggc accgcgactc   1020
cgtgacgttc tcgcgacgca atgccaccg gctggccctg gtgctgccc gcaccaaccat   1080
caccatgaa tttgggtgcc ggcattggtt ctgcacggcc ggctgcgtcc ccgagggcgt   1140
gacgtttgcc tggttctcgg gggacgaccc ctcaccggcg gctaagtccg ccgttacggc   1200

```

-continued

```

ccaggagtcg tgcgaccacc ccgggctggc tacggctcgg tccacctgc ccatttcgta 1260
cgactacagc gactacatct gtcggttgac cggatatecg gccgggattc ccgttctaga 1320
gcaccacggc agtcaccagc ccccaccag ggaccccacc gagcggcagg tgatcgagge 1380
gatcgagtgg gtggggattg gaatcggggg tctcgcggcg ggggtcctgg tcgtaacggc 1440
aatcgtgtac gtcgtccgca catcacagtc gcggcagcgt catcggcggg aacgcgagac 1500
ccccccgta cctttttaat atctatatag tttggteccc cctctatccc gccaccgct 1560
ggggcgtata aagccgccac cctctcttcc ctcaggtcat ccttggtcga tcccgaacga 1620
cacacg 1626

```

<210> SEQ ID NO 38

<211> LENGTH: 497

<212> TYPE: PRT

<213> ORGANISM: Herpes Simplex Virus-1

<400> SEQUENCE: 38

```

Met Ala Leu Leu Leu Thr Thr Val Ile Ala Leu Thr Cys Leu Gly Gly
 1          5          10          15
Phe Ala Ser Pro Gly Pro Val Pro Pro Ser Thr Ala Leu Arg Glu Leu
 20          25          30
Ile Glu Glu Leu Val Asn Ile Thr Gln Asn Gln Lys Ala Pro Leu Cys
 35          40          45
Asn Gly Ser Met Val Trp Ser Ile Asn Leu Thr Ala Gly Met Tyr Cys
 50          55          60
Ala Ala Leu Glu Ser Leu Ile Asn Val Ser Gly Cys Ser Ala Ile Glu
 65          70          75          80
Lys Thr Gln Arg Met Leu Ser Gly Phe Cys Pro His Lys Val Ser Ala
 85          90          95
Gly Gln Phe Ser Ser Leu His Val Arg Asp Thr Lys Ile Glu Val Ala
 100         105         110
Gln Phe Val Lys Asp Leu Leu Leu His Leu Lys Lys Leu Phe Arg Glu
 115         120         125
Gly Gln Phe Asn Glu Phe Ser Thr Arg Met Glu Phe Arg Leu Gln Ile
 130         135         140
Trp Arg Tyr Ser Met Gly Pro Ser Pro Pro Ile Ala Pro Ala Pro Asp
 145         150         155         160
Leu Glu Glu Val Leu Thr Asn Ile Thr Ala Pro Pro Gly Gly Leu Leu
 165         170         175
Val Tyr Asp Ser Ala Pro Asn Leu Thr Asp Pro His Val Leu Trp Ala
 180         185         190
Glu Gly Ala Gly Pro Gly Ala Asp Pro Pro Leu Tyr Ser Val Thr Gly
 195         200         205
Pro Leu Pro Thr Gln Arg Leu Ile Ile Gly Glu Val Thr Pro Ala Thr
 210         215         220
Gln Gly Met Tyr Tyr Leu Ala Trp Gly Arg Met Asp Ser Pro His Glu
 225         230         235         240
Tyr Gly Thr Trp Val Arg Val Arg Met Phe Arg Pro Pro Ser Leu Thr
 245         250         255
Leu Gln Pro His Ala Val Met Glu Gly Gln Pro Phe Lys Ala Thr Cys
 260         265         270
Thr Ala Ala Ala Tyr Tyr Pro Arg Asn Pro Val Glu Phe Asp Trp Phe

```

-continued

275			280			285									
Glu	Asp	Asp	Arg	Gln	Val	Phe	Asn	Pro	Gly	Gln	Ile	Asp	Thr	Gln	Thr
290						295					300				
His	Glu	His	Pro	Asp	Gly	Phe	Thr	Thr	Val	Ser	Thr	Val	Thr	Ser	Glu
305					310					315					320
Ala	Val	Gly	Gly	Gln	Val	Pro	Pro	Arg	Thr	Phe	Thr	Cys	Gln	Met	Thr
				325					330					335	
Trp	His	Arg	Asp	Ser	Val	Thr	Phe	Ser	Arg	Arg	Asn	Ala	Thr	Gly	Leu
			340					345					350		
Ala	Leu	Val	Leu	Pro	Arg	Pro	Thr	Ile	Thr	Met	Glu	Phe	Gly	Val	Arg
	355						360					365			
His	Val	Val	Cys	Thr	Ala	Gly	Cys	Val	Pro	Glu	Gly	Val	Thr	Phe	Ala
370						375					380				
Trp	Phe	Leu	Gly	Asp	Asp	Pro	Ser	Pro	Ala	Ala	Lys	Ser	Ala	Val	Thr
385				390						395					400
Ala	Gln	Glu	Ser	Cys	Asp	His	Pro	Gly	Leu	Ala	Thr	Val	Arg	Ser	Thr
				405					410					415	
Leu	Pro	Ile	Ser	Tyr	Asp	Tyr	Ser	Glu	Tyr	Ile	Cys	Arg	Leu	Thr	Gly
			420					425					430		
Tyr	Pro	Ala	Gly	Ile	Pro	Val	Leu	Glu	His	His	Gly	Ser	His	Gln	Pro
	435						440				445				
Pro	Pro	Arg	Asp	Pro	Thr	Glu	Arg	Gln	Val	Ile	Glu	Ala	Ile	Glu	Trp
	450					455					460				
Val	Gly	Ile	Gly	Ile	Gly	Val	Leu	Ala	Ala	Gly	Val	Leu	Val	Val	Thr
465					470					475					480
Ala	Ile	Val	Tyr	Val	Val	Arg	Thr	Ser	Gln	Ser	Arg	Gln	Arg	His	Arg
				485					490					495	

Arg

<210> SEQ ID NO 39
 <211> LENGTH: 1593
 <212> TYPE: DNA
 <213> ORGANISM: Herpes Simplex Virus-1

<400> SEQUENCE: 39

```

atgggggggg ctgccgccag gttgggggcc gtgattttgt ttgtcgtcat agtgggcctc    60
catgggggtcc ggggcaataa tgccttgccg gatgcctctc tcaagatggc cgacccaat    120
cgcttttcgcg gcaaagacct tccggtcctc gagatggcgc ttttggtgac cacggtcatt    180
gctctcaactt gccttgccgg ctttgctccc ccaggccctg tgctccctc tacagccctc    240
agggagctca ttgaggagct ggtcaacatc acccagaacc agaaggctcc gctctgcaat    300
ggcagcatgg tatggagcat caactgaca gctggcatgt actgtgcagc cctggaatcc    360
ctgatcaacg tgtcaggctg cagtgccatc gagaagacc agaggatgct gagcggattc    420
tgcccgcaca aggtctcagc tgggcagttt tccagcttgc atgtccgaga caccaaaatc    480
gaggtggccc agtttgtaaa ggacctgctc ttacatttaa agaaactttt tcgcgaggga    540
cagttcaacg gtacctgga ccggtgacc gacctccgg gggtcggcg cgtgtaccac    600
atccaggcgg gcctaccgga cccgttccag cccccagcc tcccgatcac ggtttactac    660
gccgtgttgg agcgcgctg ccgcagcgtg ctctaaacg caccgtcgga ggccccccag    720
attgtccgcg ggcctccga agacgtccgg aaacaacctt acaacctgac catcgcttgg    780
    
```

-continued

```

tttggatgg gaggaactg tgctatcccc atcacggtca tggagtacac cgaatgctcc 840
tacaacaagt ctctgggggc ctgtcccatc cgaacgcagc cccgctggaa ctactatgac 900
agcttcagcg ccgtcagcga ggataacctg gggttcctga tgcacgcccc cgcgtttgag 960
accgccggca cgtacctgcg gctcgtgaag ataaacgact ggacggagat tacacagttt 1020
atcctggagc accgagccaa gggctcctgt aagtacgccc ttccgctgcg catccccccg 1080
tcagcctgcc tctcccccca ggcctaccag cagggggtga cggtgacag catcgggatg 1140
ctgccccgct tcatccccga gaaccagcgc accgtcgcgc tatacagctt gaagatcgcc 1200
gggtggcacg ggcccaaggc cccatacacg agcacctgc tgcccccgga gctgtccgag 1260
acccccaaag ccacgcagcc agaactcgcc cgggaagacc cggaggattc ggcctcttg 1320
gaggaccccc tggggacggt ggtgcccga atccccaaa actggacat accgtcgatc 1380
caggacgccc cgacgcctta ccatccccg gccaccccg acaacatggg cctgatcgcc 1440
ggcgcggtgg ggcgcagtct cctggtagcc ctggtcattt gcggaattgt gtactggatg 1500
cgccgcgcga ctcaaaaagc cccaaagcgc atacgcctcc cccacatccg ggaagacgac 1560
cagccgtcct cgcaccagcc cttgttttac tag 1593

```

<210> SEQ ID NO 40

<211> LENGTH: 530

<212> TYPE: PRT

<213> ORGANISM: Herpes Simplex Virus-1

<400> SEQUENCE: 40

```

Met Gly Gly Ala Ala Ala Arg Leu Gly Ala Val Ile Leu Phe Val Val
1           5           10          15
Ile Val Gly Leu His Gly Val Arg Gly Lys Tyr Ala Leu Ala Asp Ala
20          25          30
Ser Leu Lys Met Ala Asp Pro Asn Arg Phe Arg Gly Lys Asp Leu Pro
35          40          45
Val Leu Glu Met Ala Leu Leu Leu Thr Thr Val Ile Ala Leu Thr Cys
50          55          60
Leu Gly Gly Phe Ala Ser Pro Gly Pro Val Pro Pro Ser Thr Ala Leu
65          70          75          80
Arg Glu Leu Ile Glu Glu Leu Val Asn Ile Thr Gln Asn Gln Lys Ala
85          90          95
Pro Leu Cys Asn Gly Ser Met Val Trp Ser Ile Asn Leu Thr Ala Gly
100         105         110
Met Tyr Cys Ala Ala Leu Glu Ser Leu Ile Asn Val Ser Gly Cys Ser
115        120        125
Ala Ile Glu Lys Thr Gln Arg Met Leu Ser Gly Phe Cys Pro His Lys
130        135        140
Val Ser Ala Gly Gln Phe Ser Ser Leu His Val Arg Asp Thr Lys Ile
145        150        155        160
Glu Val Ala Gln Phe Val Lys Asp Leu Leu Leu His Leu Lys Lys Leu
165        170        175
Phe Arg Glu Gly Gln Phe Asn Gly Thr Pro Asp Arg Leu Thr Asp Pro
180        185        190
Pro Gly Val Arg Arg Val Tyr His Ile Gln Ala Gly Leu Pro Asp Pro
195        200        205

```

-continued

Phe Gln Pro Pro Ser Leu Pro Ile Thr Val Tyr Tyr Ala Val Leu Glu
 210 215 220

Arg Ala Cys Arg Ser Val Leu Leu Asn Ala Pro Ser Glu Ala Pro Gln
 225 230 235 240

Ile Val Arg Gly Ala Ser Glu Asp Val Arg Lys Gln Pro Tyr Asn Leu
 245 250 255

Thr Ile Ala Trp Phe Arg Met Gly Gly Asn Cys Ala Ile Pro Ile Thr
 260 265 270

Val Met Glu Tyr Thr Glu Cys Ser Tyr Asn Lys Ser Leu Gly Ala Cys
 275 280 285

Pro Ile Arg Thr Gln Pro Arg Trp Asn Tyr Tyr Asp Ser Phe Ser Ala
 290 295 300

Val Ser Glu Asp Asn Leu Gly Phe Leu Met His Ala Pro Ala Phe Glu
 305 310 315 320

Thr Ala Gly Thr Tyr Leu Arg Leu Val Lys Ile Asn Asp Trp Thr Glu
 325 330 335

Ile Thr Gln Phe Ile Leu Glu His Arg Ala Lys Gly Ser Cys Lys Tyr
 340 345 350

Ala Leu Pro Leu Arg Ile Pro Pro Ser Ala Cys Leu Ser Pro Gln Ala
 355 360 365

Tyr Gln Gln Gly Val Thr Val Asp Ser Ile Gly Met Leu Pro Arg Phe
 370 375 380

Ile Pro Glu Asn Gln Arg Thr Val Ala Val Tyr Ser Leu Lys Ile Ala
 385 390 395 400

Gly Trp His Gly Pro Lys Ala Pro Tyr Thr Ser Thr Leu Leu Pro Pro
 405 410 415

Glu Leu Ser Glu Thr Pro Asn Ala Thr Gln Pro Glu Leu Ala Pro Glu
 420 425 430

Asp Pro Glu Asp Ser Ala Leu Leu Glu Asp Pro Val Gly Thr Val Val
 435 440 445

Pro Gln Ile Pro Pro Asn Trp His Ile Pro Ser Ile Gln Asp Ala Ala
 450 455 460

Thr Pro Tyr His Pro Pro Ala Thr Pro Asn Asn Met Gly Leu Ile Ala
 465 470 475 480

Gly Ala Val Gly Gly Ser Leu Leu Val Ala Leu Val Ile Cys Gly Ile
 485 490 495

Val Tyr Trp Met Arg Arg Arg Thr Gln Lys Ala Pro Lys Arg Ile Arg
 500 505 510

Leu Pro His Ile Arg Glu Asp Asp Gln Pro Ser Ser His Gln Pro Leu
 515 520 525

Phe Tyr
 530

<210> SEQ ID NO 41
 <211> LENGTH: 1416
 <212> TYPE: DNA
 <213> ORGANISM: Herpes Simplex Virus-1
 <400> SEQUENCE: 41

atggcgcttt tgttgaccac ggtcattgct ctcacttgcc ttggcggcctt tgctccccea 60
 ggcctgtgc ctccctctac agccctcagg gagctcattg aggagctggt caacatcacc 120
 cagaaccaga aggctccgct ctgcaatggc agcatggtat ggagcatcaa cctgacagct 180

-continued

```

ggcatgtact gtgcagccct ggaatocctg atcaacgtgt caggctgcag tgccatcgag 240
aagaccaga ggatgctgag cggattctgc cgcacaagg tctcagctgg gcagttttcc 300
agcttgcagt tccgagacac caaaatcgag gtggcccagt ttgtaaagga cctgctctta 360
catttaaaga aactttttcg cgagggacag ttcaacgta ccgggtcccg gcgcgtgtac 420
cacatccagg cgggcctacc ggaccogtgc cagccccca gcctcccgat cacggtttac 480
tacgccgtgt tggagcgcgc ctgccgcagc gtgctcctaa acgcaccgtc ggaggccccc 540
cagattgtcc gcggggcctc cgaagacgtc cggaaacaac cctacaacct gaccatcgct 600
tggtttcgga tgggaggcaa ctgtgctatc cccatcacgg tcatggagta caccgaatgc 660
tcctacaaca agtctctggg gccctgtccc atccgaacgc agccccgctg gaactactat 720
gacagcttca gcgccgtcag cgaggataac ctggggttcc tgatgcacgc ccccgcttt 780
gagaccgccg gcacgtacct gcggctcgtg aagataaacg actggacgga gattacacag 840
tttatcctgg agcaccgagc caagggtccc tgtaagtacg cccttccgct gcgcatcccc 900
ccgtcagcct gcctctcccc ccaggcctac cagcaggggg tgacgggtgga cagcatcggg 960
atgtgcccc gtttcatccc cgagaaccag cgcaccgtcg ccgtatacag cttgaagatc 1020
gccgggtggc acggggccca gcccctac acgagcacc tgctgcccc ggagctgtcc 1080
gagaccccc acgccacgca gccagaactc gcccgggaag accccgagga ttcggccctc 1140
ttggaggacc ccgtggggac ggtggtgccg caaatcccac caaactggca cataccgtcg 1200
atccaggacg ccgcgacgcc ttaccatccc ccggccccc cgaacaacat gggcctgatc 1260
gccggcggg tggggcgag tctcctggta gccctggta tttcggaat tgtgtactgg 1320
atgcccggcc gactcaaaa agccccaaag cgcatacggc tccccacat ccggaagac 1380
gaccagccgt cctcgacca gccctgttt tactag 1416

```

<210> SEQ ID NO 42

<211> LENGTH: 471

<212> TYPE: PRT

<213> ORGANISM: Herpes Simplex Virus-1

<400> SEQUENCE: 42

```

Met Ala Leu Leu Leu Thr Thr Val Ile Ala Leu Thr Cys Leu Gly Gly
1           5           10          15
Phe Ala Ser Pro Gly Pro Val Pro Pro Ser Thr Ala Leu Arg Glu Leu
20          25          30
Ile Glu Glu Leu Val Asn Ile Thr Gln Asn Gln Lys Ala Pro Leu Cys
35          40          45
Asn Gly Ser Met Val Trp Ser Ile Asn Leu Thr Ala Gly Met Tyr Cys
50          55          60
Ala Ala Leu Glu Ser Leu Ile Asn Val Ser Gly Cys Ser Ala Ile Glu
65          70          75          80
Lys Thr Gln Arg Met Leu Ser Gly Phe Cys Pro His Lys Val Ser Ala
85          90          95
Gly Gln Phe Ser Ser Leu His Val Arg Asp Thr Lys Ile Glu Val Ala
100         105         110
Gln Phe Val Lys Asp Leu Leu Leu His Leu Lys Lys Leu Phe Arg Glu
115         120         125
Gly Gln Phe Asn Gly Thr Gly Ser Arg Arg Val Tyr His Ile Gln Ala

```


-continued

```

cagaaccaga aggctccgct ctgcaatggc agcatggtat ggagcatcaa cctgacagct 180
ggcatgtact gtgcagccct ggaatccctg atcaacgtgt caggetgcag tgccatcgag 240
aagaccacaga g gatgctgag cggattctgc cgcacaagg tctcagctgg gcagttttcc 300
agcttgcagt tccgagacac caaaatcgag gtggcccagt ttgtaaagga cctgctctta 360
catttaaaga aactttttcg cgagggacag ttcaacggta cgggggtccg gcgctcgtac 420
cacatccagg cgggcttacc ggaccggttc cagcccccca gcctcccgat cagggtttac 480
taecccgtgt tggagcgcgc ctgccgcagc gtgctcctaa acgcaccgtc ggaggccccc 540
cagattgtcc gggggcctc cgaagacgtc cggaaacaac cctacaacct gaccatcgct 600
tggtttcgga tgggaggcaa ctgtgctatc cccatcacgg tcatggagta caccgaatgc 660
tctacaaca agtctctggg ggcctgtccc atccgaacgc agccccgtg gaactactat 720
gacagettca gcgccgtcag cgaggataac ctggggttcc tgatgcaagc ccccgcttt 780
gagaccgccc gcacgtacct gcggctcgtg aagataaacg actggacgga gattacacag 840
tttatcctgg agcaccgagc caagggtccc tgtaagtacg ccttccgct gcgcatcccc 900
ccgtcagcct gcctctcccc ccaggcctac cagcaggggg tgacggtgga cagcatcggg 960
atgctgcccc gtttcatccc cgagaaccag cgcaccgtcg ccgtatacag cttgaagatc 1020
gccgggtggc acgggcccac ggccccatac acgagcacc tgtgcccccc ggagctgtcc 1080
gagaccccc acgccacgca gccagaactc gccccggaag accccgagga ttcggccctc 1140
ttggaggacc ccgtggggac ggtggtgccc caaatccac caaactggca catacctcgt 1200
atccaggagc ccgcgagccc ttaccatccc ccggccaccc cgaacaacat gggcctgate 1260
gccggcgcgg tgggcggcag tctcctggta gccctggtca tttgcggaat tgtgtactgg 1320
atgcgccgcc gactcaaaa agccccaaa cgcatacgcc tccccacat ccgggaagac 1380
gaccagccgt cctcgcacca gcccttgttt tactag 1416

```

<210> SEQ ID NO 44

<211> LENGTH: 471

<212> TYPE: PRT

<213> ORGANISM: Herpes Simplex Virus-1

<400> SEQUENCE: 44

```

Met Ala Leu Leu Leu Thr Thr Val Ile Ala Leu Thr Cys Leu Gly Gly
1           5           10          15
Phe Ala Ser Pro Gly Pro Val Pro Pro Ser Thr Ala Leu Arg Glu Leu
20          25          30
Ile Glu Glu Leu Val Asn Ile Thr Gln Asn Gln Lys Ala Pro Leu Cys
35          40          45
Asn Gly Ser Met Val Trp Ser Ile Asn Leu Thr Ala Gly Met Tyr Cys
50          55          60
Ala Ala Leu Glu Ser Leu Ile Asn Val Ser Gly Cys Ser Ala Ile Glu
65          70          75          80
Lys Thr Gln Arg Met Leu Ser Gly Phe Cys Pro His Lys Val Ser Ala
85          90          95
Gly Gln Phe Ser Ser Leu His Val Arg Asp Thr Lys Ile Glu Val Ala
100         105         110
Gln Phe Val Lys Asp Leu Leu Leu His Leu Lys Lys Leu Phe Arg Glu
115         120         125

```

-continued

Gly Gln Phe Asn Gly Thr Gly Val Arg Arg Ser Tyr His Ile Gln Ala
 130 135 140

Gly Leu Pro Asp Pro Phe Gln Pro Pro Ser Leu Pro Ile Thr Val Tyr
 145 150 155 160

Tyr Ala Val Leu Glu Arg Ala Cys Arg Ser Val Leu Leu Asn Ala Pro
 165 170 175

Ser Glu Ala Pro Gln Ile Val Arg Gly Ala Ser Glu Asp Val Arg Lys
 180 185 190

Gln Pro Tyr Asn Leu Thr Ile Ala Trp Phe Arg Met Gly Gly Asn Cys
 195 200 205

Ala Ile Pro Ile Thr Val Met Glu Tyr Thr Glu Cys Ser Tyr Asn Lys
 210 215 220

Ser Leu Gly Ala Cys Pro Ile Arg Thr Gln Pro Arg Trp Asn Tyr Tyr
 225 230 235 240

Asp Ser Phe Ser Ala Val Ser Glu Asp Asn Leu Gly Phe Leu Met His
 245 250 255

Ala Pro Ala Phe Glu Thr Ala Gly Thr Tyr Leu Arg Leu Val Lys Ile
 260 265 270

Asn Asp Trp Thr Glu Ile Thr Gln Phe Ile Leu Glu His Arg Ala Lys
 275 280 285

Gly Ser Cys Lys Tyr Ala Leu Pro Leu Arg Ile Pro Pro Ser Ala Cys
 290 295 300

Leu Ser Pro Gln Ala Tyr Gln Gln Gly Val Thr Val Asp Ser Ile Gly
 305 310 315 320

Met Leu Pro Arg Phe Ile Pro Glu Asn Gln Arg Thr Val Ala Val Tyr
 325 330 335

Ser Leu Lys Ile Ala Gly Trp His Gly Pro Lys Ala Pro Tyr Thr Ser
 340 345 350

Thr Leu Leu Pro Pro Glu Leu Ser Glu Thr Pro Asn Ala Thr Gln Pro
 355 360 365

Glu Leu Ala Pro Glu Asp Pro Glu Asp Ser Ala Leu Leu Glu Asp Pro
 370 375 380

Val Gly Thr Val Val Pro Gln Ile Pro Pro Asn Trp His Ile Pro Ser
 385 390 395 400

Ile Gln Asp Ala Ala Thr Pro Tyr His Pro Pro Ala Thr Pro Asn Asn
 405 410 415

Met Gly Leu Ile Ala Gly Ala Val Gly Gly Ser Leu Leu Val Ala Leu
 420 425 430

Val Ile Cys Gly Ile Val Tyr Trp Met Arg Arg Arg Thr Gln Lys Ala
 435 440 445

Pro Lys Arg Ile Arg Leu Pro His Ile Arg Glu Asp Asp Gln Pro Ser
 450 455 460

Ser His Gln Pro Leu Phe Tyr
 465 470

1. A recombinant herpes simplex virus (HSV) particle having at least one protein on its surface, comprising:

- (a) an altered gD, wherein the alteration reduces binding of gD to one or more of its cellular receptors, said alteration comprising
 - (i) a heterologous peptide ligand on the surface of the recombinant HSV particle forming a fusion protein with the altered gD; and
 - (ii) an amino acid alteration;

wherein said recombinant HSV particle preferentially binds to cells expressing a binding partner to said heterologous peptide ligand.

2. The recombinant herpes simplex virus (HSV) particle of claim 1 further comprising an altered viral surface protein, wherein the alteration reduces binding of the viral surface protein to a sulfated proteoglycan.

3. The recombinant HSV particle of claim 1, wherein the alteration is a conservative amino acid substitution.

4. The recombinant HSV particle of claim 1, wherein the amino acid alteration is selected from the group consisting of an amino acid deletion, an amino acid substitution and an amino acid insertion.

5. The recombinant HSV particle of claim 4 wherein the amino acid alteration is an amino acid substitution.

6. The recombinant HSV particle of claim 5 wherein the amino acid substitution is at position 34.

7. The recombinant HSV particle of claim 6, wherein the amino acid substitution is V34S.

8. The recombinant HSV particle of claim 7, wherein the particle is selected from the group consisting of HSV R5141 and HSV R5161.

9. The recombinant HSV particle of claim 1, wherein a polynucleotide encoding the fusion protein is joined to a heterologous coding region for a leader sequence.

10. The recombinant HSV particle of claim 9, wherein the leader sequence is an HSV leader sequence.

11. The recombinant HSV particle of claim 10, wherein the leader sequence is an HSV gD leader sequence.

12. The recombinant HSV particle of claim 1, wherein a polynucleotide encoding the fusion protein is joined to a heterologous expression control element.

13. The recombinant HSV particle of claim 12, wherein the expression control element is a promoter.

14. The recombinant HSV particle of claim 1, wherein the viral surface protein is selected from the group consisting of gB and gC.

15. The recombinant HSV particle of claim 1, wherein the alteration of gD reduces binding to at least one protein selected from the group consisting of HveA and Nectin-1.

16. The recombinant HSV particle of claim 1, wherein the ligand forms a second fusion protein with a viral surface protein selected from the group consisting of gB and gC.

17. The recombinant HSV particle of claim 1, wherein the binding partner is a cell surface receptor for said ligand.

18. (canceled)

19. (canceled)

20. The recombinant HSV particle of claim 1, wherein the ligand is a cytokine.

21. The recombinant HSV particle of claim 20, wherein the cytokine is IL13.

22. The recombinant HSV particle of claim 1, wherein the ligand is a single-chain antibody.

23. A pharmaceutical composition comprising the recombinant HSV particle of claim 1 and a pharmaceutically acceptable carrier, diluent, or excipient.

24. A kit comprising the pharmaceutical composition according to claim 23 and a set of instructions for administering the composition to a subject in need.

25. A method of targeting a recombinant HSV particle to a cell comprising

(a) identifying a ligand for a ligand binding partner exhibited on the surface of a target cell; and

(b) creating an HSV particle according to claim 1, wherein the ligand binds to the binding partner exhibited on the surface of said target cell.

26. The method of claim 25, wherein said altered viral surface protein is selected from the group consisting of gB and gC.

27. The method of claim 25, wherein the alteration to gD reduces binding of gD to at least one cellular receptor for gD selected from the group consisting of HveA and Nectin-1.

28. The method of claim 25, wherein the ligand forms a second fusion protein with gC.

29. (canceled)

30. (canceled)

31. The method of claim 28, wherein the ligand is a cytokine.

32. The method of claim 31, wherein the cytokine is IL13.

33. The method of claim 25, wherein the ligand is a single-chain antibody.

34. The method of claim 25, wherein the amino acid substitution is V34S.

35.-39. (canceled)

40. A method of treating a cell-based disease comprising delivering a therapeutically effective amount of a recombinant HSV particle according to claim 1 to a subject in need.

41. (canceled)

42. The method according to claim 40 wherein the disease is cancer.

43. A method of ameliorating a symptom associated with a disease comprising administering a therapeutically effective amount of a recombinant HSV particle according to claim 1 to a subject in need.

44. (canceled)

45. The method according to claim 43 wherein the disease is characterized by hyperproliferative cells.

46. A method of delivering a therapeutically useful peptide to a cell comprising:

(a) inserting a coding region for a therapeutically useful peptide into the DNA of a recombinant HSV particle according to claim 1, thereby producing a recombinant HSV clone; and

(b) delivering a therapeutically effective amount of said recombinant HSV clone to said cell.

47. (canceled)

48. The method according to claim 46 wherein the recombinant HSV clone is delivered in vivo.

49. (canceled)

50. (canceled)

51. The use according to claim 41 wherein the disease is cancer.

52. The use according to claim 44 wherein the disease is characterized by hyperproliferative cells.

53. The use according to claim 47 wherein the recombinant HSV clone is delivered in vivo.