



US008821894B2

(12) **United States Patent**  
**Schneewind et al.**

(10) **Patent No.:** **US 8,821,894 B2**  
(45) **Date of Patent:** **\*Sep. 2, 2014**

(54) **COMPOSITIONS AND METHODS RELATED TO PROTEIN A (SPA) VARIANTS**

(75) Inventors: **Olaf Schneewind**, Chicago, IL (US);  
**Alice G. Cheng**, Chicago, IL (US);  
**Dominique M. Missiakas**, Chicago, IL (US); **Hwan Keun Kim**, Chicago, IL (US)

(73) Assignee: **The University of Chicago**, Chicago, IL (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.  
  
This patent is subject to a terminal disclaimer.

2004/0043037	A1	3/2004	Kunsch et al.	424/190.1
2004/0265962	A1	12/2004	Bailey et al.	435/69.1
2005/0106597	A1	5/2005	Choi	435/6.15
2005/0255478	A1	11/2005	Kimmerly	435/6.15
2006/0205016	A1	9/2006	Silverman	435/6
2007/0020746	A1	1/2007	Kunsch et al.	435/252.3
2008/0311146	A1	12/2008	Castado	424/243.1
2009/0317421	A1	12/2009	Missiakas	424/184.1
2010/0272743	A1	10/2010	Kimmerly	424/190.1
2011/0027265	A1	2/2011	Bubeck-Wardenburg	424/184.1
2011/0206676	A1	8/2011	Missiakas	424/184.1
2011/0262477	A1	10/2011	Cheng	424/190.1
2012/0114686	A1	5/2012	Schneewind	424/190.1
2012/0282247	A1	11/2012	Schneewind	424/150.1
2013/0136746	A1	5/2013	Schneewind	424/150.1
2013/0189249	A1	7/2013	Bubeck-Wardenburg	424/184.1
2013/0230550	A1	9/2013	Schneewind	424/190.1
2013/0236419	A1	9/2013	Schneewind	424/165.15

(21) Appl. No.: **13/807,598**

(22) PCT Filed: **Jul. 1, 2011**

(86) PCT No.: **PCT/US2011/042845**

§ 371 (c)(1),  
(2), (4) Date: **Mar. 19, 2013**

(87) PCT Pub. No.: **WO2012/003474**

PCT Pub. Date: **Jan. 5, 2012**

(65) **Prior Publication Data**

US 2013/0171183 A1 Jul. 4, 2013

**Related U.S. Application Data**

(60) Provisional application No. 61/361,218, filed on Jul. 2, 2010, provisional application No. 61/370,725, filed on Aug. 4, 2010.

(51) **Int. Cl.**  
**A61K 39/085** (2006.01)  
**C07K 14/31** (2006.01)

(52) **U.S. Cl.**  
USPC ..... **424/243.1**; 424/190.1; 424/235.1;  
530/324

(58) **Field of Classification Search**  
None  
See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

5,294,177	A	3/1994	Rasnick et al.	297/270.1
5,648,240	A	7/1997	Hook	435/69.3
5,801,234	A	9/1998	Hodgson	536/23.7
5,840,846	A	11/1998	Hook	530/350
6,008,341	A	12/1999	Foster	536/23.7
6,288,214	B1	9/2001	Hook	530/387.1
6,403,337	B1	6/2002	Bailey et al.	435/69.7
6,593,114	B1	7/2003	Kunsch et al.	435/91.41
6,703,492	B1	3/2004	Kimmerly	536/23.1
6,737,248	B2	5/2004	Kunsch et al.	435/69.1
6,753,149	B2	6/2004	Bailey et al.	435/6.15
6,833,253	B2	12/2004	Choi	435/69.1
2002/0169288	A1	11/2002	Hook	424/190.1

**FOREIGN PATENT DOCUMENTS**

EP	0786519	7/1997
EP	0594610	9/1998
WO	WO 98/57994	12/1998
WO	WO 00/02523	1/2000
WO	WO 00/12132	3/2000
WO	WO 00/12689	3/2000
WO	WO 00/15238	3/2000
WO	WO 01/34809	5/2001
WO	WO 01/60852	8/2001
WO	WO 01/98499	12/2001
WO	WO 02/059148	8/2002
WO	WO 02/094868	11/2002
WO	WO 2006/032472	3/2006
WO	WO 2006/032475	3/2006
WO	WO 2006/032500	3/2006
WO	WO 2007/113222	10/2007
WO	WO 2007/113223	10/2007
WO	WO 2012/122533	9/2012

**OTHER PUBLICATIONS**

Abdallah et al., "A specific secretion system mediates PPE41 transport in pathogenic mycobacteria", *Mol. Microbiol.*, 62, 667-679, 2006.

Abdallah et al., "Type VII secretion—mycobacteria show the way", *Nat. Rev. Microbiol.*, 5:883-891, 2007.

(Continued)

*Primary Examiner* — Padma V Baskar

(74) *Attorney, Agent, or Firm* — Fulbright & Jaworski LLP

(57) **ABSTRACT**

The present invention concerns methods and compositions for treating or preventing a bacterial infection, particularly infection by a *Staphylococcus* bacterium. The invention provides methods and compositions for stimulating an immune response against the bacteria. In certain embodiments, the methods and compositions involve a non-toxicogenic Protein A (SpA) variant.

**11 Claims, 12 Drawing Sheets**

(56)

## References Cited

## OTHER PUBLICATIONS



- Andersen et al., "Recall of Long-Lived Immunity to *Mycobacterium tuberculosis* Infection in Mice", *J. Immunol.*, 154:3359-3372, 1995.
- Archer, "Staphylococcus aureus: A Well-Armed Pathogen", *Clin. Infect. Dis.*, 26:1179-1181, 1998.
- Baba et al., "Genome Sequence of *Staphylococcus aureus* Strain Newman and comparative analysis of Staphylococcal Genomes: Polymorphism and Evolution of Two Major Pathogenicity Islands", *J. Bacteriol.* 190(1):300-310, 2007.
- Bae et al., "Staphylococcus aureus virulence genes identified by bursa aurealis mutagenesis and nematode killing", *Proc. Natl. Acad. Sci. USA*, 101(33):12312-12317, 2004.
- Bjerketorp et al., "The von Willebrand factor-binding protein (vWbp) of *Staphylococcus aureus* is a coagulase", *FEMS Microbiol. Lett.*, 234:309-314, 2004.
- Brown et al., "Determining Protein-Protein Interactions by Oxidative Cross-Linking of a Glycine-Glycine-Histidine Fusion Protein", *Biochemistry*, 37:4397-4406, 1998.
- Burts et al., "EssA and EssB are secreted by an ESAT-6-like system that is required for the pathogenesis of *Staphylococcus aureus* infections", *Proc. Natl. Acad. Sci. USA*, 102(4):1169-1174, 2005.
- Burts et al., "EsaC substrate for the ESAT-6 Secretion Pathway and its role in persistent infections of *S. aureus*", *Mol. Microbiol.*, 69(3):736-746, 2008.
- Cedergren et al., "Mutational analysis of the interaction between staphylococcal protein A and human IgG1", *Protein Eng.*, 6(4):441-448, 1993.
- Cespedes et al., "The Clonality of *Staphylococcus aureus* Nasal Carriage", *J. Infect. Dis.*, 191(3):444-452, 2005.
- Cheng et al., "Genetic requirements for *Staphylococcus aureus* abscess formation and persistence in host tissues", *FASEB J.*, 23:3393-3404, 2009.
- Dalbey and Wickner, "Leader Peptidase Catalyzes the Release of Exported Proteins from the Outer Surface of the *Escherichia coli* Plasma Membrane", *J. Biol. Chem.*, 260(29):15925-15931, 1985.
- DeDent et al., "Distribution of Protein A on the Surface of *Staphylococcus aureus*", *J. Bacteriol.* 189:4473-4484, 2007.
- DeDent et al., "Signal peptides direct surface proteins to two distinct envelope locations of *Staphylococcus aureus*", *EMBO J.* 27:2656-2668, 2008.
- Deisenhofer et al., "Crystallization, Crystal Structure Analysis and Atomic Model of the Complex Formed by a Human Fc Fragment and Fragment B of Protein A from *Staphylococcus aureus*", *Hoppe-Seihl Zeitsch. Physiol. Chem.* 359:975-985, 1978.
- Deisenhofer, "Crystallographic Refinement and Atomic Models of a Human Fc Fragment and Its Complex with Fragment B of Protein A from *Staphylococcus aureus* at 2.9- and 2.8-Å Resolution", *Biochemistry* 20(9):2361-2370, 1981.
- Dinges et al., "Exotoxins of *Staphylococcus aureus*", *Clin. Microbiol. Rev.*, 13(1):16-34, 2000.
- Duthie and Lorenz, "Staphylococcal coagulase: Mode of action and Antigenicity", *J. Gen. Microbiol.*, 6:95-107, 1952.
- Ekstedt and Yotis, "Effect of Coagulase on the Virulence of Coagulase Negative Strains", *Ann. N.Y. Acad. Sci.*, 80:496-500, 1960.
- Field and Smith, "The Coagulase Test for *Staphylococci*", *J. Comp. Pathol.*, 55:63-69, 1945.
- Fortune et al., "Mutually dependent secretion of proteins required for mycobacterial virulence", *Proc Natl. Acad. Sci. USA*, 102(30):10676-10681, 2005.
- Foster, "Immune Evasion by *Staphylococci*", *Nat. Rev. Microbiol.*, 3:948-958, 2005.
- Fournier et al., "Purification and Characterization of *Staphylococcus aureus* Type 8 Capsular Polysaccharide", *Infect. Immun.*, 45(1):87-93, 1984.
- Galan and Collmer, "Type III Secretion Machines: Bacterial Devices for Protein Delivery into Host Cells", *Science*, 284:1322-1328, 1999.
- Gomez et al., "Staphylococcus aureus protein A induces airway epithelial inflammatory responses by activating TNFR1", *Nature Med.* 10(8):842-848, 2004.
- Gomez et al., "Mechanisms of Signal Transduction: *Staphylococcus aureus* Protein A Activates TNFR1 Signaling through Conserved IgG Binding Domains", *J. Biol. Chem.* 281(29):20190-20196, 2006.
- Gomez et al., "Staphylococcus aureus protein A activates TACE through EGFR-dependent signaling", *EMBO J.* 26:701-709, 2007.
- Goodyear and Silverman, "Staphylococcal toxin induced preferential and prolonged in vivo deletion of innate-like B lymphocytes", *Proc. Nat. Acad. Sci. USA*, 101(31):11392-11397, 2004.
- Goodyear and Silverman, "Death by a B Cell Superantigen: In Vivo VH-targeted Apoptotic Supraclonal B Cell Deletion by a Staphylococcal Toxin", *J. Exp. Med.*, 197(9):1125-1139, 2003.
- Gouda et al., "Three-Dimensional Solution Structure of the B Domain of Staphylococcal Protein A: Comparisons of the Solution and Crystal Structures", *Biochemistry*, 31(40):9665-72, 1992.
- Graille et al., "Crystal structure of a *Staphylococcus aureus* protein A domain complexed with the Fab fragment of a human IgM antibody: Structural basis for recognition of B-cell receptors and superantigen activity", *Proc. Nat. Acad. Sci. USA* 97(10):5399-5404, 2000.
- Guss et al., "Region X, the cell-wall-attachment part of staphylococcal protein A", *Eur. J. Biochem.* 138:413-420, 1984.
- Hartleib et al., "Protein A is the von Willebrand factor binding protein on *Staphylococcus aureus*", *Blood* 96:2149-2156, 2000.
- Hsu et al., "The primary mechanism of attenuation of bacillus Calmette-Guerin is a loss of secreted lytic function required for invasion of lung interstitial tissue", *Proc. Natl. Acad. Sci. USA*, 100(21):12420-12425, 2003.
- Jansson et al., "All individual domains of staphylococcal protein A show Fab binding", *FEMS Immunol. Med. Microbiol.* 20:69-78, 1998.
- Jensen, "A Normally Occurring *Staphylococcus* Antibody in Human Serum", *Acta Path. Microbiol. Scand.* 44:421-428, 1958.
- Jonsson et al., "Virulence of *Staphylococcus aureus* in a Mouse Mastitis Model: Studies of Alpha Hemolysin, Coagulase, and Protein A as Possible Virulence Determinants with Protoplast Fusion and Gene Cloning", *Infection and Immunity*, 49(3):765-769, 1985.
- Kennedy et al., "Epidemic community-associated methicillin-resistant *Staphylococcus aureus*: Recent clonal expansion and diversification", *Proc. Natl. Acad. Sci. USA* 105(4):1327-1332, 2008.
- Kuklin et al., "A Novel *Staphylococcus aureus* Vaccine: Iron Surface Determinant B Induces Rapid Antibody Responses in Rhesus Macaques and Specific Increased Survival in a Murine *S. aureus* Sepsis Model", *Infect. Immun.*, 74(4):2215-23, 2006.
- Lagergard et al., "Determination of Neutralizing Antibodies and Specific Immunoglobulin Isotype Levels in Infants after Vaccination against Diphtheria", *Eur. J. Clin. Microbiol. Infect. Dis.*, 11(4):341-345, 1992.
- Lam et al., "Abscess-Forming Factor(s) Produced by *Staphylococcus aureus*", *J. Bacteriol.*, 86:87-91, 1963.
- Lee, "The prospects for developing a vaccine against *Staphylococcus aureus*", *Trends Microbiol.* 4(4):162-166, 1996.
- Lowy, "Staphylococcus aureus Infections", *New Engl. J. Med.*, 339(8):520-532, 1998.
- MacGurn et al., "A non-RD1 gene cluster is required for Snm secretion in *Mycobacterium tuberculosis*", *Mol. Microbiol.*, 57(6):1653-1663, 2005.
- Mazmanian et al., "Staphylococcus aureus Sortase, an Enzyme that Anchors Surface Proteins to the Cell Wall", *Science*, 285(5428):760-3, 1999.
- Mazmanian et al., "Staphylococcus aureus sortase mutants defective in the display of surface proteins and in the pathogenesis of animal infections", *Proc. Natl. Acad. Sci. USA*, 97(10):5510-5515, 2000.
- Mazmanian et al., "Sortase-catalysed anchoring of surface proteins to the cell wall of *Staphylococcus aureus*", *Mol. Microbiol.*, 40(5):1049-1057, 2001.
- McLaughlin et al., "A *Mycobacterium* ESX-1-Secreted Virulence Factor with Unique Requirements for Export", *PLoS Pathog.*, 3(8):1051-1061, 2007.
- Moreau et al., "Structure of the type 5 capsular polysaccharide of *Staphylococcus aureus*", *Carbohydrate Res.*, 201(2):285-297, 1990.
- Moreillon et al., "Role of *Staphylococcus aureus* Coagulase and Clumping Factor in Pathogenesis of Experimental Endocarditis", *Infect. Immun.*, 63(12):4738-4743, 1995.

(56)

## References Cited

## OTHER PUBLICATIONS

- Novick, "Autoinduction and signal transduction in the regulation of staphylococcal virulence", *Mol. Microbiol.*, 48(6):1429-1449, 2003.
- O'Brien et al., "Multiple mechanisms for the activation of human platelet aggregation by *Staphylococcus aureus*: roles for the clumping factors ClfA and ClfB, the serine-aspartate repeat protein SdrE and protein A", *Mol. Microbiol.* 44(4):1033-1044, 2002.
- O'Seaghda et al., "Staphylococcus aureus protein A binding to von Willebrand factor A1 domain is mediated by conserved IgG binding regions", *FEBS J.* 273:4831-4841, 2006.
- Pallen, "The ESAT-6/WXG100 superfamily—and a new Gram-positive secretion system?", *Trends Microbiol.*, 10(5):209-212, 2002.
- Palmqvist et al., "Bacterial cell wall-expressed protein A triggers supraclonal B-cell responses upon in vivo infection with *Staphylococcus aureus*", *Microbes. Infect.*, 7:1501-1511, 2005.
- Panizzi et al., "Fibrinogen Substrate Recognition by Staphylocoagulase-(Pro) thrombin Complexes", *J. Biol. Chem.*, 281(2):1179-1187, 2006.
- Phonimdaeng et al., "The coagulase of *Staphylococcus aureus* 8325-4. Sequence analysis and virulence of site-specific coagulase-deficient mutants", *Mol. Microbiol.*, 4(3):393-404, 1990.
- Pym et al., "Loss of RD1 contributed to the attenuation of the live tuberculosis vaccines *Mycobacterium bovis* BCG and *Mycobacterium microti*", *Molecular Microbiology*, 46(3):709-717, 2002.
- Roben et al., "VH3 Family Antibodies Bind Domain D of Staphylococcal Protein A1", *J. Immunol.* 154:6437-6445, 1995.
- Said-Salim et al., "Community-Acquired Methicillin-Resistant *Staphylococcus aureus*: An Emerging Pathogen", *Infect. Control Hosp. Epidemiol.* 24(6):451-455, 2003.
- Schneewind et al., "Sorting of Protein A to the Staphylococcal Cell Wall", *Cell* 70:267-281, 1992.
- Search Report and Written Opinion in PCT/US11/42845 mailed Feb. 10, 2012.
- Shaw et al., "The role and regulation of the extracellular proteases of *Staphylococcus aureus*", *Microbiology*, 150:217-228, 2004.
- Sheagren, "Staphylococcus aureus: The Persistent Pathogen", *N. Engl. J. Med.* 310(21):1368-1373, 1984.
- Sibbald et al., "Mapping the Pathways to Staphylococcal Pathogenesis by Comparative Scretomics", *Microbiol. Mol Biol. Rev.*, 70(3):755-788, 2006.
- Sjodahl, "Repetitive Sequences in Protein A from *Staphylococcus aureus*", *Eur. J. Biochem.* 73:343-351, 1977.
- Sjoquist et al., "Protein A Isolated from *Staphylococcus aureus* after Digestion with Lysostaphin", *Eur. J. Biochem.* 29 :572-578, 1972.
- Smith et al., "The Role of Coagulase in Staphylococcal Infections", *Brit. J. Exp. Pathol.*, 28:57-67, 1947.
- Stanley et al., "Acute infection and macrophage subversion by *Mycobacterium tuberculosis* require a specialized secretion system", *Proc. Natl. Acad. Sci. USA*, 100(22):13001-13006, 2003.
- Stranger-Jones et al., "Vaccine assembly from surface proteins of *Staphylococcus aureus*", *Proc. Nat. Acad. Sci. USA*, 103(45):16942-16947, 2006.
- Ton-That et al., "Purification and characterization of sortase, the transpeptidase that cleaves surface proteins of *Staphylococcus aureus* at the LPXTG motif", *Proc. Natl. Acad. Sci. USA*, 96(22):12424-12429, 1999.
- Uhlen et al., "Complete Sequence of the Staphylococcal Gene Encoding Protein A: A gene evolved through multiple duplications", *J. Biol. Chem.* 259(3):1695-1702, 1984.
- van Wely et al., "Translocation of proteins across the cell envelope of Gram-positive bacteria", *FEMS Microbiol. Rev.*, 25:437-454, 2001.
- Weiss et al., "Effect of srtA and srtB gene expression on the virulence of *Staphylococcus aureus* in animal models of infection", *J. Antimicrob. Chemother.*, 53:480-486, 2004.
- Xu et al., "A unique *Mycobacterium* ESX-1 protein co-secreted with CFP-10/ESAT-6 and is necessary for inhibiting phagosome maturation", *Mol. Microbiol.*, 66(3):787-800, 2007.



FIGs. 1A-1B




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7






FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12A-12C

## COMPOSITIONS AND METHODS RELATED TO PROTEIN A (SPA) VARIANTS

The present application is a national phase application under 35 U.S.C. §371 of International Patent Application No. PCT/US2011/042845 filed Jul. 1, 2011, which claims the benefit of priority to U.S. Provisional Patent Application Ser. No. 61/361,218 filed Jul. 2, 2010, and 61/370,725 filed Aug. 4, 2010, hereby incorporated by reference in their entirety.

This invention was made with government support under AI057153, AI052474, and GM007281 awarded by the National Institutes of Health. The government has certain rights in the invention.

### BACKGROUND OF THE INVENTION

#### I. Field of the Invention

The present invention relates generally to the fields of immunology, microbiology, and pathology. More particularly, it concerns methods and compositions involving bacterial Protein A variants, which can be used to invoke an immune response against the bacteria.

#### II. Background

The number of both community acquired and hospital acquired infections have increased over recent years with the increased use of intravascular devices. Hospital acquired (nosocomial) infections are a major cause of morbidity and mortality, more particularly in the United States, where it affects more than 2 million patients annually. The most frequent infections are urinary tract infections (33% of the infections), followed by pneumonia (15.5%), surgical site infections (14.8%) and primary bloodstream infections (13%) (Emorl and Gaynes, 1993).

The major nosocomial pathogens include *Staphylococcus aureus*, coagulase-negative Staphylococci (mostly *Staphylococcus epidermidis*), *enterococcus* spp., *Escherichia coli* and *Pseudomonas aeruginosa*. Although these pathogens cause approximately the same number of infections, the severity of the disorders they can produce combined with the frequency of antibiotic resistant isolates balance this ranking towards *S. aureus* and *S. epidermidis* as being the most significant nosocomial pathogens.

Staphylococci can cause a wide variety of diseases in humans and other animals through either toxin production or invasion. Staphylococcal toxins are also a common cause of food poisoning, as the bacteria can grow in improperly-stored food.

*Staphylococcus epidermidis* is a normal skin commensal which is also an important opportunistic pathogen responsible for infections of impaired medical devices and infections at sites of surgery. Medical devices infected by *S. epidermidis* include cardiac pacemakers, cerebrospinal fluid shunts, continuous ambulatory peritoneal dialysis catheters, orthopedic devices and prosthetic heart valves.

*Staphylococcus aureus* is the most common cause of nosocomial infections with a significant morbidity and mortality. It is the cause of some cases of osteomyelitis, endocarditis, septic arthritis, pneumonia, abscesses, and toxic shock syndrome. *S. aureus* can survive on dry surfaces, increasing the chance of transmission. Any *S. aureus* infection can cause the staphylococcal scalded skin syndrome, a cutaneous reaction to exotoxin absorbed into the bloodstream. It can also cause a type of septicemia called pyaemia that can be life-threatening. Problematically, Methicillin-resistant *Staphylococcus aureus* (MRSA) has become a major cause of hospital-acquired infections.

*S. aureus* and *S. epidermidis* infections are typically treated with antibiotics, with penicillin being the drug of choice, whereas vancomycin is used for methicillin resistant isolates. The percentage of staphylococcal strains exhibiting wide-spectrum resistance to antibiotics has become increasingly prevalent, posing a threat for effective antimicrobial therapy. In addition, the recent emergence of vancomycin resistant *S. aureus* strain has aroused fear that MRSA strains are emerging and spreading for which no effective therapy is available.

An alternative to antibiotic treatment for staphylococcal infections is under investigation that uses antibodies directed against staphylococcal antigens. This therapy involves administration of polyclonal antisera (WO00/15238, WO00/12132) or treatment with monoclonal antibodies against lipoteichoic acid (WO98/57994).

An alternative approach would be the use of active vaccination to generate an immune response against staphylococci. The *S. aureus* genome has been sequenced and many of the coding sequences have been identified (WO02/094868, EP0786519), which can lead to the identification of potential antigens. The same is true for *S. epidermidis* (WO01/34809). As a refinement of this approach, others have identified proteins that are recognized by hyperimmune sera from patients who have suffered staphylococcal infection (WO01/98499, WO02/059148).

*S. aureus* secretes a plethora of virulence factors into the extracellular milieu (Archer, 1998; Dinges et al., 2000; Foster, 2005; Shaw et al., 2004; Sibbald et al., 2006). Like most secreted proteins, these virulence factors are translocated by the Sec machinery across the plasma membrane. Proteins secreted by the Sec machinery bear an N-terminal leader peptide that is removed by leader peptidase once the pre-protein is engaged in the Sec translocon (Dalbey and Wickner, 1985; van Wely et al., 2001). Recent genome analysis suggests that Actinobacteria and members of the Firmicutes encode an additional secretion system that recognizes a subset of proteins in a Sec-independent manner (Pallen, 2002). ESAT-6 (early secreted antigen target 6 kDa) and CFP-10 (culture filtrate antigen 10 kDa) of *Mycobacterium tuberculosis* represent the first substrates of this novel secretion system termed ESX-1 or 5 nm in *M. tuberculosis* (Andersen et al., 1995; Hsu et al., 2003; Pym et al., 2003; Stanley et al., 2003). In *S. aureus*, two ESAT-6 like factors designated ExsA and ExsB are secreted by the Ess pathway (ESAT-6 secretion system) (Burts et al., 2005).

The first generation of vaccines targeted against *S. aureus* or against the exoproteins it produces have met with limited success (Lee, 1996). There remains a need to develop effective vaccines against staphylococcal infections. Additional compositions for treating staphylococcal infections are also needed.

### SUMMARY OF THE INVENTION

Protein A (SpA) (SEQ ID NO:33), a cell wall anchored surface protein of *Staphylococcus aureus*, provides for bacterial evasion from innate and adaptive immune responses. Protein A binds immunoglobulins at their Fc portion, interacts with the VH3 domain of B cell receptors inappropriately stimulating B cell proliferation and apoptosis, binds to von Willebrand factor A1 domains to activate intracellular clotting, and also binds to the TNF Receptor-1 to contribute to the pathogenesis of staphylococcal pneumonia. Due to the fact that Protein A captures immunoglobulin and displays toxic attributes, the possibility that this surface molecule may function as a vaccine in humans has not been rigorously pursued. Here the inventors demonstrate that Protein A variants no

longer able to bind to immunoglobulins, which are thereby removed of their toxigenic potential, i.e., are non-toxicogenic, stimulate humoral immune responses that protect against staphylococcal disease.

In certain embodiments the SpA variant is a full length SpA variant comprising a variant A, B, C, D, and/or E domain. In certain aspects, the SpA variant comprises or consists of the amino acid sequence that is 80, 90, 95, 98, 99, or 100% identical to the amino acid sequence of SEQ ID NO:34. In other embodiments the SpA variant comprises a segment of SpA. The SpA segment can comprise at least or at most 1, 2, 3, 4, 5 or more IgG binding domains. The IgG domains can be at least or at most 1, 2, 3, 4, 5 or more variant A, B, C, D, or E domains. In certain aspects the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant A domains. In a further aspect the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant B domains. In still a further aspect the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant C domains. In yet a further aspect the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant D domains. In certain aspects the SpA variant comprises at least or at most 1, 2, 3, 4, 5, or more variant E domains. In a further aspect the SpA variant comprises a combination of A, B, C, D, and E domains in various combinations and permutations. The combinations can include all or part of a SpA signal peptide segment, a SpA region X segment, and/or a SpA sorting signal segment. In other aspects the SpA variant does not include a SpA signal peptide segment, a SpA region X segment, and/or a SpA sorting signal segment. In certain aspects a variant A domain comprises a substitution at position(s) 7, 8, 34, and/or 35 of SEQ ID NO:4. In another aspect a variant B domain comprises a substitution at position(s) 7, 8, 34, and/or 35 of SEQ ID NO:6. In still another aspect a variant C domain comprises a substitution at position(s) 7, 8, 34, and/or 35 of SEQ ID NO:5. In certain aspects a variant D domain comprises a substitution at position(s) 9, 10, 36, and/or 37 of SEQ ID NO:2. In a further aspect a variant E domain comprises a substitution at position(s) 6, 7, 33, and/or 34 of SEQ ID NO:3.

In certain aspects, an SpA domain D variant or its equivalent can comprise a mutation at position 9 and 36; 9 and 37; 9 and 10; 36 and 37; 10 and 36; 10 and 37; 9, 36, and 37; 10, 36, and 37, 9, 10 and 36; or 9,10 and 37 of SEQ ID NO:2. In a further aspect, analogous mutations can be included in one or more of domains A, B, C, or E.

In further aspects, the amino acid glutamine (Q) at position 9 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an asparagine (N), an aspartic acid (D), a cysteine (C), a glutamic acid (E), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the glutamine at position 9 can be substituted with an arginine (R). In a further aspect, the glutamine at position 9 of SEQ ID NO:2, or its equivalent, can be substituted with a lysine or a glycine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

In another aspect, the amino acid glutamine (Q) at position 10 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an asparagine (N), an aspartic acid (D), a cysteine (C), a glutamic acid (E), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the glutamine at position 10 can be substituted with an arginine

(R). In a further aspect, the glutamine at position 10 of SEQ ID NO:2, or its equivalent, can be substituted with a lysine or a glycine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

In certain aspects, the aspartic acid (D) at position 36 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an asparagine (N), an arginine (R), a cysteine (C), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a glutamine (Q), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the aspartic acid at position 36 can be substituted with a glutamic acid (E). In certain aspects, an aspartic acid at position 36 of SEQ ID NO:2, or its equivalent, can be substituted with an alanine or a serine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

In another aspect, the aspartic acid (D) at position 37 of SEQ ID NO:2 (or its analogous amino acid in other SpA domains) can be replaced with an alanine (A), an asparagine (N), an arginine (R), a cysteine (C), a phenylalanine (F), a glycine (G), a histidine (H), an isoleucine (I), a lysine (K), a leucine (L), a methionine (M), a proline (P), a glutamine (Q), a serine (S), a threonine (T), a valine (V), a tryptophane (W), or a tyrosine (Y). In some aspects the aspartic acid at position 37 can be substituted with a glutamic acid (E). In certain aspects, an aspartic acid at position 37 of SEQ ID NO:2, or its equivalent, can be substituted with an alanine or a serine. Any 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more of the substitutions can be explicitly excluded.

In a particular embodiment the amino at position 9 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a valine (V). In certain aspects the amino acid at position 9 of SEQ ID NO:2 is replaced by a glycine. In a further aspect the amino acid at position 9 of SEQ ID NO:2 is replaced by a lysine.

In a particular embodiment the amino at position 10 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a valine (V). In certain aspects the amino acid at position 10 of SEQ ID NO:2 is replaced by a glycine. In a further aspect the amino acid at position 10 of SEQ ID NO:2 is replaced by a lysine.

In a particular embodiment the amino at position 36 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a valine (V). In certain aspects the amino acid at position 36 of SEQ ID NO:2 is replaced by a serine. In a further aspect the amino acid at position 36 of SEQ ID NO:2 is replaced by an alanine.

In a particular embodiment the amino at position 37 of SEQ ID NO:2 (or an analogous amino acid in another SpA domain) is replaced by an alanine (A), a glycine (G), an isoleucine (I), a leucine (L), a proline (P), a serine (S), or a valine (V). In certain aspects the amino acid at position 37 of SEQ ID NO:2 is replaced by a serine. In a further aspect the amino acid at position 37 of SEQ ID NO:2 is replaced by an alanine.

In certain aspects the SpA variant includes a substitution of (a) one or more amino acid substitution in an IgG Fc binding sub-domain of SpA domain A, B, C, D, and/or E that disrupts or decreases binding to IgG Fc, and (b) one or more amino acid substitution in a  $V_H3$  binding sub-domain of SpA domain A, B, C, D, and/or E that disrupts or decreases binding to  $V_H3$ . In still further aspects the amino acid sequence of a SpA variant comprises an amino acid sequence that is at least 50%, 60%, 70%, 80%, 90%, 95%, or 100% identical, includ-

ing all values and ranges there between, to the amino acid sequence of SEQ ID NOs:2-6.

In a further aspect the SpA variant includes (a) one or more amino acid substitution in an IgG Fc binding sub-domain of SpA domain D, or at a corresponding amino acid position in other IgG domains, that disrupts or decreases binding to IgG Fc, and (b) one or more amino acid substitution in a  $V_H3$  binding sub-domain of SpA domain D, or at a corresponding amino acid position in other IgG domains, that disrupts or decreases binding to  $V_H3$ . In certain aspects amino acid residue F5, Q9, Q10, S11, F13, Y14, L17, N28, I31, and/or K35 (SEQ ID NO:2, QQNNFNKDQQSAFYEILNMPNLNEAQRNGFIQSLKDDPSQSTNVLGEAKKLNES) of the IgG Fc binding sub-domain of domain D are modified or substituted. In certain aspects amino acid residue Q26, G29, F30, S33, D36, D37, Q40, N43, and/or E47 (SEQ ID NO:2) of the  $V_H3$  binding sub-domain of domain D are modified or substituted such that binding to Fc or  $V_H3$  is attenuated. In further aspects corresponding modifications or substitutions can be engineered in corresponding positions of the domain A, B, C, and/or E. Corresponding positions are defined by alignment of the domain D amino acid sequence with one or more of the amino acid sequences from other IgG binding domains of SpA, for example see FIG. 2A. In certain aspects the amino acid substitution can be any of the other 20 amino acids. In a further aspect conservative amino acid substitutions can be specifically excluded from possible amino acid substitutions. In other aspects only non-conservative substitutions are included. In any event, any substitution or combination of substitutions that reduces the binding of the domain such that SpA toxicity is significantly reduced is contemplated. The significance of the reduction in binding refers to a variant that produces minimal to no toxicity when introduced into a subject and can be assessed using in vitro methods described herein.

In certain embodiments, a variant SpA comprises at least or at most 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more variant SpA domain D peptides. In certain aspects 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19 or more amino acid residues of the variant SpA are substituted or modified—including but not limited to amino acids F5, Q9, Q10, S11, F13, Y14, L17, N28, I31, and/or K35 (SEQ ID NO:2) of the IgG Fc binding sub-domain of domain D and amino acid residue Q26, G29, F30, S33, D36, D37, Q40, N43, and/or E47 (SEQ ID NO:2) of the  $V_H3$  binding sub-domain of domain D. In one aspect of the invention glutamine residues at position 9 and/or 10 of SEQ ID NO:2 (or corresponding positions in other domains) are mutated. In another aspect, aspartic acid residues 36 and/or 37 of SEQ ID NO:2 (or corresponding positions in other domains) are mutated. In a further aspect, glutamine 9 and 10, and aspartic acid residues 36 and 37 are mutated. Purified non-toxicogenic SpA or SpA-D mutants/variants described herein are no longer able to significantly bind (i.e., demonstrate attenuated or disrupted binding affinity) Fc $\gamma$  or F(ab) $_2$   $V_H3$  and also do not stimulate B cell apoptosis. These non-toxicogenic Protein A variants can be used as subunit vaccines and raise humoral immune responses and confer protective immunity against *S. aureus* challenge. Compared to wild-type full-length Protein A or the wild-type SpA-domain D, immunization with SpA-D variants resulted in an increase in Protein A specific antibody. Using a mouse model of staphylococcal challenge and abscess formation, it was observed that immunization with the non-toxicogenic Protein A variants generated significant protection from staphylococcal infection and abscess formation. As virtually all *S. aureus* strains express Protein A, immunization of humans with the non-toxicogenic Protein A variants can neutralize this virulence

factor and thereby establish protective immunity. In certain aspects the protective immunity protects or ameliorates infection by drug resistant strains of *Staphylococcus*, such as USA300 and other MRSA strains.

Embodiments include the use of Protein A variants in methods and compositions for the treatment of bacterial and/or staphylococcal infection. This application also provides an immunogenic composition comprising a Protein A variant or immunogenic fragment thereof. In certain aspects, the immunogenic fragment is a Protein A domain D segment. Furthermore, the present invention provides methods and compositions that can be used to treat (e.g., limiting staphylococcal abscess formation and/or persistence in a subject) or prevent bacterial infection. In some cases, methods for stimulating an immune response involve administering to the subject an effective amount of a composition including or encoding all or part of a Protein A variant polypeptide or antigen, and in certain aspects other bacterial proteins. Other bacterial proteins include, but are not limited to (i) a secreted virulence factor, and/or a cell surface protein or peptide, or (ii) a recombinant nucleic acid molecule encoding a secreted virulence factor, and/or a cell surface protein or peptide.

In other aspects, the subject can be administered all or part of a Protein A variant, such as a variant Protein A domain D segment. The polypeptide of the invention can be formulated in a pharmaceutically acceptable composition. The composition can further comprise one or more of at least or at most 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19 additional staphylococcal antigen or immunogenic fragment thereof (e.g., Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla (e.g., H35 mutants), IsdC, SasF, vWbp, or vWh). Additional staphylococcal antigens that can be used in combination with a Protein A variant include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa (GenBank CAC80837), Aap (GenBank accession AJ249487), Ant (GenBank accession NP\_372518), autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg $^{2+}$  transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein (see PCT publications WO2007/113222, WO2007/113223, WO2006/032472, WO2006/032475, WO2006/032500, each of which is incorporated herein by reference in their entirety). The staphylococcal antigen or immunogenic fragment can be administered concurrently with the Protein A variant. The staphylococcal antigen or immunogenic fragment and the Protein A variant can be administered in the same composition. The Protein A variant can also be a recombinant nucleic acid molecule encoding a Protein A variant. A recombinant nucleic acid molecule can encode the Protein A variant and at least one staphylococcal antigen or immunogenic fragment thereof. As used herein, the term “modulate” or “modulation” encompasses the meanings of the words “enhance,” or “inhibit.” “Modulation” of activity may be either an increase or a decrease in activity. As used herein, the term “modulator” refers to compounds that effect the function of a moiety,







TABLE 1-continued

SpA and staphylococcal antigen combinations.						
vWh						+
Coa	+	+	+	+	+	+
Hla		+	+	+	+	+
Hla <sub>H35A</sub>			+	+	+	+
IsdC				+	+	+
SasF					+	+
vWbp						+
vWh						+
Hla	+	+	+	+	+	+
Hla <sub>H35A</sub>			+	+	+	+
IsdC				+	+	+
SasF					+	+
vWbp						+
vWh						+
Hla <sub>H35A</sub>			+	+	+	+
IsdC				+	+	+
SasF					+	+
vWbp						+
vWh						+
IsdC			+	+	+	+
SasF				+	+	+
vWbp						+
vWh						+
SasF				+	+	+
vWbp						+
vWh						+
vWbp						+
vWh						+
vWh						+

In still further aspects, the isolated Protein A variant is multimerized, e.g., dimerized or a linear fusion of two or more polypeptides or peptide segments. In certain aspects of the invention, a composition comprises multimers or concatamers of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more isolated cell surface proteins or segments thereof. Concatamers are linear polypeptides having one or more repeating peptide units. SpA polypeptides or fragments can be consecutive or separated by a spacer or other peptide sequences, e.g., one or more additional bacterial peptide. In a further aspect, the other polypeptides or peptides contained in the multimer or concatamer can include, but are not limited to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 of Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, vWh or immunogenic fragments thereof. Additional staphylococcal antigens that can be used in combination with a Protein A variant include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg2+ transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein.

The term "Protein A variant" or "SpA variant" refers to polypeptides that include a SpA IgG domain having two or more amino acid substitutions that disrupt binding to Fc and V<sub>H</sub>3. In certain aspect, a SpA variant includes a variant domain D peptide, as well as variants of SpA polypeptides

and segments thereof that are non-toxicogenic and stimulate an immune response against *staphylococcus* bacteria Protein A and/or bacteria expressing such.

Embodiments of the present invention include methods for eliciting an immune response against a *staphylococcus* bacterium or staphylococci in a subject comprising providing to the subject an effective amount of a Protein A variant or a segment thereof. In certain aspects, the methods for eliciting an immune response against a *staphylococcus* bacterium or staphylococci in a subject comprising providing to the subject an effective amount of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or more secreted proteins and/or cell surface proteins or segments/fragments thereof. A secreted protein or cell surface protein includes, but is not limited to Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, and/or vWh proteins and immunogenic fragments thereof. Additional staphylococcal antigens that can be used in combination with a Protein A variant include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg2+ transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein.

Embodiments of the invention include compositions that include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%

identical or similar to Protein A, or a second protein or peptide that is a secreted bacterial protein or a bacterial cell surface protein. In a further embodiment of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a Protein A domain D polypeptide (SEQ ID NO:2), domain E (SEQ ID NO:3), domain A (SEQ ID NO:4), domain C (SEQ ID NO:5), domain B (SEQ ID NO:6), or a nucleic acid sequence encoding a Protein A domain D, domain E, domain A, domain C, or domain B polypeptide. In certain aspects a Protein A polypeptide segment will have an amino acid sequence of SEQ ID NO:8. Similarity or identity, with identity being preferred, is known in the art and a number of different programs can be used to identify whether a protein (or nucleic acid) has sequence identity or similarity to a known sequence. Sequence identity and/or similarity is determined using standard techniques known in the art, including, but not limited to, the local sequence identity algorithm of Smith & Waterman (1981), by the sequence identity alignment algorithm of Needleman & Wunsch (1970), by the search for similarity method of Pearson & Lipman (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Drive, Madison, Wis.), the Best Fit sequence program described by Devereux et al. (1984), preferably using the default settings, or by inspection. Preferably, percent identity is calculated by using alignment tools known to and readily ascertainable to those of skill in the art. Percent identity is essentially the number of identical amino acids divided by the total number of amino acids compared times one hundred.

Still further embodiments include methods for stimulating in a subject a protective or therapeutic immune response against a *staphylococcus* bacterium comprising administering to the subject an effective amount of a composition including (i) a SpA variant, e.g., a variant SpA domain D polypeptide or peptide thereof; or, (ii) a nucleic acid molecule encoding such a SpA variant polypeptide or peptide thereof, or (iii) administering a SpA variant domain D polypeptide with any combination or permutation of bacterial proteins described herein. In a preferred embodiment the composition is not a *staphylococcus* bacterium. In certain aspects the subject is a human or a cow. In a further aspect the composition is formulated in a pharmaceutically acceptable formulation. The staphylococci may be *Staphylococcus aureus*.

Yet still further embodiments include vaccines comprising a pharmaceutically acceptable composition having an isolated SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described herein, wherein the composition is capable of stimulating an immune response against a *staphylococcus* bacterium. The vaccine may comprise an isolated SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described. In certain aspects of the invention the isolated SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described are multimerized, e.g., dimerized or concatamerized. In a further aspect, the vaccine composition is contaminated by less than about 10, 9, 8, 7, 6, 5, 4, 3, 2, 1, 0.5, 0.25, 0.05% (or any range derivable therein) of other Staphylococcal proteins. A composition may further comprise an isolated non-SpA polypeptide. Typically the vaccine comprises an adjuvant. In certain aspects a protein or peptide of the invention is linked (covalently or non-covalently) to the adjuvant, preferably the adjuvant is chemically conjugated to the protein.

In still yet further embodiments, a vaccine composition is a pharmaceutically acceptable composition having a recombinant nucleic acid encoding all or part of a SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described herein, wherein the composition is capable of stimulating an immune response against a *staphylococcus* bacteria. The vaccine composition may comprise a recombinant nucleic acid encoding all or part of a SpA variant polypeptide, or any other combination or permutation of protein(s) or peptide(s) described herein. In certain embodiments the recombinant nucleic acid contains a heterologous promoter. Preferably the recombinant nucleic acid is a vector. More preferably the vector is a plasmid or a viral vector. In some aspects the vaccine includes a recombinant, non-staphylococcus bacterium containing the nucleic acid. The recombinant non-staphylococci may be *Salmonella* or another gram-positive bacteria. The vaccine may comprise a pharmaceutically acceptable excipient, more preferably an adjuvant.

Still further embodiments include methods for stimulating in a subject a protective or therapeutic immune response against a *staphylococcus* bacterium comprising administering to the subject an effective amount of a composition of a SpA variant polypeptide or segment/fragment thereof and further comprising one or more of a Eap, Ebh, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, or vWh protein or peptide thereof. In a preferred embodiment the composition comprises a non-staphylococcus bacterium. In a further aspect the composition is formulated in a pharmaceutically acceptable formulation. The staphylococci for which a subject is being treated may be *Staphylococcus aureus*. Methods of the invention also include SpA variant compositions that contain 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or more secreted virulence factors and/or cell surface proteins, such as Eap, Ebh, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, or vWh in various combinations. In certain aspects a vaccine formulation includes Eap, Ebh, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, and vWh. In certain aspects an antigen combination can include (1) a SpA variant and IsdA; (2) SpA variant and ClfB; (3) SpA variant and SdrD; (4) SpA variant and Hla or Hla variant; (5) SpA variant and ClfB, SdrD, and Hla or Hla variant; (6) SpA variant, IsdA, SdrD, and Hla or Hla variant; (7) SpA variant, IsdA, ClfB, and Hla or Hla variant; (8) SpA variant, IsdA, ClfB, and SdrD; (9) SpA variant, IsdA, ClfB, SdrD and Hla or Hla variant; (10) SpA variant, IsdA, ClfB, and SdrD; (11) SpA variant, IsdA, SdrD, and Hla or Hla variant; (12) SpA variant, IsdA, and Hla or Hla variant; (13) SpA variant, IsdA, ClfB, and Hla or Hla variant; (14) SpA variant, ClfB, and SdrD; (15) SpA variant, ClfB, and Hla or Hla variant; or (16) SpA variant, SdrD, and Hla or Hla variant.

In certain aspects, a bacterium delivering a composition of the invention will be limited or attenuated with respect to prolonged or persistent growth or abscess formation. In yet a further aspect, SpA variant(s) can be overexpressed in an attenuated bacterium to further enhance or supplement an immune response or vaccine formulation.

The term "EsxA protein" refers to a protein that includes isolated wild-type EsxA polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria EsxA proteins.

The term "EsxB protein" refers to a protein that includes isolated wild-type EsxB polypeptides from *staphylococcus*

bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria EsxB proteins.

The term “SdrD protein” refers to a protein that includes isolated wild-type SdrD polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria SdrD proteins.

The term “SdrE protein” refers to a protein that includes isolated wild-type SdrE polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria SdrE proteins.

The term “IsdA protein” refers to a protein that includes isolated wild-type IsdA polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria IsdA proteins.

The term “IsdB protein” refers to a protein that includes isolated wild-type IsdB polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria IsdB proteins.

The term “Eap protein” refers to a protein that includes isolated wild-type Eap polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Eap proteins.

The term “Ebh protein” refers to a protein that includes isolated wild-type Ebh polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Ebh proteins.

The term “Emp protein” refers to a protein that includes isolated wild-type Emp polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Emp proteins.

The term “EsaB protein” refers to a protein that includes isolated wild-type EsaB polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria EsaB proteins.

The term “EsaC protein” refers to a protein that includes isolated wild-type EsaC polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria EsaC proteins.

The term “SdrC protein” refers to a protein that includes isolated wild-type SdrC polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria SdrC proteins.

The term “ClfA protein” refers to a protein that includes isolated wild-type ClfA polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria ClfA proteins.

The term “ClfB protein” refers to a protein that includes isolated wild-type ClfB polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria ClfB proteins.

The term “Coa protein” refers to a protein that includes isolated wild-type Coa polypeptides from *staphylococcus*

bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Coa proteins.

The term “Hla protein” refers to a protein that includes isolated wild-type Hla polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria Hla proteins.

The term “IsdC protein” refers to a protein that includes isolated wild-type IsdC polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria IsdC proteins.

The term “SasF protein” refers to a protein that includes isolated wild-type SasF polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria SasF proteins.

The term “vWbp protein” refers to a protein that includes isolated wild-type vWbp (von Willebrand factor binding protein) polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria vWbp proteins.

The term “vWh protein” refers to a protein that includes isolated wild-type vWh (von Willebrand factor binding protein homolog) polypeptides from *staphylococcus* bacteria and segments thereof, as well as variants that stimulate an immune response against *staphylococcus* bacteria vWh proteins.

An immune response refers to a humoral response, a cellular response, or both a humoral and cellular response in an organism. An immune response can be measured by assays that include, but are not limited to, assays measuring the presence or amount of antibodies that specifically recognize a protein or cell surface protein, assays measuring T-cell activation or proliferation, and/or assays that measure modulation in terms of activity or expression of one or more cytokines.

In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an EsxA protein. In certain aspects the EsxA protein will have all or part of the amino acid sequence of SEQ ID NO:11.

In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an EsxB protein. In certain aspects the EsxB protein will have all or part of the amino acid sequence of SEQ ID NO:12.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an SdrD protein. In certain aspects the SdrD protein will have all or part of the amino acid sequence of SEQ ID NO:13.

In further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an SdrE protein. In certain aspects the SdrE protein will have all or part of the amino acid sequence of SEQ ID NO:14.

In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or

99% identical or similar to an IsdA protein. In certain aspects the IsdA protein will have all or part of the amino acid sequence of SEQ ID NO:15.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an IsdB protein. In certain aspects the IsdB protein will have all or part of the amino acid sequence of SEQ ID NO:16.

Embodiments of the invention include compositions that include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a EsaB protein. In certain aspects the EsaB protein will have all or part of the amino acid sequence of SEQ ID NO:17.

In a further embodiment of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a ClfB protein. In certain aspects the ClfB protein will have all or part of the amino acid sequence of SEQ ID NO:18.

In still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an IsdC protein. In certain aspects the IsdC protein will have all or part of the amino acid sequence of SEQ ID NO:19.

In yet further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a SasF protein. In certain aspects the SasF protein will have all or part of the amino acid sequence of SEQ ID NO:20.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a SdrC protein. In certain aspects the SdrC protein will have all or part of the amino acid sequence of SEQ ID NO:21.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a ClfA protein. In certain aspects the ClfA protein will have all or part of the amino acid sequence of SEQ ID NO:22.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an Eap protein. In certain aspects the Eap protein will have all or part of the amino acid sequence of SEQ ID NO:23.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an Ebh protein. In certain aspects the Ebh protein will have all or part of the amino acid sequence of SEQ ID NO:24.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an Emp protein. In certain aspects the Emp protein will have all or part of the amino acid sequence of SEQ ID NO:25.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to an EsaC protein. In certain

aspects the EsaC protein will have all or part of the amino acid sequence of SEQ ID NO:26. Sequence of EsaC polypeptides can be found in the protein databases and include, but are not limited to accession numbers ZP\_02760162 (GI:168727885), NP\_645081.1 (GI:21281993), and NP\_370813.1 (GI:15923279), each of which is incorporated herein by reference as of the priority date of this application.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a Coa protein. In certain aspects the Coa protein will have all or part of the amino acid sequence of SEQ ID NO:27.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a Hla protein. In certain aspects the Hla protein will have all or part of the amino acid sequence of SEQ ID NO:28.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a vWa protein. In certain aspects the vWa protein will have all or part of the amino acid sequence of SEQ ID NO:29.

In yet still further embodiments of the invention a composition may include a polypeptide, peptide, or protein that is or is at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical or similar to a vWbp protein. In certain aspects the vWbp protein will have all or part of the amino acid sequence of SEQ ID NO:32.

In certain aspects, a polypeptide or segment/fragment can have a sequence that is at least 85%, at least 90%, at least 95%, at least 98%, or at least 99% or more identical to the amino acid sequence of the reference polypeptide. The term "similarity" refers to a polypeptide that has a sequence that has a certain percentage of amino acids that are either identical with the reference polypeptide or constitute conservative substitutions with the reference polypeptides.

The polypeptides described herein may include 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or more variant amino acids within at least, or at most 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 300, 400, 500, 550, 1000 or more contiguous amino acids, or any range derivable therein, of SEQ ID NO:2-30, or SEQ ID NO:32-34.

A polypeptide segment as described herein may include 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73,

74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 300, 400, 500, 550, 1000 or more contiguous amino acids, or any range derivable therein, of SEQ ID NO:2-30, or SEQ ID NO:33-34.

The compositions may be formulated in a pharmaceutically acceptable composition. In certain aspects of the invention the *staphylococcus* bacterium is an *S. aureus* bacterium.

In further aspects, a composition may be administered more than one time to the subject, and may be administered 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20 or more times. The administration of the compositions include, but is not limited to oral, parenteral, subcutaneous, intramuscular, intravenous, or various combinations thereof, including inhalation or aspiration.

In still further embodiments, a composition comprises a recombinant nucleic acid molecule encoding a polypeptide described herein or segments/fragments thereof. Typically a recombinant nucleic acid molecule encoding a polypeptide described herein contains a heterologous promoter. In certain aspects, a recombinant nucleic acid molecule of the invention is a vector, in still other aspects the vector is a plasmid. In certain embodiments the vector is a viral vector. In certain aspects a composition includes a recombinant, non-*staphylococcus* bacterium containing or expressing a polypeptide described herein. In particular aspects the recombinant non-staphylococcus bacteria is *Salmonella* or another gram-positive bacteria. A composition is typically administered to mammals, such as human subjects, but administration to other animals that are capable of eliciting an immune response is contemplated. In further aspects the *staphylococcus* bacterium containing or expressing the polypeptide is *Staphylococcus aureus*. In further embodiments the immune response is a protective immune response.

In further embodiments a composition comprises a recombinant nucleic acid molecule encoding all or part of one or more of a Eap, Ehb, Emp, EsaB, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, SpA, vWbp, or vWh protein or peptide or variant thereof. Additional staphylococcal antigens that can be used in combination with the polypeptides described herein include, but are not limited to 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-

1, SSP-2, and/or Vitronectin binding protein. In particular aspects, a bacteria is a recombinant non-staphylococcus bacterium, such as a *Salmonella* or other gram-positive bacteria.

Compositions of the invention are typically administered to human subjects, but administration to other animals that are capable of eliciting an immune response to a *staphylococcus* bacterium is contemplated, particularly cattle, horses, goats, sheep and other domestic animals, i.e., mammals.

In certain aspects the *staphylococcus* bacterium is a *Staphylococcus aureus*. In further embodiments the immune response is a protective immune response. In still further aspects, the methods and compositions of the invention can be used to prevent, ameliorate, reduce, or treat infection of tissues or glands, e.g., mammary glands, particularly mastitis and other infections. Other methods include, but are not limited to prophylactically reducing bacterial burden in a subject not exhibiting signs of infection, particularly those subjects suspected of or at risk of being colonized by a target bacteria, e.g., patients that are or will be at risk or susceptible to infection during a hospital stay, treatment, and/or recovery.

Any embodiment discussed with respect to one aspect of the invention applies to other aspects of the invention as well. In particular, any embodiment discussed in the context of a SpA variant polypeptide or peptide or nucleic acid may be implemented with respect to other antigens, such as Eap, Ehb, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, vWh, 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein (or nucleic acids), and vice versa. It is also understood that any one or more of Eap, Ehb, Emp, EsaC, EsxA, EsxB, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, Coa, Hla, IsdC, SasF, vWbp, vWh, 52 kDa vitronectin binding protein (WO 01/60852), Aaa, Aap, Ant, autolysin glucosaminidase, autolysin amidase, Cna, collagen binding protein (U.S. Pat. No. 6,288,214), EFB (FIB), Elastin binding protein (EbpS), EPB, FbpA, fibrinogen binding protein (U.S. Pat. No. 6,008,341), Fibronectin binding protein (U.S. Pat. No. 5,840,846), FnbA, FnbB, GehD (US 2002/0169288), HarA, HBP, Immunodominant ABC transporter, IsaA/PisA, laminin receptor, Lipase GehD, MAP, Mg<sup>2+</sup> transporter, MHC II analogue (U.S. Pat. No. 5,648,240), MRPII, Npase, RNA III activating protein (RAP), SasA, SasB, SasC, SasD, SasK, SBI, SdrF (WO 00/12689), SdrG/Fig (WO 00/12689), SdrH (WO 00/12689), SEA exotoxins (WO 00/02523), SEB exotoxins (WO 00/02523), SitC and Ni ABC transporter, SitC/MntC/saliva binding protein (U.S. Pat. No. 5,801,234), SsaA, SSP-1, SSP-2, and/or Vitronectin binding protein can be specifically excluded from a claimed composition.

Embodiments of the invention include compositions that contain or do not contain a bacterium. A composition may or may not include an attenuated or viable or intact staphylococcal bacterium. In certain aspects, the composition comprises a bacterium that is not a staphylococcal bacterium or does not

contain staphylococcal bacteria. In certain embodiments a bacterial composition comprises an isolated or recombinantly expressed staphylococcal Protein A variant or a nucleotide encoding the same. The composition may be or include a recombinantly engineered *staphylococcus* bacterium that has been altered in a way that comprises specifically altering the bacterium with respect to a secreted virulence factor or cell surface protein. For example, the bacteria may be recombinantly modified to express more of the virulence factor or cell surface protein than it would express if unmodified.

The term “isolated” can refer to a nucleic acid or polypeptide that is substantially free of cellular material, bacterial material, viral material, or culture medium (when produced by recombinant DNA techniques) of their source of origin, or chemical precursors or other chemicals (when chemically synthesized). Moreover, an isolated compound refers to one that can be administered to a subject as an isolated compound; in other words, the compound may not simply be considered “isolated” if it is adhered to a column or embedded in an agarose gel. Moreover, an “isolated nucleic acid fragment” or “isolated peptide” is a nucleic acid or protein fragment that is not naturally occurring as a fragment and/or is not typically in the functional state.

Moieties of the invention, such as polypeptides, peptides, antigens, or immunogens, may be conjugated or linked covalently or noncovalently to other moieties such as adjuvants, proteins, peptides, supports, fluorescence moieties, or labels. The term “conjugate” or “immunoconjugate” is broadly used to define the operative association of one moiety with another agent and is not intended to refer solely to any type of operative association, and is particularly not limited to chemical “conjugation.” Recombinant fusion proteins are particularly contemplated. Compositions of the invention may further comprise an adjuvant or a pharmaceutically acceptable excipient. An adjuvant may be covalently or noncovalently coupled to a polypeptide or peptide of the invention. In certain aspects, the adjuvant is chemically conjugated to a protein, polypeptide, or peptide.

The term “providing” is used according to its ordinary meaning to indicate “to supply or furnish for use.” In some embodiments, the protein is provided directly by administering the protein, while in other embodiments, the protein is effectively provided by administering a nucleic acid that encodes the protein. In certain aspects the invention contemplates compositions comprising various combinations of nucleic acid, antigens, peptides, and/or epitopes.

The subject will have (e.g., are diagnosed with a staphylococcal infection), will be suspected of having, or will be at risk of developing a staphylococcal infection. Compositions of the present invention include immunogenic compositions wherein the antigen(s) or epitope(s) are contained in an amount effective to achieve the intended purpose. More specifically, an effective amount means an amount of active ingredients necessary to stimulate or elicit an immune response, or provide resistance to, amelioration of, or mitigation of infection. In more specific aspects, an effective amount prevents, alleviates or ameliorates symptoms of disease or infection, or prolongs the survival of the subject being treated. Determination of the effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. For any preparation used in the methods of the invention, an effective amount or dose can be estimated initially from in vitro studies, cell culture, and/or animal model assays. For example, a dose can be formulated in animal models to achieve a desired immune

response or circulating antibody concentration or titer. Such information can be used to more accurately determine useful doses in humans.

The embodiments in the Example section are understood to be embodiments of the invention that are applicable to all aspects of the invention.

The use of the term “or” in the claims is used to mean “and/or” unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and “and/or.” It is also contemplated that anything listed using the term “or” may also be specifically excluded.

Throughout this application, the term “about” is used to indicate that a value includes the standard deviation of error for the device or method being employed to determine the value.

Following long-standing patent law, the words “a” and “an,” when used in conjunction with the word “comprising” in the claims or specification, denotes one or more, unless specifically noted.

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating specific embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

## DESCRIPTION OF THE DRAWINGS

So that the matter in which the above-recited features, advantages and objects of the invention as well as others which will become clear are attained and can be understood in detail, more particular descriptions and certain embodiments of the invention briefly summarized above are illustrated in the appended drawings. These drawings form a part of the specification. It is to be noted, however, that the appended drawings illustrate certain embodiments of the invention and therefore are not to be considered limiting in their scope.

FIGS. 1A-1B. (FIG. 1A) Primary structure of the Protein A precursor with an N-terminal YSIRK motif signal peptide, five immunoglobulin binding domains as tandem repeats designated E, D, A, B, C, region X, and the LPXTG sorting signal. (FIG. 1B) Following synthesis of the Protein A precursor, staphylococci secrete this product via the Sec pathway, and sortase A cleaves the LPXTG sorting signal between the T and G residues. Nucleophilic attack of the amino group within lipid II at the sortase-Protein A thioester-linked intermediate forms the amide bond that links Protein A to the cell wall envelope and enables its display on the bacterial surface.

FIG. 2. Three dimensional model of the molecular interactions between the SpA-domain D of Protein A, the VH3 Fab domain of the B cell receptor, and of the Fcγ domain of immunoglobulin. The model is derived from two crystal structures (Graille et al., 2000 and Gouda et al., 1992) that revealed side chain residues involved in the formation of ionic bonds that enable these complexes. Gln-9 and Gln-10 of SpA-D promote binding to Fcγ, whereas Asp-36 and Asp-37 enable complex formation with VH3 Fab.

FIG. 3. Left panel—Coomassie Blue stained SDS-PAGE reveals the migrational position of purified His-tagged SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub>, human IgG, and sortase A (SrtA), a control protein. Right panel—Coomassie Blue stained SDS-PAGE to reveal the elution of Protein A immunoglobulin complexes eluted following affinity chromatogra-

25

phy of human IgG on Ni-NTA columns pre-charged with His-tagged SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub> or SrtA.

FIG. 4. ELISA assays to quantify human immunoglobulin (hIgG), human F(ab)<sub>2</sub> IgG fragments and human Fc fragments of immunoglobulin (hFc). Plates were coated with equal amounts of His-tagged SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub> or SrtA. hIgG-HRP, F(ab)<sub>2</sub>-HRP and hFc-HRP were added onto the plates and incubated for an hour. Absorbance at 450 nm was recorded and plotted to determine the half maximal titers.

FIG. 5. Purified SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub> or a PBS mock control were injected into the peritoneum of mice and analyzed for their ability to reduce the B cell population in the spleen of experimental BALB/c mice. Animals were killed 4 hours following injection, their spleen removed, tissue homogenized and stained with CD 19 antibodies directed against B cells. The number of B cells was quantified by FACS sorting.

FIG. 6 Generation of a non-toxicogenic protein A vaccine. a, Translational protein A (SpA) product of *S. aureus* Newman and USA300 LAC with an N-terminal signal peptide (white box), five immunoglobulin binding domains (IgBDs designated E, D, A, B and C), variable region X and C-terminal sorting signal (black box). b, Amino acid sequence of the five IgBDs as well as nontoxicogenic SpA-D<sub>KKAA</sub>, with the positions of triple  $\alpha$ -helical bundles (H1, H2 and H3) as well as glutamine (Q) 9, 10 and aspartate (D) 36, 37 indicated. c, Coomassie Blue-stained SDS-PAGE of SpA, SpA-D, SpA-D<sub>KKAA</sub> or SrtA purified on Ni-NTA sepharose in the presence or absence of human immunoglobulin (hIgG). d, ELISA examining the association of immobilized SpA, SpA-D or SpA-D<sub>KKAA</sub> with human IgG as well as its Fc or F(ab)<sub>2</sub> fragments and von Willebrand factor (vWF). e, CD19+ B lymphocytes in splenic tissue of BALB/c mice that had been mock immunized or treated with SpA-D or SpA-D<sub>KKAA</sub> were quantified by FACS.

FIG. 7 Non-toxicogenic protein A vaccine prevents abscess formation. Histopathology of renal tissue isolated during necropsy of BALB/c mice that had been mock immunized (PBS) or vaccinated with SpA, SpA-D as well as SpA-D<sub>KKAA</sub> and challenged with *S. aureus* Newman. Thin sectioned tissues were stained with hematoxylin-eosin. White arrows identify polymorphonuclear leukocyte (PMN) infiltrates. Dark arrows identify staphylococcal abscess communities.

FIG. 8 Antibodies raised by the non-toxicogenic protein A vaccine block the B cell superantigen function of SpA. a, Rabbit antibodies raised against SpA-D<sub>KKAA</sub> were purified on a matrix with immobilized antigen and analyzed by Coomassie Blue-stained SDS-PAGE. Antibodies were cleaved with pepsin and F(ab)<sub>2</sub> fragments were purified by a second round of affinity chromatography on SpA-D<sub>KKAA</sub> matrix. b, SpA-D<sub>KKAA</sub> specific F(ab)<sub>2</sub> interfere with the binding of SpA or SpA-D to human immunoglobulin (hIgG) or, c, to von Willebrand Factor (vWF).

FIG. 9 Full-length non-toxicogenic protein A generates improved immune responses. a, Full-length SpA<sub>KKAA</sub> was purified on Ni-NTA sepharose and analyzed by Coomassie-Blue stained SDS-PAGE. b, CD 19+ B lymphocytes in splenic tissue of BALB/c mice that had been mock immunized or treated with SpA or SpA<sub>KKAA</sub> were quantified by FACS. c, ELISA examining the association of immobilized SpA or SpA<sub>KKAA</sub> with human IgG as well as its Fc or F(ab)<sub>2</sub> fragments or von Willebrand factor (vWF). d, Human or mouse serum antibody titers to diphtheria toxoid (CRM197) and non-toxicogenic SpA<sub>KKAA</sub> or SpA-D<sub>KKAA</sub>. Human volunteers with a history of DTaP immunization and staphylococcal infection (n=16) as well as mice (n=20) that had been

26

infected with *S. aureus* Newman or USA 300 LAC or immunized with SpA<sub>KKAA</sub> or SpA-D<sub>KKAA</sub> were examined by quantitative dot blot.

FIG. 10 Staphylococcal infection does not generate protective immunity. BALB/c mice (n=20) were infected with *S. aureus* Newman or mock challenged (PBS) for thirty days and infection cleared with chloramphenicol treatment. Both cohorts of animals were then challenged with *S. aureus* Newman and bacterial load (CFU) in kidney tissue homogenate analyzed following necropsy on day 4.

FIG. 11 Comparison of abscess formation in mice treated with PBS, SpA, SpA-D, and SpA-D<sub>KKAA</sub>.

FIGS. 12A-12C (A) ELISA examining the association of immobilized SpA, SpA-D, SpA-DKKAA or SpA-DGGSS with human IgG as well as its Fc or F(ab)<sub>2</sub> fragments and IgM. Statistical significance of SpA-DKKAA and SpA-DGGSS binding to each ligand was compared against SpA-D; SpA-D binding was compared against SpA (n=3); \* signifies P<0.05; \*\* signifies P<0.01. (B) ELISA examining the level of cross-reactive antibodies of hyper-immune sera samples collected from actively immunized mice (n=5) with SpA-D, SpA-DKKAA and SpA-DGGSS. (C) Abscess formation in mice treated with PBS, SpA-D, SpA-D<sub>KKAA</sub> and SpA-D<sub>GGSS</sub>.

#### DETAILED DESCRIPTION

*Staphylococcus aureus* is a commensal of the human skin and nares, and the leading cause of bloodstream, skin and soft tissue infections (Klevens et al., 2007). Recent dramatic increases in the mortality of staphylococcal diseases are attributed to the spread of methicillin-resistant *S. aureus* (MRSA) strains often not susceptible to antibiotics (Kennedy et al., 2008). In a large retrospective study, the incidence of MRSA infections was 4.6% of all hospital admissions in the United States (Klevens et al., 2007). The annual health care costs for 94,300 MRSA infected individuals in the United States exceed \$2.4 billion (Klevens et al., 2007). The current MRSA epidemic has precipitated a public health crisis that needs to be addressed by development of a preventive vaccine (Boucher and Corey, 2008). To date, an FDA licensed vaccine that prevents *S. aureus* diseases is not available.

The inventors describe here the use of Protein A, a cell wall anchored surface protein of staphylococci, for the generation of variants that can serve as subunit vaccines. The pathogenesis of staphylococcal infections is initiated as bacteria invade the skin or blood stream via trauma, surgical wounds, or medical devices (Lowy, 1998). Although the invading pathogen may be phagocytosed and killed, staphylococci can also escape innate immune defenses and seed infections in organ tissues, inducing inflammatory responses that attract macrophages, neutrophils, and other phagocytes (Lowy, 1998). The responsive invasion of immune cells to the site of infection is accompanied by liquefaction necrosis as the host seeks to prevent staphylococcal spread and allow for removal of necrotic tissue debris (Lam et al., 1963). Such lesions can be observed by microscopy as hypercellular areas containing necrotic tissue, leukocytes, and a central nidus of bacteria (Lam et al., 1963). Unless staphylococcal abscesses are surgically drained and treated with antibiotics, disseminated infection and septicemia produce a lethal outcome (Sheagren, 1984).

## II. STAPHYLOCOCCAL ANTIGENS

## A. Staphylococcal Protein A (SpA)

All *Staphylococcus aureus* strains express the structural gene for Protein A (spa) (Jensen, 1958; Said-Salim et al., 2003), a well characterized virulence factor whose cell wall anchored surface protein product (SpA) encompasses five highly homologous immunoglobulin binding domains designated E, D, A, B, and C (Sjodahl, 1977). These domains display ~80% identity at the amino acid level, are 56 to 61 residues in length, and are organized as tandem repeats (Uhlen et al., 1984). SpA is synthesized as a precursor protein with an N-terminal YSIRK/GS signal peptide and a C-terminal LPXTG motif sorting signal (DeDent et al., 2008; Schneewind et al., 1992). Cell wall anchored Protein A is displayed in great abundance on the staphylococcal surface (DeDent et al., 2007; Sjoquist et al., 1972). Each of its immunoglobulin binding domains is composed of anti-parallel  $\alpha$ -helices that assemble into a three helix bundle and bind the Fc domain of immunoglobulin G (IgG) (Deisenhofer, 1981; Deisenhofer et al., 1978), the VH3 heavy chain (Fab) of IgM (i.e., the B cell receptor) (Graille et al., 2000), the von Willebrand factor at its A1 domain [vWF A1 is a ligand for platelets] (O'Seaghda et al., 2006) and the tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ) receptor I (TNFRI) (Gomez et al., 2006), which is displayed on surfaces of airway epithelia (Gomez et al., 2004; Gomez et al., 2007).

SpA impedes neutrophil phagocytosis of staphylococci through its attribute of binding the Fc component of IgG (Jensen, 1958; Uhlen et al., 1984). Moreover, SpA is able to activate intravascular clotting via its binding to von Willebrand factor A1 domains (Hartleib et al., 2000). Plasma proteins such as fibrinogen and fibronectin act as bridges between staphylococci (ClfA and ClfB) and the platelet integrin GPIIb/IIIa (O'Brien et al., 2002), an activity that is supplemented through Protein A association with vWF A1, which allows staphylococci to capture platelets via the GPIIb- $\alpha$  platelet receptor (Foster, 2005; O'Seaghda et al., 2006). SpA also binds TNFRI and this interaction contributes to the pathogenesis of staphylococcal pneumonia (Gomez et al., 2004). SpA activates proinflammatory signaling through TNFRI mediated activation of TRAF2, the p38/c-Jun kinase, mitogen activate protein kinase (MAPK) and the Rel-transcription factor NF-KB. SpA binding further induces TNFRI shedding, an activity that appears to require the TNF-converting enzyme (TACE) (Gomez et al., 2007). All of the aforementioned SpA activities are mediated through its five IgG binding domains and can be perturbed by the same amino acid substitutions, initially defined by their requirement for the interaction between Protein A and human IgG1 (Cedergren et al., 1993).

SpA also functions as a B cell superantigen by capturing the Fab region of VH3 bearing IgM, the B cell receptor (Gomez et al., 2007; Goodyear et al., 2003; Goodyear and Silverman, 2004; Roben et al., 1995). Following intravenous challenge, staphylococcal Protein A (SpA) mutations show a reduction in staphylococcal load in organ tissues and dramatically diminished ability to form abscesses (described herein). During infection with wildtype *S. aureus*, abscesses are formed within forty-eight hours and are detectable by light microscopy of hematoxylin-eosin stained, thin-sectioned kidney tissue, initially marked by an influx of polymorphonuclear leukocytes (PMNs). On day 5 of infection, abscesses increase in size and enclosed a central population of staphylococci, surrounded by a layer of eosinophilic, amorphous material and a large cuff of PMNs. Histopathology revealed massive necrosis of PMNs in proximity to the staphylococcal

nidus at the center of abscess lesions as well as a mantle of healthy phagocytes. The inventors also observed a rim of necrotic PMNs at the periphery of abscess lesions, bordering the eosinophilic pseudocapsule that separated healthy renal tissue from the infectious lesion. Staphylococcal variants lacking Protein A are unable to establish the histopathology features of abscesses and are cleared during infection.

In previous studies, Cedergren et al. (1993) engineered five individual substitutions in the Fc fragment binding sub-domain of the B domain of SpA, L17D, N28A, I31A and K35A. These authors created these proteins to test data gathered from a three dimensional structure of a complex between one domain of SpA and Fc<sub>1</sub>. Cedergren et al. determined the effects of these mutations on stability and binding, but did not contemplate use of such substitutions for the production of a vaccine antigen.

Brown et al. (1998) describe studies designed to engineer new proteins based on SpA that allow the use of more favorable elution conditions when used as affinity ligands. The mutations studied included single mutations of Q13A, Q14H, N15A, N15H, F17H, Y18F, L21H, N32H, or K39H. Brown et al. report that Q13A, N15A, N15H, and N32H substitutions made little difference to the dissociation constant values and that the Y18F substitution resulted in a 2 fold decrease in binding affinity as compared to wild type SpA. Brown et al. also report that L21H and F17H substitutions decrease the binding affinity by five-fold and a hundred-fold respectively. The authors also studied analogous substitutions in two tandem domains. Thus, the Brown et al. studies were directed to generating a SpA with a more favorable elution profile, hence the use of His substitutions to provide a pH sensitive alteration in the binding affinity. Brown et al. is silent on the use of SpA as a vaccine antigen.

Graille et al. (2000) describe a crystal structure of domain D of SpA and the Fab fragment of a human IgM antibody. Graille et al. define by analysis of a crystal structure the D domain amino acid residues that interact with the Fab fragment as residues Q26, G29, F30, Q32, S33, D36, D37, Q40, N43, E47, or L51, as well as the amino acid residues that form the interface between the domain D sub-domains. Graille et al. define the molecular interactions of these two proteins, but is silent in regard to any use of substitutions in the interacting residues in producing a vaccine antigen.

O'Seaghda et al. (2006) describe studies directed at elucidating which sub-domain of domain D binds vWF. The authors generated single mutations in either the Fc or VH3 binding sub-domains, i.e., amino acid residues F5A, Q9A, Q10A, F13A, Y14A, L17A, N28A, I31A, K35A, G29A, F30A, S33A, D36A, D37A, Q40A, E47A, or Q32A. The authors discovered that vWF binds the same sub-domain that binds Fc. O'Seaghda et al. define the sub-domain of domain D responsible for binding vWF, but is silent in regard to any use of substitutions in the interacting residues in producing a vaccine antigen.

Gomez et al. (2006) describe the identification of residues responsible for activation of the TNFRI by using single mutations of F5A, F13A, Y14A, L17A, N21A, I31A, Q32A, and K35A. Gomez et al. is silent in regard to any use of substitutions in the interacting residues in producing a vaccine antigen.

Recombinant affinity tagged Protein A, a polypeptide encompassing the five IgG domains (EDCAB) (Sjodahl, 1977) but lacking the C-terminal Region X (Guss et al., 1984), was purified from recombinant *E. coli* and used as a vaccine antigen (Stranger-Jones et al., 2006). Because of the attributes of SpA in binding the Fc portion of IgG, a specific humoral immune response to Protein A could not be mea-

sured (Stranger-Jones et al., 2006). The inventors have overcome this obstacle through the generation of SpA-DQ9,10K; D36,37A. BALB/c mice immunized with recombinant Protein A (SpA) displayed significant protection against intravenous challenge with *S. aureus* strains: a 2.951 log reduction in staphylococcal load as compared to the wild-type ( $P > 0.005$ ; Student's t-test) (Stranger-Jones et al., 2006). SpA specific antibodies may cause phagocytic clearance prior to abscess formation and/or impact the formation of the aforementioned eosinophilic barrier in abscesses that separate staphylococcal communities from immune cells since these do not form during infection with Protein A mutant strains. Each of the five SpA domains (i.e., domains formed from three helix bundles designated E, D, A, B, and C) exerts similar binding properties (Jansson et al., 1998). The solution and crystal structure of the domain D has been solved both with and without the Fc and VH3 (Fab) ligands, which bind Protein A in a non-competitive manner at distinct sites (Graille et al., 2000). Mutations in residues known to be involved in IgG binding (F5, Q9, Q10, S11, F13, Y14, L17, N28, I31 and K35) are also required for vWF A1 and TNFR1 binding (Cedergren et al., 1993; Gomez et al., 2006; O'Seaghda et al., 2006), whereas residues important for the VH3 interaction (Q26, G29, F30, S33, D36, D37, Q40, N43, E47) appear to have no impact on the other binding activities (Graille et al., 2000; Jansson et al., 1998). SpA specifically targets a subset of B cells that express VH3 family related IgM on their surface, i.e., VH3 type B cell receptors (Roben et al., 1995). Upon interaction with SpA, these B cells proliferate and commit to apoptosis, leading to preferential and prolonged deletion of innate-like B lymphocytes (i.e., marginal zone B cells and follicular B2 cells) (Goodyear et al., 2003; Goodyear et al., 2004).

Molecular Basis of Protein A Surface Display and Function.

Protein A is synthesized as a precursor in the bacterial cytoplasm and secreted via its YSIRK signal peptide at the cross wall, i.e. the cell division septum of staphylococci (FIG. 1) (DeDent et al., 2007; DeDent et al., 2008). Following cleavage of the C-terminal LPXTG sorting signal, Protein A is anchored to bacterial peptidoglycan crossbridges by sortase A (Mazmanian et al., 1999; Schneewind et al., 1995; Mazmanian et al., 2000). Protein A is the most abundant surface protein of staphylococci; the molecule is expressed by virtually all *S. aureus* strains (Cespedes et al., 2005; Kennedy et al., 2008; Said-Salim et al., 2003). Staphylococci turn over 15-20% of their cell wall per division cycle (Navarre and Schneewind, 1999). Murine hydrolases cleave the glycan strands and wall peptides of peptidoglycan, thereby releasing Protein A with its attached C-terminal cell wall disaccharide tetrapeptide into the extracellular medium (Ton-That et al., 1999). Thus, by physiological design, Protein A is both anchored to the cell wall and displayed on the bacterial surface but also released into surrounding tissues during host infection (Marraffini et al., 2006).

Protein A captures immunoglobulins on the bacterial surface and this biochemical activity enables staphylococcal escape from host innate and acquired immune responses (Jensen, 1958; Goodyear et al., 2004). Interestingly, region X of Protein A (Guss et al., 1984), a repeat domain that tethers the IgG binding domains to the LPXTG sorting signal /cell wall anchor, is perhaps the most variable portion of the staphylococcal genome (Said-Salim, 2003; Schneewind et al., 1992). Each of the five immunoglobulin binding domains of Protein A (SpA), formed from three helix bundles and designated E, D, A, B, and C, exerts similar structural and functional properties (Sjodahl, 1977; Jansson et al., 1998). The

solution and crystal structure of the domain D has been solved both with and without the Fc and V<sub>H</sub>3 (Fab) ligands, which bind Protein A in a non-competitive manner at distinct sites (Graille 2000).

In the crystal structure complex, the Fab interacts with helix II and helix III of domain D via a surface composed of four VH region  $\beta$ -strands (Graille 2000). The major axis of helix II of domain D is approximately 50° to the orientation of the strands, and the interhelical portion of domain D is most proximal to the C0 strand. The site of interaction on Fab is remote from the Ig light chain and the heavy chain constant region. The interaction involves the following domain D residues: Asp-36 of helix II, Asp-37 and Gln-40 in the loop between helix II and helix III and several other residues (Graille 2000). Both interacting surfaces are composed predominantly of polar side chains, with three negatively charged residues on domain D and two positively charged residues on the 2A2 Fab buried by the interaction, providing an overall electrostatic attraction between the two molecules. Of the five polar interactions identified between Fab and domain D, three are between side chains. A salt bridge is formed between Arg-H19 and Asp-36 and two hydrogen bonds are made between Tyr-H59 and Asp-37 and between Asn-H82a and Ser-33. Because of the conservation of Asp-36 and Asp-37 in all five IgG binding domains of Protein A, the inventors mutated these residues.

The SpA-D sites responsible for Fab binding are structurally separate from the domain surface that mediates Fc $\gamma$  binding. The interaction of Fc $\gamma$  with domain D primarily involves residues in helix I with lesser involvement of helix II (Gouda et al., 1992; Deisenhofer, 1981). With the exception of the Gln-32, a minor contact in both complexes, none of the residues that mediate the Fc $\gamma$  interaction are involved in Fab binding. To examine the spatial relationship between these different Ig-binding sites, the SpA domains in these complexes have been superimposed to construct a model of a complex between Fab, the SpA-domain D, and the Fc $\gamma$  molecule. In this ternary model, Fab and Fc $\gamma$  form a sandwich about opposite faces of the helix II without evidence of steric hindrance of either interaction. These findings illustrate how, despite its small size (i.e., 56-61 aa), an SpA domain can simultaneously display both activities, explaining experimental evidence that the interactions of Fab with an individual domain are noncompetitive. Residues for the interaction between SpA-D and Fc $\gamma$  are Gln-9 and Gln-10.

In contrast, occupancy of the Fc portion of IgG on the domain D blocks its interaction with vWF A1 and probably also TNFR1 (O'Seaghda et al., 2006). Mutations in residues essential for IgG Fc binding (F5, Q9, Q10, S11, F13, Y14, L17, N28, I31 and K35) are also required for vWF A1 and TNFR1 binding (O'Seaghda et al., 2006; Cedergren et al., 1993; Gomez et al., 2006), whereas residues critical for the VH3 interaction (Q26, G29, F30, S33, D36, D37, Q40, N43, E47) have no impact on the binding activities of IgG Fc, vWF A1 or TNFR1 (Jansson et al., 1998; Graille et al., 2000). The Protein A immunoglobulin Fab binding activity targets a subset of B cells that express V<sub>H</sub>3 family related IgM on their surface, i.e., these molecules function as VH3 type B cell receptors (Roben et al., 1995). Upon interaction with SpA, these B cells rapidly proliferate and then commit to apoptosis, leading to preferential and prolonged deletion of innate-like B lymphocytes (i.e., marginal zone B cells and follicular B2 cells) (Goodyear and Silverman, 2004; Goodyear and Silverman, 2003). More than 40% of circulating B cells are targeted by the Protein A interaction and the V<sub>H</sub>3 family represents the largest family of human B cell receptors to impart protective humoral responses against pathogens (Goodyear and Silver-

man, 2004; Goodyear and Silverman, 2003). Thus, Protein A functions analogously to staphylococcal superantigens (Roben et al., 1995), albeit that the latter class of molecules, for example SEB, TSST-1, TSST-2, form complexes with the T cell receptor to inappropriately stimulate host immune responses and thereby precipitating characteristic disease features of staphylococcal infections (Roben et al., 1995; Tiedemann et al., 1995). Together these findings document the contributions of Protein A in establishing staphylococcal infections and in modulating host immune responses.

In sum, Protein A domains can be viewed as displaying two different interfaces for binding with host molecules and any development of Protein A based vaccines must consider the generation of variants that do not perturb host cell signaling, platelet aggregation, sequestration of immunoglobulins or the induction of B cell proliferation and apoptosis. Such Protein A variants should also be useful in analyzing vaccines for the ability of raising antibodies that block the aforementioned SpA activities and occupy the five repeat domains at their dual binding interfaces. This goal is articulated and pursued here for the first time and methods are described in detail for the generation of Protein A variants that can be used as a safe vaccine for humans. To perturb IgG Fc $\gamma$ , vWF AI and TNFR1 binding, glutamine (Q) 9 and 10 [numbering derived from the SpA domain D as described in Uhlen et al., 1984] were mutated, and generated lysine substitutions for both glutamines with the expectation that these abolish the ligand attributes at the first binding interface. To perturb IgM Fab VH3 binding, aspartate (D) 36 and 37 were mutated, each of which is required for the association with the B cell receptor. D36 and D37 were both substituted with alanine. Q9, 10K and D36, 37A mutations are here combined in the recombinant molecule SpA-DQ9, 10K; D36, 37A and tested for the binding attributes of Protein A. Further, SpA-D and SpA-DQ9, 10K; D36, 37A are subjected to immunization studies in mice and rabbits and analyzed for [1] the production of specific antibodies (SpA-D Ab); [2] the ability of SpA-D Ab to block the association between Protein A and its four different ligands; and, [3] the attributes of SpA-D Ab to generate protective immunity against staphylococcal infections. (See Examples section below).

#### B. Staphylococcal Coagulases

Coagulases are enzymes produced by *Staphylococcus* bacteria that convert fibrinogen to fibrin. Coa and vW<sub>h</sub> activate prothrombin without proteolysis (Friedrich et al., 2003). The coagulase-prothrombin complex recognizes fibrinogen as a specific substrate, converting it directly into fibrin. The crystal structure of the active complex revealed binding of the D1 and D2 domains to prothrombin and insertion of its Ile<sup>16</sup>-Val<sup>2</sup> N-terminus into the Ile<sup>16</sup> pocket, inducing a functional active site in the zymogen through conformational change (Friedrich et al., 2003). Exosite I of  $\alpha$ -thrombin, the fibrinogen recognition site, and proexosite I on prothrombin are blocked by the D2 of Coa (Friedrich et al., 2003). Nevertheless, association of the tetrameric (Coa-prothrombin)<sub>2</sub> complex binds fibrinogen at a new site with high affinity (Panizzi et al., 2006). This model explains the coagulant properties and efficient fibrinogen conversion by coagulase (Panizzi et al., 2006).

Fibrinogen is a large glycoprotein (Mr ~340,000), formed by three pairs of A $\alpha$ -, B $\beta$ -, and  $\gamma$ -chains covalently linked to form a "dimer of trimers," where A and B designate the fibrinopeptides released by thrombin cleavage (Panizzi et al., 2006). The elongated molecule folds into three separate domains, a central fragment E that contains the N-termini of all six chains and two flanking fragments D formed mainly by the C-termini of the B $\beta$ - and  $\gamma$ -chains. These globular

domains are connected by long triple-helical structures. Coagulase-prothrombin complexes, which convert human fibrinogen to the self-polymerizing fibrin, are not targeted by circulating thrombin inhibitors (Panizzi et al., 2006). Thus, staphylococcal coagulases bypass the physiological blood coagulation pathway.

All *S. aureus* strains secrete coagulase and vWbp (Bjerker-torp et al., 2004; Field and Smith, 1945). Although early work reported important contributions of coagulase to the pathogenesis of staphylococcal infections (Ekstedt and Yotis, 1960; Smith et al., 1947), more recent investigations with molecular genetics tools challenged this view by observing no virulence phenotypes with endocarditis, skin abscess and mastitis models in mice (Moreillon et al., 1995; Phonimdaeng et al., 1990). Generating isogenic variants of *S. aureus* Newman, a fully virulent clinical isolate (Duthie et al., 1952), it is described herein that coa mutants indeed display virulence defects in a lethal bacteremia and renal abscess model in mice. In the inventors experience, *S. aureus* 8325-4 is not fully virulent and it is presumed that mutational lesions in this strain may not be able to reveal virulence defects in vivo. Moreover, antibodies raised against Coa or vWbp perturb the pathogenesis of *S. aureus* Newman infections to a degree mirroring the impact of gene deletions. Coa and vWbp contribute to staphylococcal abscess formation and lethal bacteremia and may also function as protective antigens in subunit vaccines.

Biochemical studies document the biological value of antibodies against Coa and vWbp. By binding to antigen and blocking its association with clotting factors, the antibodies prevent the formation of Coa-prothrombin and vWbp-prothrombin complexes. Passive transfer studies revealed protection of experimental animals against staphylococcal abscess formation and lethal challenge by Coa and vWbp antibodies. Thus, Coa and vWbp neutralizing antibodies generate immune protection against staphylococcal disease.

Earlier studies revealed a requirement of coagulase for resisting phagocytosis in blood (Smith et al., 1947) and the inventors observed a similar phenotype for  $\Delta$ coa mutants in lepirudin-treated mouse blood (see Example 3 below). As vWbp displays higher affinity for human prothrombin than the mouse counterpart, it is suspected the same may be true for  $\Delta$ vWbp variants in human blood. Further, expression of Coa and vWbp in abscess lesions as well as their striking distribution in the eosinophilic pseudocapsule surrounding (staphylococcal abscess communities (SACs) or the peripheral fibrin wall, suggest that secreted coagulases contribute to the establishment of these lesions. This hypothesis was tested and, indeed,  $\Delta$ coa mutants were defective in the establishment of abscesses. A corresponding test, blocking Coa function with specific antibodies, produced the same effect. Consequently, it is proposed that the clotting of fibrin is a critical event in the establishment of staphylococcal abscesses that can be targeted for the development of protective vaccines. Due to their overlapping function on human prothrombin, both Coa and vWbp are considered excellent candidates for vaccine development.

#### C. Other Staphylococcal Antigens

Research over the past several decades identified *S. aureus* exotoxins, surface proteins and regulatory molecules as important virulence factors (Foster, 2005; Mazmanian et al., 2001; Novick, 2003). Much progress has been achieved regarding the regulation of these genes. For example, staphylococci perform a bacterial census via the secretion of auto-inducing peptides that bind to a cognate receptor at threshold concentration, thereby activating phospho-relay reactions and transcriptional activation of many of the exotoxin genes (Novick, 2003). The pathogenesis of staphylococcal infec-

tions relies on these virulence factors (secreted exotoxins, exopolysaccharides, and surface adhesins). The development of staphylococcal vaccines is hindered by the multifaceted nature of staphylococcal invasion mechanisms. It is well established that live attenuated microorganisms are highly effective vaccines; immune responses elicited by such vaccines are often of greater magnitude and of longer duration than those produced by non-replicating immunogens. One explanation for this may be that live attenuated strains establish limited infections in the host and mimic the early stages of natural infection. Embodiments of the invention are directed to compositions and methods including variant SpA polypeptides and peptides, as well as other immunogenic extracellular proteins, polypeptides, and peptides (including both secreted and cell surface proteins or peptides) of gram positive bacteria for the use in mitigating or immunizing against infection. In particular embodiments the bacteria is a *staphylococcus* bacteria. Extracellular proteins, polypeptides, or peptides include, but are not limited to secreted and cell surface proteins of the targeted bacteria.

The human pathogen *S. aureus* secretes EsxA and EsxB, two ESAT-6 like proteins, across the bacterial envelope (Burts et al., 2005, which is incorporated herein by reference). Staphylococcal esxA and esxB are clustered with six other genes in the order of transcription: esxA esaA essA esaB essB essC esaC esxB. The acronyms esa, ess, and esx stand for ESAT-6 secretion accessory, system, and extracellular, respectively, depending whether the encoded proteins play an accessory (esa) or direct (ess) role for secretion, or are secreted (esx) in the extracellular milieu. The entire cluster of eight genes is herein referred to as the Ess cluster. EsxA, esxB, essA, essB, and essC are all required for synthesis or secretion of EsxA and EsxB. Mutants that fail to produce EsxA, EsxB, and EssC display defects in the pathogenesis of *S. aureus* murine abscesses, suggesting that this specialized secretion system may be a general strategy of human bacterial pathogenesis. Secretion of non-WXG100 substrates by the ESX-1 pathway has been reported for several antigens including EspA, EspB, Rv3483c, and Rv3615c (Fortune et al., 2005; MacGurn et al., 2005; McLaughlin et al., 2007; Xu et al., 2007). The alternate ESX-5 pathway has also been shown to secrete both WXG100 and non-WXG100 proteins in pathogenic mycobacteria (Abdallah et al., 2007; Abdallah et al., 2006).

The *Staphylococcus aureus* Ess pathway can be viewed as a secretion module equipped with specialized transport components (Ess), accessory factors (Esa) and cognate secretion substrates (Esx). EssA, EssB and EssC are required for EsxA and EsxB secretion. Because EssA, EssB and EssC are predicted to be transmembrane proteins, it is contemplated that these proteins form a secretion apparatus. Some of the proteins in the ess gene cluster may actively transport secreted substrates (acting as motor) while others may regulate transport (regulator). Regulation may be achieved, but need not be limited to, transcriptional or post-translational mechanisms for secreted polypeptides, sorting of specific substrates to defined locations (e.g., extracellular medium or host cells), or timing of secretion events during infection. At this point, it is unclear whether all secreted Esx proteins function as toxins or contribute indirectly to pathogenesis.

Staphylococci rely on surface protein mediated-adhesion to host cells or invasion of tissues as a strategy for escape from immune defenses. Furthermore, *S. aureus* utilize surface proteins to sequester iron from the host during infection. The majority of surface proteins involved in staphylococcal pathogenesis carry C-terminal sorting signals, i.e., they are covalently linked to the cell wall envelope by sortase. Further, staphylococcal strains lacking the genes required for surface

protein anchoring, i.e., sortase A and B, display a dramatic defect in the virulence in several different mouse models of disease. Thus, surface protein antigens represent a validated vaccine target as the corresponding genes are essential for the development of staphylococcal disease and can be exploited in various embodiments of the invention. The sortase enzyme superfamily are Gram-positive transpeptidases responsible for anchoring surface protein virulence factors to the peptidoglycan cell wall layer. Two sortase isoforms have been identified in *Staphylococcus aureus*, SrtA and SrtB. These enzymes have been shown to recognize a LPXTG motif in substrate proteins. The SrtB isoform appears to be important in heme iron acquisition and iron homeostasis, whereas the SrtA isoform plays a critical role in the pathogenesis of Gram-positive bacteria by modulating the ability of the bacterium to adhere to host tissue via the covalent anchoring of adhesins and other proteins to the cell wall peptidoglycan. In certain embodiments the SpA variants described herein can be used in combination with other staphylococcal proteins such as Coa, Eap, Ehb, Emp, EsaC, EsaB, EsxA, EsxB, Hla, SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, IsdC, SasF, vWbp, and/or vWh proteins.

Certain aspects of the invention include methods and compositions concerning proteinaceous compositions including polypeptides, peptides, or nucleic acid encoding SpA variant(s) and other staphylococcal antigens such as other proteins transported by the Ess pathway, or sortase substrates. These proteins may be modified by deletion, insertion, and/or substitution.

The Esx polypeptides include the amino acid sequence of Esx proteins from bacteria in the *Staphylococcus* genus. The Esx sequence may be from a particular *staphylococcus* species, such as *Staphylococcus aureus*, and may be from a particular strain, such as Newman. In certain embodiments, the EsxA sequence is SAV0282 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number Q99WU4 (gi|68565539), which is hereby incorporated by reference. In other embodiments, the EsxB sequence is SAV0290 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number Q99WT7 (gi|68565532), which is hereby incorporated by reference. In further embodiments, other polypeptides transported by the Ess pathway may be used, the sequences of which may be identified by one of skill in the art using databases and internet accessible resources.

The sortase substrate polypeptides include, but are not limited to the amino acid sequence of SdrC, SdrD, SdrE, IsdA, IsdB, ClfA, ClfB, IsdC or SasF proteins from bacteria in the *Staphylococcus* genus. The sortase substrate polypeptide sequence may be from a particular *staphylococcus* species, such as *Staphylococcus aureus*, and may be from a particular strain, such as Newman. In certain embodiments, the SdrD sequence is from strain N315 and can be accessed using Genbank Accession Number NP\_373773.1 (gi|15926240), which is incorporated by reference. In other embodiments, the SdrE sequence is from strain N315 and can be accessed using Genbank Accession Number NP\_373774.1 (gi|15926241), which is incorporated by reference. In other embodiments, the IsdA sequence is SAV1130 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number NP\_371654.1 (gi|15924120), which is incorporated by reference. In other embodiments, the IsdB sequence is SAV1129 from strain Mu50 (which is the same amino acid sequence for Newman) and can be accessed using Genbank Accession Number NP\_371653.1 (gi|15924119), which is

incorporated by reference. In further embodiments, other polypeptides transported by the Ess pathway or processed by sortase may be used, the sequences of which may be identified by one of skill in the art using databases and internet accessible resources.

Examples of various proteins that can be used in the context of the present invention can be identified by analysis of database submissions of bacterial genomes, including but not limited to accession numbers NC\_002951 (GI:57650036 and GenBank CP000046), NC\_002758 (GI:57634611 and GenBank BA000017), NC\_002745 (GI:29165615 and GenBank BA000018), NC\_003923 (GI:21281729 and GenBank BA000033), NC\_002952 (GI:49482253 and GenBank BX571856), NC\_002953 (GI:49484912 and GenBank BX571857), NC\_007793 (GI:87125858 and GenBank CP000255), NC\_007795 (GI:87201381 and GenBank CP000253) each of which are incorporated by reference.

As used herein, a "protein" or "polypeptide" refers to a molecule comprising at least ten amino acid residues. In some embodiments, a wild-type version of a protein or polypeptide are employed, however, in many embodiments of the invention, a modified protein or polypeptide is employed to generate an immune response. The terms described above may be used interchangeably. A "modified protein" or "modified polypeptide" or a "variant" refers to a protein or polypeptide whose chemical structure, particularly its amino acid sequence, is altered with respect to the wild-type protein or polypeptide. In some embodiments, a modified/variant protein or polypeptide has at least one modified activity or function (recognizing that proteins or polypeptides may have multiple activities or functions). It is specifically contemplated that a modified/variant protein or polypeptide may be altered with respect to one activity or function yet retain a wild-type activity or function in other respects, such as immunogenicity.

In certain embodiments the size of a protein or polypeptide (wild-type or modified) may comprise, but is not limited to, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, 550, 575, 600, 625, 650, 675, 700, 725, 750, 775, 800, 825, 850, 875, 900, 925, 950, 975, 1000, 1100, 1200, 1300, 1400, 1500, 1750, 2000, 2250, 2500 amino molecules or greater, and any range derivable therein, or derivative of a corresponding amino sequence described or referenced herein. It is contemplated that polypeptides may be mutated by truncation, rendering them shorter than their corresponding wild-type form, but also they might be altered by fusing or conjugating a heterologous protein sequence with a particular function (e.g., for targeting or localization, for enhanced immunogenicity, for purification purposes, etc.).

As used herein, an "amino molecule" refers to any amino acid, amino acid derivative, or amino acid mimic known in the art. In certain embodiments, the residues of the proteinaceous molecule are sequential, without any non-amino molecule interrupting the sequence of amino molecule residues. In other embodiments, the sequence may comprise one or more non-amino molecule moieties. In particular embodiments, the sequence of residues of the proteinaceous molecule may be interrupted by one or more non-amino molecule moieties.

Accordingly, the term "proteinaceous composition" encompasses amino molecule sequences comprising at least

one of the 20 common amino acids in naturally synthesized proteins, or at least one modified or unusual amino acid.

Proteinaceous compositions may be made by any technique known to those of skill in the art, including (i) the expression of proteins, polypeptides, or peptides through standard molecular biological techniques, (ii) the isolation of proteinaceous compounds from natural sources, or (iii) the chemical synthesis of proteinaceous materials. The nucleotide as well as the protein, polypeptide, and peptide sequences for various genes have been previously disclosed, and may be found in the recognized computerized databases. One such database is the National Center for Biotechnology Information's Genbank and GenPept databases (on the World Wide Web at [ncbi.nlm.nih.gov/](http://ncbi.nlm.nih.gov/)). The coding regions for these genes may be amplified and/or expressed using the techniques disclosed herein or as would be known to those of ordinary skill in the art.

Amino acid sequence variants of SpA, coagulases and other polypeptides of the invention can be substitutional, insertional, or deletion variants. A variation in a polypeptide of the invention may affect 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, or more non-contiguous or contiguous amino acids of the polypeptide, as compared to wild-type. A variant can comprise an amino acid sequence that is at least 50%, 60%, 70%, 80%, or 90%, including all values and ranges there between, identical to any sequence provided or referenced herein, e.g., SEQ ID NO:2-8 or SEQ ID NO:11-30. A variant can include 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more substitute amino acids. A polypeptide processed or secreted by the Ess pathway or other surface proteins (see Table 1) or sortase substrates from any *staphylococcus* species and strain are contemplated for use in compositions and methods described herein.

Deletion variants typically lack one or more residues of the native or wild-type protein. Individual residues can be deleted or a number of contiguous amino acids can be deleted. A stop codon may be introduced (by substitution or insertion) into an encoding nucleic acid sequence to generate a truncated protein. Insertional mutants typically involve the addition of material at a non-terminal point in the polypeptide. This may include the insertion of one or more residues. Terminal additions, called fusion proteins, may also be generated. These fusion proteins include multimers or concatamers of one or more peptide or polypeptide described or referenced herein.

Substitutional variants typically contain the exchange of one amino acid for another at one or more sites within the protein, and may be designed to modulate one or more properties of the polypeptide, with or without the loss of other functions or properties. Substitutions may be conservative, that is, one amino acid is replaced with one of similar shape and charge. Conservative substitutions are well known in the art and include, for example, the changes of: alanine to serine; arginine to lysine; asparagine to glutamine or histidine; aspartate to glutamate; cysteine to serine; glutamine to asparagine; glutamate to aspartate; glycine to proline; histidine to asparagine or glutamine; isoleucine to leucine or valine; leucine to valine or isoleucine; lysine to arginine; methionine to leucine or isoleucine; phenylalanine to tyrosine, leucine or methionine; serine to threonine; threonine to serine; tryptophan to tyrosine; tyrosine to tryptophan or phenylalanine; and valine to isoleucine or leucine. Alternatively, substitutions may be non-conservative such that a function or activity of the polypeptide is affected. Non-conservative changes typically involve substituting a residue with one that is chemically dissimilar, such as a polar or charged amino acid for a non-polar or uncharged amino acid, and vice versa.

TABLE 2

Exemplary surface proteins of <i>S. aureus</i> strains.								
SAV #	SA#	Surface	MW2	Mu50	N315	Newman	MRSA252*	MSSA476*
SAV0111	SA0107	Spa	492	450	450	520	516	492
SAV2503	SA2291	FnBPA	1015	1038	1038	741	—	1015
SAV2502	SA2290	FnBPB	943	961	961	677	965	957
SAV0811	SA0742	CIfA	946	935	989	933	1029	928
SAV2630	SA2423	CIfB	907	877	877	913	873	905
Np	Np	Np	1183	—	—	—	1183	1183
SAV0561	SA0519	SdrC	955	953	953	947	906	957
SAV0562	SA0520	SdrD	1347	1385	1385	1315	—	1365
SAV0563	SA0521	SdrE	1141	1141	1141	1166	1137	1141
Np	Np	Pls	—	—	—	—	—	—
SAV2654	SA2447	SasA	2275	2271	2271	2271	1351	2275
SAV2160	SA1964	SasB	686	2481	2481	2481	2222	685
	SA1577	SasC	2186	213	2186	2186	2189	2186
SAV0134	SA0129	SasD	241	241	241	241	221	241
SAV1130	SA0977	SasE/IsdA	350	350	350	350	354	350
SAV2646	SA2439	SasF	635	635	635	635	627	635
SAV2496		SasG	1371	525	927	—	—	1371
SAV0023	SA0022	SasH	772	—	772	772	786	786
SAV1731	SA1552	SasI	895	891	891	891	534	895
SAV1129	SA0976	SasJ/IsdB	645	645	645	645	652	645
	SA2381	SasK	198	211	211	—	—	197
	Np	SasL	—	232	—	—	—	—
SAV1131	SA0978	IsdC	227	227	227	227	227	227

Proteins of the invention may be recombinant, or synthesized in vitro. Alternatively, a non-recombinant or recombinant protein may be isolated from bacteria. It is also contemplated that a bacteria containing such a variant may be implemented in compositions and methods of the invention. Consequently, a protein need not be isolated.

The term "functionally equivalent codon" is used herein to refer to codons that encode the same amino acid, such as the six codons for arginine or serine, and also refers to codons that encode biologically equivalent amino acids (see Table 2, below).

TABLE 3

Codon Table			
Amino Acids			Codons
Alanine	Ala	A	GCA GCC GCG GCU
Cysteine	Cys	C	UGC UGU
Aspartic acid	Asp	D	GAC GAU
Glutamic acid	Glu	E	GAA GAG
Phenylalanine	Phe	F	UUC UUU
Glycine	Gly	G	GGA GGC GGG GGU
Histidine	His	H	CAC CAU
Isoleucine	Ile	I	AUA AUC AUU
Lysine	Lys	K	AAA AAG
Leucine	Leu	L	UUA UUG CUA CUC CUG CUU
Methionine	Met	M	AUG
Asparagine	Asn	N	AAC AAU
Proline	Pro	P	CCA CCC CCG CCU
Glutamine	Gln	Q	CAA CAG
Arginine	Arg	R	AGA AGG CGA CGC CGG CGU

TABLE 3-continued

Codon Table							
	Amino Acids						Codons
	Serine	Ser	S	AGC	AGU	UCA	UCC UCG UCU
	Threonine	Thr	T	ACA	ACC	ACG	ACU
	Valine	Val	V	GUA	GUC	GUG	GUU
	Tryptophan	Trp	W	UGG			
	Tyrosine	Tyr	Y	UAC	UAU		

It also will be understood that amino acid and nucleic acid sequences may include additional residues, such as additional N- or C-terminal amino acids, or 5' or 3' sequences, respectively, and yet still be essentially as set forth in one of the sequences disclosed herein, so long as the sequence meets the criteria set forth above, including the maintenance of biological protein activity (e.g., immunogenicity) where protein expression is concerned. The addition of terminal sequences particularly applies to nucleic acid sequences that may, for example, include various non-coding sequences flanking either of the 5' or 3' portions of the coding region.

The following is a discussion based upon changing of the amino acids of a protein to create a variant polypeptide or peptide. For example, certain amino acids may be substituted for other amino acids in a protein structure with or without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of antibodies or binding sites on substrate molecules. Since it is the interactive capacity and nature of a protein that defines that protein's functional activity, certain amino acid substitutions can be made in a protein sequence, and in its underlying DNA coding sequence, and nevertheless produce a protein with a desirable property. It is thus contemplated by the inventors that various changes may be made in the DNA sequences of genes.

It is contemplated that in compositions of the invention, there is between about 0.001 mg and about 10 mg of total

polypeptide, peptide, and/or protein per ml. The concentration of protein in a composition can be about, at least about or at most about 0.001, 0.010, 0.050, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, 8.5, 9.0, 9.5, 10.0 mg/ml or more (or any range derivable therein). Of this, about, at least about, or at most about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100% may be an SpA variant or a coagulase, and may be used in combination with other peptides or polypeptides, such as other bacterial peptides and/or antigens.

The present invention contemplates the administration of variant SpA polypeptides or peptides to effect a preventative therapy or therapeutic effect against the development of a disease or condition associated with infection by a *staphylococcus* pathogen.

In certain aspects, combinations of staphylococcal antigens are used in the production of an immunogenic composition that is effective at treating or preventing staphylococcal infection. Staphylococcal infections progress through several different stages. For example, the staphylococcal life cycle involves commensal colonization, initiation of infection by accessing adjoining tissues or the bloodstream, and/or anaerobic multiplication in the blood. The interplay between *S. aureus* virulence determinants and the host defense mechanisms can induce complications such as endocarditis, metastatic abscess formation, and sepsis syndrome. Different molecules on the surface of the bacterium are involved in different steps of the infection cycle. Combinations of certain antigens can elicit an immune response which protects against multiple stages of staphylococcal infection. The effectiveness of the immune response can be measured either in animal model assays and/or using an opsonophagocytic assay.

#### D. Polypeptides and Polypeptide Production

The present invention describes polypeptides, peptides, and proteins and immunogenic fragments thereof for use in various embodiments of the present invention. For example, specific polypeptides are assayed for or used to elicit an immune response. In specific embodiments, all or part of the proteins of the invention can also be synthesized in solution or on a solid support in accordance with conventional techniques. Various automatic synthesizers are commercially available and can be used in accordance with known protocols. See, for example, Stewart and Young, (1984); Tam et al., (1983); Merrifield, (1986); and Barany and Merrifield (1979), each incorporated herein by reference.

Alternatively, recombinant DNA technology may be employed wherein a nucleotide sequence which encodes a peptide of the invention is inserted into an expression vector, transformed or transfected into an appropriate host cell and cultivated under conditions suitable for expression.

One embodiment of the invention includes the use of gene transfer to cells, including microorganisms, for the production and/or presentation of polypeptides or peptides. The gene for the polypeptide or peptide of interest may be transferred into appropriate host cells followed by culture of cells under the appropriate conditions. The generation of recombinant expression vectors, and the elements included therein, are well known in the art and briefly discussed herein. Alternatively, the protein to be produced may be an endogenous protein normally synthesized by the cell that is isolated and purified.

Another embodiment of the present invention uses autologous B lymphocyte cell lines, which are transfected with a viral vector that expresses an immunogenic product, and more specifically, a protein having immunogenic activity. Other examples of mammalian host cell lines include, but are not limited to Vero and HeLa cells, other B- and T-cell lines, such as CEM, 721.221, H9, Jurkat, Raji, as well as cell lines of Chinese hamster ovary, W138, BHK, COS-7, 293, HepG2, 3T3, RIN and MDCK cells. In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or that modifies and processes the gene product in the manner desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed.

A number of selection systems may be used including, but not limited to HSV thymidine kinase, hypoxanthine-guanine phosphoribosyltransferase, and adenine phosphoribosyltransferase genes, in tk-, hgprrt- or aprt- cells, respectively. Also, anti-metabolite resistance can be used as the basis of selection: for dhfr, which confers resistance to trimethoprim and methotrexate; gpt, which confers resistance to mycophenolic acid; neo, which confers resistance to the aminoglycoside G418; and hygromycin, which confers resistance to hygromycin.

Animal cells can be propagated in vitro in two modes: as non-anchorage-dependent cells growing in suspension throughout the bulk of the culture or as anchorage-dependent cells requiring attachment to a solid substrate for their propagation (i.e., a monolayer type of cell growth).

Non-anchorage dependent or suspension cultures from continuous established cell lines are the most widely used means of large scale production of cells and cell products. However, suspension cultured cells have limitations, such as tumorigenic potential and lower protein production than adherent cells.

Where a protein is specifically mentioned herein, it is preferably a reference to a native or recombinant protein or optionally a protein in which any signal sequence has been removed. The protein may be isolated directly from the staphylococcal strain or produced by recombinant DNA techniques. Immunogenic fragments of the protein may be incorporated into the immunogenic composition of the invention. These are fragments comprising at least 10 amino acids, 20 amino acids, 30 amino acids, 40 amino acids, 50 amino acids, or 100 amino acids, including all values and ranges there between, taken contiguously from the amino acid sequence of the protein. In addition, such immunogenic fragments are immunologically reactive with antibodies generated against the Staphylococcal proteins or with antibodies generated by infection of a mammalian host with Staphylococci. Immunogenic fragments also include fragments that when administered at an effective dose, (either alone or as a hapten bound to a carrier), elicit a protective or therapeutic immune response against Staphylococcal infection, in certain aspects it is protective against *S. aureus* and/or *S. epidermidis* infection. Such an immunogenic fragment may include, for example, the protein lacking an N-terminal leader sequence, and/or a transmembrane domain and/or a C-terminal anchor domain. In a preferred aspect the immunogenic fragment according to the invention comprises substantially all of the extracellular domain of a protein which has at least 80% identity, at least 85% identity, at least 90% identity, at least 95% identity, or at

least 97-99% identity, including all values and ranges there between, to a sequence selected segment of a polypeptide described or referenced herein.

Also included in immunogenic compositions of the invention are fusion proteins composed of one or more Staphylococcal proteins, or immunogenic fragments of staphylococcal proteins. Such fusion proteins may be made recombinantly and may comprise one portion of at least 1, 2, 3, 4, 5, or 6 staphylococcal proteins or segments. Alternatively, a fusion protein may comprise multiple portions of at least 1, 2, 3, 4 or 5 staphylococcal proteins. These may combine different Staphylococcal proteins and/or multiples of the same protein or protein fragment, or immunogenic fragments in the same protein (forming a multimer or a concatamer). Alternatively, the invention also includes individual fusion proteins of Staphylococcal proteins or immunogenic fragments thereof, as a fusion protein with heterologous sequences such as a provider of T-cell epitopes or purification tags, for example:  $\beta$ -galactosidase, glutathione-S-transferase, green fluorescent proteins (GFP), epitope tags such as FLAG, myc tag, poly histidine, or viral surface proteins such as influenza virus haemagglutinin, or bacterial proteins such as tetanus toxoid, diphtheria toxoid, or CRM197.

## II. NUCLEIC ACIDS

In certain embodiments, the present invention concerns recombinant polynucleotides encoding the proteins, polypeptides, peptides of the invention. The nucleic acid sequences for SpA, coagulases and other bacterial proteins are included, all of which are incorporated by reference, and can be used to prepare peptides or polypeptides.

As used in this application, the term "polynucleotide" refers to a nucleic acid molecule that either is recombinant or has been isolated free of total genomic nucleic acid. Included within the term "polynucleotide" are oligonucleotides (nucleic acids of 100 residues or less in length), recombinant vectors, including, for example, plasmids, cosmids, phage, viruses, and the like. Polynucleotides include, in certain aspects, regulatory sequences, isolated substantially away from their naturally occurring genes or protein encoding sequences. Polynucleotides may be single-stranded (coding or antisense) or double-stranded, and may be RNA, DNA (genomic, cDNA or synthetic), analogs thereof, or a combination thereof. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide.

In this respect, the term "gene," "polynucleotide," or "nucleic acid" is used to refer to a nucleic acid that encodes a protein, polypeptide, or peptide (including any sequences required for proper transcription, post-translational modification, or localization). As will be understood by those in the art, this term encompasses genomic sequences, expression cassettes, cDNA sequences, and smaller engineered nucleic acid segments that express, or may be adapted to express, proteins, polypeptides, domains, peptides, fusion proteins, and mutants. A nucleic acid encoding all or part of a polypeptide may contain a contiguous nucleic acid sequence of: 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 441, 450, 460, 470, 480, 490, 500, 510, 520, 530, 540, 550, 560, 570, 580, 590, 600, 610, 620, 630, 640, 650, 660, 670, 680, 690, 700, 710, 720, 730, 740, 750, 760, 770, 780, 790, 800, 810, 820, 830, 840, 850, 860, 870, 880, 890, 900, 910, 920, 930, 940, 950, 960, 970, 980, 990, 1000, 1010, 1020, 1030, 1040, 1050, 1060, 1070, 1080, 1090, 1095, 1100, 1500, 2000, 2500, 3000, 3500, 4000, 4500,

5000, 5500, 6000, 6500, 7000, 7500, 8000, 9000, 10000, or more nucleotides, nucleosides, or base pairs, including all values and ranges therebetween, of a polynucleotide encoding one or more amino acid sequence described or referenced herein. It also is contemplated that a particular polypeptide may be encoded by nucleic acids containing variations having slightly different nucleic acid sequences but, nonetheless, encode the same or substantially similar protein (see Table 3 above).

In particular embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors incorporating nucleic acid sequences that encode a variant SpA or coagulase. The term "recombinant" may be used in conjunction with a polynucleotide or polypeptide and generally refers to a polypeptide or polynucleotide produced and/or manipulated in vitro or that is a replication product of such a molecule.

In other embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors incorporating nucleic acid sequences that encode a variant SpA or coagulase polypeptide or peptide to generate an immune response in a subject. In various embodiments the nucleic acids of the invention may be used in genetic vaccines.

The nucleic acid segments used in the present invention can be combined with other nucleic acid sequences, such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant nucleic acid protocol. In some cases, a nucleic acid sequence may encode a polypeptide sequence with additional heterologous coding sequences, for example to allow for purification of the polypeptide, transport, secretion, post-translational modification, or for therapeutic benefits such as targeting or efficacy. As discussed above, a tag or other heterologous polypeptide may be added to the modified polypeptide-encoding sequence, wherein "heterologous" refers to a polypeptide that is not the same as the modified polypeptide.

In certain other embodiments, the invention concerns isolated nucleic acid segments and recombinant vectors that include within their sequence a contiguous nucleic acid sequence from SEQ ID NO:1 (SpA domain D) or SEQ ID NO:3 (SpA) or any other nucleic acid sequences encoding coagulases or other secreted virulence factors and/or surface proteins including proteins transported by the Ess pathway, processed by sortase, or proteins incorporated herein by reference.

In certain embodiments, the present invention provides polynucleotide variants having substantial identity to the sequences disclosed herein; those comprising at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% or higher sequence identity, including all values and ranges there between, compared to a polynucleotide sequence of this invention using the methods described herein (e.g., BLAST analysis using standard parameters).

The invention also contemplates the use of polynucleotides which are complementary to all the above described polynucleotides.

### A. Vectors

Polypeptides of the invention may be encoded by a nucleic acid molecule comprised in a vector. The term "vector" is used to refer to a carrier nucleic acid molecule into which a heterologous nucleic acid sequence can be inserted for introduction into a cell where it can be replicated and expressed. A nucleic acid sequence can be "heterologous," which means

that it is in a context foreign to the cell in which the vector is being introduced or to the nucleic acid in which is incorporated, which includes a sequence homologous to a sequence in the cell or nucleic acid but in a position within the host cell or nucleic acid where it is ordinarily not found. Vectors include DNAs, RNAs, plasmids, cosmids, viruses (bacteriophage, animal viruses, and plant viruses), and artificial chromosomes (e.g., YACs). One of skill in the art would be well equipped to construct a vector through standard recombinant techniques (for example Sambrook et al., 2001; Ausubel et al., 1996, both incorporated herein by reference). In addition to encoding a variant SpA polypeptide the vector can encode other polypeptide sequences such as a one or more other bacterial peptide, a tag, or an immunogenicity enhancing peptide. Useful vectors encoding such fusion proteins include pIN vectors (Inouye et al., 1985), vectors encoding a stretch of histidines, and pGEX vectors, for use in generating glutathione S-transferase (GST) soluble fusion proteins for later purification and separation or cleavage.

The term "expression vector" refers to a vector containing a nucleic acid sequence coding for at least part of a gene product capable of being transcribed. In some cases, RNA molecules are then translated into a protein, polypeptide, or peptide. Expression vectors can contain a variety of "control sequences," which refer to nucleic acid sequences necessary for the transcription and possibly translation of an operably linked coding sequence in a particular host organism. In addition to control sequences that govern transcription and translation, vectors and expression vectors may contain nucleic acid sequences that serve other functions as well and are described herein.

#### 1. Promoters and Enhancers

A "promoter" is a control sequence. The promoter is typically a region of a nucleic acid sequence at which initiation and rate of transcription are controlled. It may contain genetic elements at which regulatory proteins and molecules may bind such as RNA polymerase and other transcription factors. The phrases "operatively positioned," "operatively linked," "under control," and "under transcriptional control" mean that a promoter is in a correct functional location and/or orientation in relation to a nucleic acid sequence to control transcriptional initiation and expression of that sequence. A promoter may or may not be used in conjunction with an "enhancer," which refers to a cis-acting regulatory sequence involved in the transcriptional activation of a nucleic acid sequence.

Naturally, it may be important to employ a promoter and/or enhancer that effectively directs the expression of the DNA segment in the cell type or organism chosen for expression. Those of skill in the art of molecular biology generally know the use of promoters, enhancers, and cell type combinations for protein expression (see Sambrook et al., 2001, incorporated herein by reference). The promoters employed may be constitutive, tissue-specific, or inducible and in certain embodiments may direct high level expression of the introduced DNA segment under specified conditions, such as large-scale production of recombinant proteins or peptides.

Various elements/promoters may be employed in the context of the present invention to regulate the expression of a gene. Examples of such inducible elements, which are regions of a nucleic acid sequence that can be activated in response to a specific stimulus, include but are not limited to Immunoglobulin Heavy Chain (Banerji et al., 1983; Gilles et al., 1983; Grosschedl et al., 1985; Atchinson et al., 1986, 1987; Imler et al., 1987; Weinberger et al., 1984; Kiledjian et al., 1988; Porton et al., 1990), Immunoglobulin Light Chain (Queen et al., 1983; Picard et al., 1984), T Cell Receptor

(Luria et al., 1987; Winoto et al., 1989; Redondo et al., 1990), HLA DQ  $\alpha$  and/or DQ  $\beta$  (Sullivan et al., 1987),  $\beta$  Interferon (Goodbourn et al., 1986; Fujita et al., 1987; Goodbourn et al., 1988), Interleukin-2 (Greene et al., 1989), Interleukin-2 Receptor (Greene et al., 1989; Lin et al., 1990), MHC Class II 5 (Koch et al., 1989), MHC Class II HLA-DR $\alpha$  (Sherman et al., 1989),  $\beta$ -Actin (Kawamoto et al., 1988; Ng et al., 1989), Muscle Creatine Kinase (MCK) (Jaynes et al., 1988; Horlick et al., 1989; Johnson et al., 1989), Prealbumin (Transthyretin) (Costa et al., 1988), Elastase I (Ornitz et al., 1987), Metallothionein (MTII) (Karin et al., 1987; Culotta et al., 1989), Collagenase (Pinkert et al., 1987; Angel et al., 1987), Albumin (Pinkert et al., 1987; Tronche et al., 1989, 1990),  $\alpha$ -Fetoprotein (Godbout et al., 1988; Campere et al., 1989),  $\gamma$ -Globin (Bodine et al., 1987; Perez-Stable et al., 1990),  $\beta$ -Globin (Trudel et al., 1987), c-fos (Cohen et al., 1987), c-Ha-Ras (Triesman, 1986; Deschamps et al., 1985), Insulin (Edlund et al., 1985), Neural Cell Adhesion Molecule (NCAM) (Hirsh et al., 1990),  $\alpha$ 1-Antitrypsin (Latimer et al., 1990), H2B (TH2B) Histone (Hwang et al., 1990), Mouse and/or Type I Collagen (Ripe et al., 1989), Glucose-Regulated Proteins (GRP94 and GRP78) (Chang et al., 1989), Rat Growth Hormone (Larsen et al., 1986), Human Serum Amyloid A (SAA) (Edbrooke et al., 1989), Troponin I (TN I) (Yutzey et al., 1989), Platelet-Derived Growth Factor (PDGF) (Pech et al., 1989), Duchenne Muscular Dystrophy (Klamut et al., 1990), SV40 (Banerji et al., 1981; Moreau et al., 1981; Sleigh et al., 1985; Firak et al., 1986; Herr et al., 1986; Imbra et al., 1986; Kadesch et al., 1986; Wang et al., 1986; Ondek et al., 1987; Kuhl et al., 1987; Schaffner et al., 1988), Polyoma (Swartzendruber et al., 1975; Vasseur et al., 1980; Katinka et al., 1980, 1981; Tyndell et al., 1981; Dandolo et al., 1983; de Villiers et al., 1984; Hen et al., 1986; Satake et al., 1988; Campbell et al., 1988), Retroviruses (Kriegler et al., 1982, 1983; Levinson et al., 1982; Kriegler et al., 1983, 1984a, b, 1988; Bosze et al., 1986; Miksicek et al., 1986; Celander et al., 1987; Thiesen et al., 1988; Celander et al., 1988; Choi et al., 1988; Reisman et al., 1989), Papilloma Virus (Campo et al., 1983; Lusky et al., 1983; Spandidos and Wilkie, 1983; Spalholz et al., 1985; Lusky et al., 1986; Cripe et al., 1987; Gloss et al., 1987; Hirochika et al., 1987; Stephens et al., 1987), Hepatitis B Virus (Bulla et al., 1986; Jameel et al., 1986; Shaul et al., 1987; Spandau et al., 1988; Vannice et al., 1988), Human Immunodeficiency Virus (Muesing et al., 1987; Hauber et al., 1988; Jakobovits et al., 1988; Feng et al., 1988; Takebe et al., 1988; Rosen et al., 1988; Berkhout et al., 1989; Laspia et al., 1989; Sharp et al., 1989; Braddock et al., 1989), Cytomegalovirus (CMV) IE (Weber et al., 1984; Boshart et al., 1985; Foecking et al., 1986), Gibbon Ape Leukemia Virus (Holbrook et al., 1987; Quinn et al., 1989).

Inducible elements include, but are not limited to MT II—Phorbol Ester (TFA)/Heavy metals (Palmiter et al., 1982; Haslinger et al., 1985; Searle et al., 1985; Stuart et al., 1985; Imagawa et al., 1987; Karin et al., 1987; Angel et al., 1987b; McNeill et al., 1989); MMTV (mouse mammary tumor virus)—Glucocorticoids (Huang et al., 1981; Lee et al., 1981; Majors et al., 1983; Chandler et al., 1983; Lee et al., 1984; Ponta et al., 1985; Sakai et al., 1988);  $\beta$ -Interferon—poly(rI) x/poly(rc) (Tavernier et al., 1983); Adenovirus 5 E2-E1A (Imperiale et al., 1984); Collagenase—Phorbol Ester (TPA) (Angel et al., 1987a); Stromelysin—Phorbol Ester (TPA) (Angel et al., 1987b); SV40—Phorbol Ester (TPA) (Angel et al., 1987b); Murine MX Gene—Interferon, Newcastle Disease Virus (Hug et al., 1988); GRP78 Gene—A23187 (Resendez et al., 1988);  $\alpha$ -2-Macroglobulin—IL-6 (Kunz et al., 1989); Vimentin—Serum (Rittling et al., 1989); MHC Class I Gene H-2kb—Interferon (Blonar et al., 1989); HSP70—E1A/SV40

Large T Antigen (Taylor et al., 1989, 1990a, 1990b); Proliferin—Phorbol Ester/TPA (Mordacq et al., 1989); Tumor Necrosis Factor—PMA (Hensel et al., 1989); and Thyroid Stimulating Hormone a Gene—Thyroid Hormone (Chatterjee et al., 1989).

The particular promoter that is employed to control the expression of peptide or protein encoding polynucleotide of the invention is not believed to be critical, so long as it is capable of expressing the polynucleotide in a targeted cell, preferably a bacterial cell. Where a human cell is targeted, it is preferable to position the polynucleotide coding region adjacent to and under the control of a promoter that is capable of being expressed in a human cell. Generally speaking, such a promoter might include either a bacterial, human or viral promoter.

In embodiments in which a vector is administered to a subject for expression of the protein, it is contemplated that a desirable promoter for use with the vector is one that is not down-regulated by cytokines or one that is strong enough that even if down-regulated, it produces an effective amount of a variant SpA for eliciting an immune response. Non-limiting examples of these are CMV IE and RSV LTR. Tissue specific promoters can be used, particularly if expression is in cells in which expression of an antigen is desirable, such as dendritic cells or macrophages. The mammalian MHC I and MHC II promoters are examples of such tissue-specific promoters.

#### 2. Initiation Signals and Internal Ribosome Binding Sites (IRES)

A specific initiation signal also may be required for efficient translation of coding sequences. These signals include the ATG initiation codon or adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals.

In certain embodiments of the invention, the use of internal ribosome entry sites (IRES) elements are used to create multigene, or polycistronic, messages. IRES elements are able to bypass the ribosome scanning model of 5' methylated Cap dependent translation and begin translation at internal sites (Pelletier and Sonenberg, 1988; Macejak and Sarnow, 1991). IRES elements can be linked to heterologous open reading frames. Multiple open reading frames can be transcribed together, each separated by an IRES, creating polycistronic messages. Multiple genes can be efficiently expressed using a single promoter/enhancer to transcribe a single message (see U.S. Pat. Nos. 5,925,565 and 5,935,819, herein incorporated by reference).

#### 3. Selectable and Screenable Markers

In certain embodiments of the invention, cells containing a nucleic acid construct of the present invention may be identified in vitro or in vivo by encoding a screenable or selectable marker in the expression vector. When transcribed and translated, a marker confers an identifiable change to the cell permitting easy identification of cells containing the expression vector. Generally, a selectable marker is one that confers a property that allows for selection. A positive selectable marker is one in which the presence of the marker allows for its selection, while a negative selectable marker is one in which its presence prevents its selection. An example of a positive selectable marker is a drug resistance marker.

#### B. Host Cells

As used herein, the terms “cell,” “cell line,” and “cell culture” may be used interchangeably. All of these terms also include their progeny, which is any and all subsequent generations. It is understood that all progeny may not be identical due to deliberate or inadvertent mutations. In the context of

expressing a heterologous nucleic acid sequence, “host cell” refers to a prokaryotic or eukaryotic cell, and it includes any transformable organism that is capable of replicating a vector or expressing a heterologous gene encoded by a vector. A host cell can, and has been, used as a recipient for vectors or viruses. A host cell may be “transfected” or “transformed,” which refers to a process by which exogenous nucleic acid, such as a recombinant protein-encoding sequence, is transferred or introduced into the host cell. A transformed cell includes the primary subject cell and its progeny.

Host cells may be derived from prokaryotes or eukaryotes, including bacteria, yeast cells, insect cells, and mammalian cells for replication of the vector or expression of part or all of the nucleic acid sequence(s). Numerous cell lines and cultures are available for use as a host cell, and they can be obtained through the American Type Culture Collection (ATCC), which is an organization that serves as an archive for living cultures and genetic materials ([www.atcc.org](http://www.atcc.org)).

#### C. Expression Systems

Numerous expression systems exist that comprise at least a part or all of the compositions discussed above. Prokaryote- and/or eukaryote-based systems can be employed for use with the present invention to produce nucleic acid sequences, or their cognate polypeptides, proteins and peptides. Many such systems are commercially and widely available.

The insect cell/baculovirus system can produce a high level of protein expression of a heterologous nucleic acid segment, such as described in U.S. Pat. Nos. 5,871,986, 4,879,236, both herein incorporated by reference, and which can be bought, for example, under the name MAXBAC® 2.0 from INVITROGEN® and BACPACK™ BACULOVIRUS EXPRESSION SYSTEM FROM CLONTECH®.

In addition to the disclosed expression systems of the invention, other examples of expression systems include STRATAGENE®'s COMPLETE CONTROL™ Inducible Mammalian Expression System, which involves a synthetic ecdysone-inducible receptor, or its pET Expression System, an *E. coli* expression system. Another example of an inducible expression system is available from INVITROGEN®, which carries the T-REX™ (tetracycline-regulated expression) System, an inducible mammalian expression system that uses the full-length CMV promoter. INVITROGEN® also provides a yeast expression system called the *Pichia methanolica* Expression System, which is designed for high-level production of recombinant proteins in the methylotrophic yeast *Pichia methanolica*. One of skill in the art would know how to express a vector, such as an expression construct, to produce a nucleic acid sequence or its cognate polypeptide, protein, or peptide.

### III. POLYSACCHARIDES

The immunogenic compositions of the invention may further comprise capsular polysaccharides including one or more of PIA (also known as PNAG) and/or *S. aureus* Type V and/or type VIII capsular polysaccharide and/or *S. epidermidis* Type I, and/or Type II and/or Type III capsular polysaccharide.

#### A. PIA (PNAG)

It is now clear that the various forms of staphylococcal surface polysaccharides identified as PS/A, PIA and SAA are the same chemical entity—PNAG (Maira-Litran et al., 2004). Therefore the term PIA or PNAG encompasses all these polysaccharides or oligosaccharides derived from them.

PIA is a polysaccharide intercellular adhesin and is composed of a polymer of  $\beta$ -(1 $\rightarrow$ 6)-linked glucosamine substituted with N-acetyl and O-succinyl constituents. This

polysaccharide is present in both *S. aureus* and *S. epidermidis* and can be isolated from either source (Joyce et al., 2003; Maira-Litran et al., 2002). For example, PNAG may be isolated from *S. aureus* strain MN8m (WO04/43407). PIA isolated from *S. epidermidis* is an integral constituent of biofilm. It is responsible for mediating cell-cell adhesion and probably also functions to shield the growing colony from the host's immune response. The polysaccharide previously known as poly-N-succinyl-β-(1→6)-glucosamine (PNSG) was recently shown not to have the expected structure since the identification of N-succinylation was incorrect (Maira-Litran et al., 2002). Therefore the polysaccharide formally known as PNSG and now found to be PNAG is also encompassed by the term PIA.

PIA (or PNAG) may be of different sizes varying from over 400 kDa to between 75 and 400 kDa to between 10 and 75 kDa to oligosaccharides composed of up to 30 repeat units (of β-(1→6)-linked glucosamine substituted with N-acetyl and O-succinyl constituents). Any size of PIA polysaccharide or oligosaccharide may be used in an immunogenic composition of the invention, in one aspect the polysaccharide is over 40 kDa. Sizing may be achieved by any method known in the art, for instance by microfluidization, ultrasonic irradiation or by chemical cleavage (WO 03/53462, EP497524, EP497525). In certain aspects PIA (PNAG) is at least or at most 40-400 kDa, 40-300 kDa, 50-350 kDa, 60-300 kDa, 50-250 kDa and 60-200 kDa.

PIA (PNAG) can have different degree of acetylation due to substitution on the amino groups by acetate. PIA produced in vitro is almost fully substituted on amino groups (95-100%). Alternatively, a deacetylated PIA (PNAG) can be used having less than 60%, 50%, 40%, 30%, 20%, 10% acetylation. Use of a deacetylated PIA (PNAG) is preferred since non-acetylated epitopes of PNAG are efficient at mediating opsonic killing of Gram positive bacteria, preferably *S. aureus* and/or *S. epidermidis*. In certain aspects, the PIA (PNAG) has a size between 40 kDa and 300 kDa and is deacetylated so that less than 60%, 50%, 40%, 30% or 20% of amino groups are acetylated.

The term deacetylated PNAG (dPNAG) refers to a PNAG polysaccharide or oligosaccharide in which less than 60%, 50%, 40%, 30%, 20% or 10% of the amino groups are acetylated. In certain aspects, PNAG is deacetylated to form dPNAG by chemically treating the native polysaccharide. For example, the native PNAG is treated with a basic solution such that the pH rises to above 10. For instance the PNAG is treated with 0.1-5 M, 0.2-4 M, 0.3-3 M, 0.5-2 M, 0.75-1.5 M or 1 M NaOH, KOH or NH<sub>4</sub>OH. Treatment is for at least 10 to 30 minutes, or 1, 2, 3, 4, 5, 10, 15 or 20 hours at a temperature of 20-100, 25-80, 30-60 or 30-50 or 35-45° C. dPNAG may be prepared as described in WO 04/43405.

The polysaccharide(s) can be conjugated or unconjugated to a carrier protein.

#### B. Type 5 and Type 8 Polysaccharides from *S. aureus*

Most strains of *S. aureus* that cause infection in man contain either Type 5 or Type 8 polysaccharides. Approximately 60% of human strains are Type 8 and approximately 30% are Type 5. The structures of Type 5 and Type 8 capsular polysaccharide antigens are described in Moreau et al., (1990) and Fournier et al., (1984). Both have FucNAcp in their repeat unit as well as ManNAcA which can be used to introduce a sulfhydryl group. The structures are:

##### Type 5

→4)-β-D-ManNAcA(3OAc)-(1→4)-α-L-FucNAc  
(1→3)-β-D-FucNAc-(1→

##### Type 8

→3)-β-D-ManNAcA(4OAc)-(1→3)-α-L-FucNAc  
(1→3)-β-D-FucNAc-(1→

Recently (Jones, 2005) NMR spectroscopy revised the structures to:

##### Type 5

→4)-β-D-ManNAcA-(1→4)-α-L-FucNAc(3OAc)-  
(1→3)-β-D-FucNAc-(1→

##### Type 8

→3)-β-D-ManNAcA(4OAc)-(1→3)-α-L-FucNAc  
(1→3)-α-D-FucNAc(1→

Polysaccharides may be extracted from the appropriate strain of *S. aureus* using method well known to of skill in the art, See U.S. Pat. No. 6,294,177. For example, ATCC 12902 is a Type 5 *S. aureus* strain and ATCC 12605 is a Type 8 *S. aureus* strain.

Polysaccharides are of native size or alternatively may be sized, for instance by microfluidisation, ultrasonic irradiation, or by chemical treatment. The invention also covers oligosaccharides derived from the type 5 and 8 polysaccharides from *S. aureus*. The type 5 and 8 polysaccharides included in the immunogenic composition of the invention are preferably conjugated to a carrier protein as described below or are alternatively unconjugated. The immunogenic compositions of the invention alternatively contains either type 5 or type 8 polysaccharide.

#### C. *S. aureus* 336 Antigen

In an embodiment, the immunogenic composition of the invention comprises the *S. aureus* 336 antigen described in U.S. Pat. No. 6,294,177. The 336 antigen comprises β-linked hexosamine, contains no O-acetyl groups, and specifically binds to antibodies to *S. aureus* Type 336 deposited under ATCC 55804. In an embodiment, the 336 antigen is a polysaccharide which is of native size or alternatively may be sized, for instance by microfluidisation, ultrasonic irradiation, or by chemical treatment. The invention also covers oligosaccharides derived from the 336 antigen. The 336 antigen can be unconjugated or conjugated to a carrier protein.

#### D. Type I, II and III Polysaccharides from *S. epidermidis*

Amongst the problems associated with the use of polysaccharides in vaccination, is the fact that polysaccharides per se are poor immunogens. It is preferred that the polysaccharides utilized in the invention are linked to a protein carrier which provide bystander T-cell help to improve immunogenicity. Examples of such carriers which may be conjugated to polysaccharide immunogens include the Diphtheria and Tetanus toxoids (DT, DT CRM197 and TT respectively), Keyhole Limpet Haemocyanin (KLH), and the purified protein derivative of Tuberculin (PPD), *Pseudomonas aeruginosa* exoprotein A (rEPA), protein D from *Haemophilus influenzae*, pneumolysin or fragments of any of the above. Fragments suitable for use include fragments encompassing T-helper epitopes. In particular the protein D fragment from *H. influenzae* will preferably contain the N-terminal 1/3 of the protein. Protein D is an IgD-binding protein from *Haemophilus influenzae* (EP 0 594 610 B1) and is a potential immunogen. In addition, staphylococcal proteins may be used as a carrier protein in the polysaccharide conjugates of the invention.

A carrier protein that would be particularly advantageous to use in the context of a staphylococcal vaccine is staphylococcal alpha toxoid. The native form may be conjugated to a polysaccharide since the process of conjugation reduces toxicity. Preferably genetically detoxified alpha toxins such as the His35Leu or His35Arg variants are used as carriers since residual toxicity is lower. Alternatively the alpha toxin is chemically detoxified by treatment with a cross-linking reagent, formaldehyde or glutaraldehyde. A genetically detoxified alpha toxin is optionally chemically detoxified,

preferably by treatment with a cross-linking reagent, formaldehyde or glutaraldehyde to further reduce toxicity.

The polysaccharides may be linked to the carrier protein(s) by any known method (for example those methods described in U.S. Pat. Nos. 4,372,945, 4,474,757, and 4,356,170). Preferably, CDAP conjugation chemistry is carried out (see WO95/08348). In CDAP, the cyanylating reagent 1-cyano-dimethylaminopyridinium tetrafluoroborate (CDAP) is preferably used for the synthesis of polysaccharide-protein conjugates. The cyanilation reaction can be performed under relatively mild conditions, which avoids hydrolysis of the alkaline sensitive polysaccharides. This synthesis allows direct coupling to a carrier protein.

Conjugation preferably involves producing a direct linkage between the carrier protein and polysaccharide. Optionally a spacer (such as adipic dihydride (ADH)) may be introduced between the carrier protein and the polysaccharide.

#### IV. IMMUNE RESPONSE AND ASSAYS

As discussed above, the invention concerns evoking or inducing an immune response in a subject against a variant SpA or coagulase peptide. In one embodiment, the immune response can protect against or treat a subject having, suspected of having, or at risk of developing an infection or related disease, particularly those related to staphylococci. One use of the immunogenic compositions of the invention is to prevent nosocomial infections by inoculating a subject prior to undergoing procedures in a hospital or other environment having an increased risk of infection.

##### A. Immunoassays

The present invention includes the implementation of serological assays to evaluate whether and to what extent an immune response is induced or evoked by compositions of the invention. There are many types of immunoassays that can be implemented. Immunoassays encompassed by the present invention include, but are not limited to, those described in U.S. Pat. No. 4,367,110 (double monoclonal antibody sandwich assay) and U.S. Pat. No. 4,452,901 (western blot). Other assays include immunoprecipitation of labeled ligands and immunocytochemistry, both in vitro and in vivo.

Immunoassays generally are binding assays. Certain preferred immunoassays are the various types of enzyme linked immunosorbent assays (ELISAs) and radioimmunoassays (RIA) known in the art. Immunohistochemical detection using tissue sections is also particularly useful. In one example, antibodies or antigens are immobilized on a selected surface, such as a well in a polystyrene microtiter plate, dipstick, or column support. Then, a test composition suspected of containing the desired antigen or antibody, such as a clinical sample, is added to the wells. After binding and washing to remove non specifically bound immune complexes, the bound antigen or antibody may be detected. Detection is generally achieved by the addition of another antibody, specific for the desired antigen or antibody, that is linked to a detectable label. This type of ELISA is known as a "sandwich ELISA." Detection also may be achieved by the addition of a second antibody specific for the desired antigen, followed by the addition of a third antibody that has binding affinity for the second antibody, with the third antibody being linked to a detectable label.

Competition ELISAs are also possible implementations in which test samples compete for binding with known amounts of labeled antigens or antibodies. The amount of reactive species in the unknown sample is determined by mixing the sample with the known labeled species before or during incu-

bation with coated wells. The presence of reactive species in the sample acts to reduce the amount of labeled species available for binding to the well and thus reduces the ultimate signal. Irrespective of the format employed, ELISAs have certain features in common, such as coating, incubating or binding, washing to remove non specifically bound species, and detecting the bound immune complexes.

Antigen or antibodies may also be linked to a solid support, such as in the form of plate, beads, dipstick, membrane, or column matrix, and the sample to be analyzed is applied to the immobilized antigen or antibody. In coating a plate with either antigen or antibody, one will generally incubate the wells of the plate with a solution of the antigen or antibody, either overnight or for a specified period. The wells of the plate will then be washed to remove incompletely-adsorbed material. Any remaining available surfaces of the wells are then "coated" with a nonspecific protein that is antigenically neutral with regard to the test antisera. These include bovine serum albumin (BSA), casein, and solutions of milk powder. The coating allows for blocking of nonspecific adsorption sites on the immobilizing surface and thus reduces the background caused by nonspecific binding of antisera onto the surface.

##### B. Diagnosis of Bacterial Infection

In addition to the use of proteins, polypeptides, and/or peptides, as well as antibodies binding these polypeptides, proteins, and/or peptides, to treat or prevent infection as described above, the present invention contemplates the use of these polypeptides, proteins, peptides, and/or antibodies in a variety of ways, including the detection of the presence of Staphylococci to diagnose an infection, whether in a patient or on medical equipment which may also become infected. In accordance with the invention, a preferred method of detecting the presence of infections involves the steps of obtaining a sample suspected of being infected by one or more staphylococcal bacteria species or strains, such as a sample taken from an individual, for example, from one's blood, saliva, tissues, bone, muscle, cartilage, or skin. Following isolation of the sample, diagnostic assays utilizing the polypeptides, proteins, peptides, and/or antibodies of the present invention may be carried out to detect the presence of staphylococci, and such assay techniques for determining such presence in a sample are well known to those skilled in the art and include methods such as radioimmunoassay, western blot analysis and ELISA assays. In general, in accordance with the invention, a method of diagnosing an infection is contemplated wherein a sample suspected of being infected with staphylococci has added to it the polypeptide, protein, peptide, antibody, or monoclonal antibody in accordance with the present invention, and staphylococci are indicated by antibody binding to the polypeptides, proteins, and/or peptides, or polypeptides, proteins, and/or peptides binding to the antibodies in the sample.

Accordingly, antibodies in accordance with the invention may be used for the prevention of infection from staphylococcal bacteria (i.e., passive immunization), for the treatment of an ongoing infection, or for use as research tools. The term "antibodies" as used herein includes monoclonal, polyclonal, chimeric, single chain, bispecific, simianized, and humanized or primatized antibodies as well as Fab fragments, such as those fragments which maintain the binding specificity of the antibodies, including the products of an Fab immunoglobulin expression library. Accordingly, the invention contemplates the use of single chains such as the variable heavy and light chains of the antibodies. Generation of any of these types of antibodies or antibody fragments is well known to those skilled in the art. Specific examples of the generation of an

antibody to a bacterial protein can be found in U.S. Patent Application Pub. No. 20030153022, which is incorporated herein by reference in its entirety.

Any of the above described polypeptides, proteins, peptides, and/or antibodies may be labeled directly with a detectable label for identification and quantification of staphylococcal bacteria. Labels for use in immunoassays are generally known to those skilled in the art and include enzymes, radioisotopes, and fluorescent, luminescent and chromogenic substances, including colored particles such as colloidal gold or latex beads. Suitable immunoassays include enzyme-linked immunosorbent assays (ELISA).

### C. Protective Immunity

In some embodiments of the invention, proteinaceous compositions confer protective immunity to a subject. Protective immunity refers to a body's ability to mount a specific immune response that protects the subject from developing a particular disease or condition that involves the agent against which there is an immune response. An immunogenically effective amount is capable of conferring protective immunity to the subject.

As used herein in the specification and in the claims section that follows, the term polypeptide or peptide refer to a stretch of amino acids covalently linked there amongst via peptide bonds. Different polypeptides have different functionalities according to the present invention. While according to one aspect, a polypeptide is derived from an immunogen designed to induce an active immune response in a recipient, according to another aspect of the invention, a polypeptide is derived from an antibody which results following the elicitation of an active immune response in, for example, an animal, and which can serve to induce a passive immune response in the recipient. In both cases, however, the polypeptide is encoded by a polynucleotide according to any possible codon usage.

As used herein the phrase "immune response" or its equivalent "immunological response" refers to the development of a humoral (antibody mediated), cellular (mediated by antigen-specific T cells or their secretion products) or both humoral and cellular response directed against a protein, peptide, carbohydrate, or polypeptide of the invention in a recipient patient. Such a response can be an active response induced by administration of immunogen or a passive response induced by administration of antibody, antibody containing material, or primed T-cells. A cellular immune response is elicited by the presentation of polypeptide epitopes in association with Class I or Class II MHC molecules, to activate antigen-specific CD4 (+) T helper cells and/or CD8 (+) cytotoxic T cells. The response may also involve activation of monocytes, macrophages, NK cells, basophils, dendritic cells, astrocytes, microglia cells, eosinophils or other components of innate immunity. As used herein "active immunity" refers to any immunity conferred upon a subject by administration of an antigen.

As used herein "passive immunity" refers to any immunity conferred upon a subject without administration of an antigen to the subject. "Passive immunity" therefore includes, but is not limited to, administration of activated immune effectors including cellular mediators or protein mediators (e.g., monoclonal and/or polyclonal antibodies) of an immune response. A monoclonal or polyclonal antibody composition may be used in passive immunization for the prevention or treatment of infection by organisms that carry the antigen recognized by the antibody. An antibody composition may include antibodies that bind to a variety of antigens that may in turn be associated with various organisms. The antibody component can be a polyclonal antiserum. In certain aspects the antibody or antibodies are affinity purified from an animal

or second subject that has been challenged with an antigen(s). Alternatively, an antibody mixture may be used, which is a mixture of monoclonal and/or polyclonal antibodies to antigens present in the same, related, or different microbes or organisms, such as gram-positive bacteria, gram-negative bacteria, including but not limited to *staphylococcus* bacteria.

Passive immunity may be imparted to a patient or subject by administering to the patient immunoglobulins (Ig) and/or other immune factors obtained from a donor or other non-patient source having a known immunoreactivity. In other aspects, an antigenic composition of the present invention can be administered to a subject who then acts as a source or donor for globulin, produced in response to challenge with the antigenic composition ("hyperimmune globulin"), that contains antibodies directed against *Staphylococcus* or other organism. A subject thus treated would donate plasma from which hyperimmune globulin would then be obtained, via conventional plasma-fractionation methodology, and administered to another subject in order to impart resistance against or to treat *staphylococcus* infection. Hyperimmune globulins according to the invention are particularly useful for immunocompromised individuals, for individuals undergoing invasive procedures or where time does not permit the individual to produce their own antibodies in response to vaccination. See U.S. Pat. Nos. 6,936,258, 6,770,278, 6,756,361, 5,548,066, 5,512,282, 4,338,298, and 4,748,018, each of which is incorporated herein by reference in its entirety, for exemplary methods and compositions related to passive immunity.

For purposes of this specification and the accompanying claims the terms "epitope" and "antigenic determinant" are used interchangeably to refer to a site on an antigen to which B and/or T cells respond or recognize. B-cell epitopes can be formed both from contiguous amino acids or noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained on exposure to denaturing solvents whereas epitopes formed by tertiary folding are typically lost on treatment with denaturing solvents. An epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids in a unique spatial conformation. Methods of determining spatial conformation of epitopes include, for example, x-ray crystallography and 2-dimensional nuclear magnetic resonance. See, e.g., Epitope Mapping Protocols (1996). Antibodies that recognize the same epitope can be identified in a simple immunoassay showing the ability of one antibody to block the binding of another antibody to a target antigen. T-cells recognize continuous epitopes of about nine amino acids for CD8 cells or about 13-15 amino acids for CD4 cells. T cells that recognize the epitope can be identified by in vitro assays that measure antigen-dependent proliferation, as determined by <sup>3</sup>H-thymidine incorporation by primed T cells in response to an epitope (Burke et al., 1994), by antigen-dependent killing (cytotoxic T lymphocyte assay, Tigges et al., 1996) or by cytokine secretion.

The presence of a cell-mediated immunological response can be determined by proliferation assays (CD4 (+) T cells) or CTL (cytotoxic T lymphocyte) assays. The relative contributions of humoral and cellular responses to the protective or therapeutic effect of an immunogen can be distinguished by separately isolating IgG and T-cells from an immunized syngeneic animal and measuring protective or therapeutic effect in a second subject.

As used herein and in the claims, the terms "antibody" or "immunoglobulin" are used interchangeably and refer to any of several classes of structurally related proteins that function

as part of the immune response of an animal or recipient, which proteins include IgG, IgD, IgE, IgA, IgM and related proteins.

Under normal physiological conditions antibodies are found in plasma and other body fluids and in the membrane of certain cells and are produced by lymphocytes of the type denoted B cells or their functional equivalent. Antibodies of the IgG class are made up of four polypeptide chains linked together by disulfide bonds. The four chains of intact IgG molecules are two identical heavy chains referred to as H-chains and two identical light chains referred to as L-chains.

In order to produce polyclonal antibodies, a host, such as a rabbit or goat, is immunized with the antigen or antigen fragment, generally with an adjuvant and, if necessary, coupled to a carrier. Antibodies to the antigen are subsequently collected from the sera of the host. The polyclonal antibody can be affinity purified against the antigen rendering it monospecific.

Monoclonal antibodies can be produced by hyperimmunization of an appropriate donor with the antigen or ex-vivo by use of primary cultures of splenic cells or cell lines derived from spleen (Anavi, 1998; Huston et al., 1991; Johnson et al., 1991; Mernaugh et al., 1995).

As used herein and in the claims, the phrase "an immunological portion of an antibody" includes a Fab fragment of an antibody, a Fv fragment of an antibody, a heavy chain of an antibody, a light chain of an antibody, a heterodimer consisting of a heavy chain and a light chain of an antibody, a variable fragment of a light chain of an antibody, a variable fragment of a heavy chain of an antibody, and a single chain variant of an antibody, which is also known as scFv. In addition, the term includes chimeric immunoglobulins which are the expression products of fused genes derived from different species, one of the species can be a human, in which case a chimeric immunoglobulin is said to be humanized. Typically, an immunological portion of an antibody competes with the intact antibody from which it was derived for specific binding to an antigen.

Optionally, an antibody or preferably an immunological portion of an antibody, can be chemically conjugated to, or expressed as, a fusion protein with other proteins. For purposes of this specification and the accompanying claims, all such fused proteins are included in the definition of antibodies or an immunological portion of an antibody.

As used herein the terms "immunogenic agent" or "immunogen" or "antigen" are used interchangeably to describe a molecule capable of inducing an immunological response against itself on administration to a recipient, either alone, in conjunction with an adjuvant, or presented on a display vehicle.

#### D. Treatment Methods

A method of the present invention includes treatment for a disease or condition caused by a *staphylococcus* pathogen. An immunogenic polypeptide of the invention can be given to induce an immune response in a person infected with *staphylococcus* or suspected of having been exposed to *staphylococcus*. Methods may be employed with respect to individuals who have tested positive for exposure to *staphylococcus* or who are deemed to be at risk for infection based on possible exposure.

In particular, the invention encompasses a method of treatment for staphylococcal infection, particularly hospital acquired nosocomial infections. The immunogenic compositions and vaccines of the invention are particularly advantageous to use in cases of elective surgery. Such patients will know the date of surgery in advance and could be inoculated

in advance. The immunogenic compositions and vaccines of the invention are also advantageous to use to inoculate health care workers.

In some embodiments, the treatment is administered in the presence of adjuvants or carriers or other staphylococcal antigens. Furthermore, in some examples, treatment comprises administration of other agents commonly used against bacterial infection, such as one or more antibiotics.

The use of peptides for vaccination can require, but not necessarily, conjugation of the peptide to an immunogenic carrier protein, such as hepatitis B surface antigen, keyhole limpet hemocyanin, or bovine serum albumin. Methods for performing this conjugation are well known in the art.

## V. VACCINE AND OTHER PHARMACEUTICAL COMPOSITIONS AND ADMINISTRATION

### A. Vaccines

The present invention includes methods for preventing or ameliorating staphylococcal infections, particularly hospital acquired nosocomial infections. As such, the invention contemplates vaccines for use in both active and passive immunization embodiments. Immunogenic compositions, proposed to be suitable for use as a vaccine, may be prepared from immunogenic SpA polypeptide(s), such as a SpA domain D variant, or immunogenic coagulases. In other embodiments SpA or coagulases can be used in combination with other secreted virulence proteins, surface proteins or immunogenic fragments thereof. In certain aspects, antigenic material is extensively dialyzed to remove undesired small molecular weight molecules and/or lyophilized for more ready formulation into a desired vehicle.

Other options for a protein/peptide-based vaccine involve introducing nucleic acids encoding the antigen(s) as DNA vaccines. In this regard, recent reports described construction of recombinant vaccinia viruses expressing either 10 contiguous minimal CTL epitopes (Thomson, 1996) or a combination of B cell, cytotoxic T-lymphocyte (CTL), and T-helper (Th) epitopes from several microbes (An, 1997), and successful use of such constructs to immunize mice for priming protective immune responses. Thus, there is ample evidence in the literature for successful utilization of peptides, peptide-pulsed antigen presenting cells (APCs), and peptide-encoding constructs for efficient in vivo priming of protective immune responses. The use of nucleic acid sequences as vaccines is exemplified in U.S. Pat. Nos. 5,958,895 and 5,620,896.

The preparation of vaccines that contain polypeptide or peptide sequence(s) as active ingredients is generally well understood in the art, as exemplified by U.S. Pat. Nos. 4,608,251; 4,601,903; 4,599,231; 4,599,230; 4,596,792; and 4,578,770, all of which are incorporated herein by reference. Typically, such vaccines are prepared as injectables either as liquid solutions or suspensions: solid forms suitable for solution in or suspension in liquid prior to injection may also be prepared. The preparation may also be emulsified. The active immunogenic ingredient is often mixed with excipients that are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vaccine may contain amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, or adjuvants that enhance the effectiveness of the vaccines. In specific embodiments, vaccines are formulated with a combination of substances, as described in U.S. Pat. Nos. 6,793,923 and 6,733,754, which are incorporated herein by reference.

Vaccines may be conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories and, in some cases, oral formulations. For suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or triglycerides: such suppositories may be formed from mixtures containing the active ingredient in the range of about 0.5% to about 10%, preferably about 1% to about 2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain about 10% to about 95% of active ingredient, preferably about 25% to about 70%.

The polypeptides and polypeptide-encoding DNA constructs may be formulated into a vaccine as neutral or salt forms. Pharmaceutically-acceptable salts include the acid addition salts (formed with the free amino groups of the peptide) and those that are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like.

Typically, vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective and immunogenic. The quantity to be administered depends on the subject to be treated, including the capacity of the individual's immune system to synthesize antibodies and the degree of protection desired. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner. However, suitable dosage ranges are of the order of several hundred micrograms of active ingredient per vaccination. Suitable regimes for initial administration and booster shots are also variable, but are typified by an initial administration followed by subsequent inoculations or other administrations.

The manner of application may be varied widely. Any of the conventional methods for administration of a vaccine are applicable. These are believed to include oral application within a solid physiologically acceptable base or in a physiologically acceptable dispersion, parenterally, by injection and the like. The dosage of the vaccine will depend on the route of administration and will vary according to the size and health of the subject.

In certain instances, it will be desirable to have multiple administrations of the vaccine, e.g., 2, 3, 4, 5, 6 or more administrations. The vaccinations can be at 1, 2, 3, 4, 5, 6, 7, 8, to 5, 6, 7, 8, 9, 10, 11, 12 twelve week intervals, including all ranges there between. Periodic boosters at intervals of 1-5 years will be desirable to maintain protective levels of the antibodies. The course of the immunization may be followed by assays for antibodies against the antigens, as described in U.S. Pat. Nos. 3,791,932; 4,174,384 and 3,949,064.

#### 1. Carriers

A given composition may vary in its immunogenicity. It is often necessary therefore to boost the host immune system, as may be achieved by coupling a peptide or polypeptide to a carrier. Exemplary and preferred carriers are keyhole limpet hemocyanin (KLH) and bovine serum albumin (BSA). Other albumins such as ovalbumin, mouse serum albumin, or rabbit serum albumin can also be used as carriers. Means for conjugating a polypeptide to a carrier protein are well known in the art and include glutaraldehyde, m-maleimidobencoyl-N-hydroxysuccinimide ester, carbodiimide, and bis-biazotized benzidine.

#### 2. Adjuvants

The immunogenicity of polypeptide or peptide compositions can be enhanced by the use of non-specific stimulators of the immune response, known as adjuvants. Suitable adjuvants include all acceptable immunostimulatory compounds, such as cytokines, toxins, or synthetic compositions. A number of adjuvants can be used to enhance an antibody response against a variant SpA polypeptide or coagulase, or any other bacterial protein or combination contemplated herein. Adjuvants can (1) trap the antigen in the body to cause a slow release; (2) attract cells involved in the immune response to the site of administration; (3) induce proliferation or activation of immune system cells; or (4) improve the spread of the antigen throughout the subject's body.

Adjuvants include, but are not limited to, oil-in-water emulsions, water-in-oil emulsions, mineral salts, polynucleotides, and natural substances. Specific adjuvants that may be used include IL-1, IL-2, IL-4, IL-7, IL-12,  $\gamma$ -interferon, GMCSF, BCG, aluminum salts, such as aluminum hydroxide or other aluminum compound, MDP compounds, such as thur-MDP and nor-MDP, CGP (MTP-PE), lipid A, and monophosphoryl lipid A (MPL). RIBI, which contains three components extracted from bacteria, MPL, trehalose dimycolate (TDM), and cell wall skeleton (CWS) in a 2% squalene/Tween 80 emulsion. MHC antigens may even be used. Others adjuvants or methods are exemplified in U.S. Pat. Nos. 6,814, 971, 5,084,269, 6,656,462, each of which is incorporated herein by reference).

Various methods of achieving adjuvant affect for the vaccine includes use of agents such as aluminum hydroxide or phosphate (alum), commonly used as about 0.05 to about 0.1% solution in phosphate buffered saline, admixture with synthetic polymers of sugars (Carbopol®) used as an about 0.25% solution, aggregation of the protein in the vaccine by heat treatment with temperatures ranging between about 70° to about 101° C. for a 30-second to 2-minute period, respectively. Aggregation by reactivating with pepsin-treated (Fab) antibodies to albumin; mixture with bacterial cells (e.g., *C. parvum*), endotoxins or lipopolysaccharide components of Gram-negative bacteria; emulsion in physiologically acceptable oil vehicles (e.g., mannide mono-oleate (Aracel A)); or emulsion with a 20% solution of a perfluorocarbon (Fluosol-DA®) used as a block substitute may also be employed to produce an adjuvant effect.

Examples of and often preferred adjuvants include complete Freund's adjuvant (a non-specific stimulator of the immune response containing killed *Mycobacterium tuberculosis*), incomplete Freund's adjuvants, and aluminum hydroxide.

In some aspects, it is preferred that the adjuvant be selected to be a preferential inducer of either a Th1 or a Th2 type of response. High levels of Th1-type cytokines tend to favor the induction of cell mediated immune responses to a given antigen, while high levels of Th2-type cytokines tend to favor the induction of humoral immune responses to the antigen.

The distinction of Th1 and Th2-type immune response is not absolute. In reality an individual will support an immune response which is described as being predominantly Th1 or predominantly Th2. However, it is often convenient to consider the families of cytokines in terms of that described in murine CD4+ T cell clones by Mosmann and Coffman (Mosmann, and Coffman, 1989). Traditionally, Th1-type responses are associated with the production of the INF- $\gamma$  and IL-2 cytokines by T-lymphocytes. Other cytokines often directly associated with the induction of Th1-type immune

responses are not produced by T-cells, such as IL-12. In contrast, Th2-type responses are associated with the secretion of IL-4, IL-5, IL-6, IL-10.

In addition to adjuvants, it may be desirable to co-administer biologic response modifiers (BRM) to enhance immune responses. BRMs have been shown to upregulate T cell immunity or downregulate suppressor cell activity. Such BRMs include, but are not limited to, Cimetidine (CIM; 1200 mg/d) (Smith/Kline, PA); or low-dose Cyclophosphamide (CYP; 300 mg/m<sup>2</sup>) (Johnson/Mead, NJ) and cytokines such as  $\gamma$ -interferon, IL-2, or IL-12 or genes encoding proteins involved in immune helper functions, such as B-7.

#### B. Lipid Components and Moieties

In certain embodiments, the present invention concerns compositions comprising one or more lipids associated with a nucleic acid or a polypeptide/peptide. A lipid is a substance that is insoluble in water and extractable with an organic solvent. Compounds other than those specifically described herein are understood by one of skill in the art as lipids, and are encompassed by the compositions and methods of the present invention. A lipid component and a non-lipid may be attached to one another, either covalently or non-covalently.

A lipid may be a naturally occurring lipid or a synthetic lipid. However, a lipid is usually a biological substance. Biological lipids are well known in the art, and include for example, neutral fats, phospholipids, phosphoglycerides, steroids, terpenes, lysolipids, glycosphingolipids, glucolipids, sulphatides, lipids with ether and ester-linked fatty acids and polymerizable lipids, and combinations thereof.

A nucleic acid molecule or a polypeptide/peptide, associated with a lipid may be dispersed in a solution containing a lipid, dissolved with a lipid, emulsified with a lipid, mixed with a lipid, combined with a lipid, covalently bonded to a lipid, contained as a suspension in a lipid or otherwise associated with a lipid. A lipid or lipid-poxvirus-associated composition of the present invention is not limited to any particular structure. For example, they may also simply be interspersed in a solution, possibly forming aggregates which are not uniform in either size or shape. In another example, they may be present in a bilayer structure, as micelles, or with a "collapsed" structure. In another non-limiting example, a lipofectamine (Gibco BRL)-poxvirus or Superfect (Qiagen)-poxvirus complex is also contemplated.

In certain embodiments, a composition may comprise about 1%, about 2%, about 3%, about 4% about 5%, about 6%, about 7%, about 8%, about 9%, about 10%, about 11%, about 12%, about 13%, about 14%, about 15%, about 16%, about 17%, about 18%, about 19%, about 20%, about 21%, about 22%, about 23%, about 24%, about 25%, about 26%, about 27%, about 28%, about 29%, about 30%, about 31%, about 32%, about 33%, about 34%, about 35%, about 36%, about 37%, about 38%, about 39%, about 40%, about 41%, about 42%, about 43%, about 44%, about 45%, about 46%, about 47%, about 48%, about 49%, about 50%, about 51%, about 52%, about 53%, about 54%, about 55%, about 56%, about 57%, about 58%, about 59%, about 60%, about 61%, about 62%, about 63%, about 64%, about 65%, about 66%, about 67%, about 68%, about 69%, about 70%, about 71%, about 72%, about 73%, about 74%, about 75%, about 76%, about 77%, about 78%, about 79%, about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or any range therebetween, of a particular lipid, lipid type, or non-lipid component such as an adjuvant, antigen, peptide, polypeptide, sugar, nucleic acid or other material disclosed herein or as would be

known to one of skill in the art. In a non-limiting example, a composition may comprise about 10% to about 20% neutral lipids, and about 33% to about 34% of a cerebroside, and about 1% cholesterol. In another non-limiting example, a liposome may comprise about 4% to about 12% terpenes, wherein about 1% of the micelle is specifically lycopene, leaving about 3% to about 11% of the liposome as comprising other terpenes; and about 10% to about 35% phosphatidyl choline, and about 1% of a non-lipid component. Thus, it is contemplated that compositions of the present invention may comprise any of the lipids, lipid types or other components in any combination or percentage range.

#### C. Combination Therapy

The compositions and related methods of the present invention, particularly administration of a secreted virulence factor or surface protein, including a variant SpA polypeptide or peptide, and/or other bacterial peptides or proteins to a patient/subject, may also be used in combination with the administration of traditional therapies. These include, but are not limited to, the administration of antibiotics such as streptomycin, ciprofloxacin, doxycycline, gentamycin, chloramphenicol, trimethoprim, sulfamethoxazole, ampicillin, tetracycline or various combinations of antibiotics.

In one aspect, it is contemplated that a polypeptide vaccine and/or therapy is used in conjunction with antibacterial treatment. Alternatively, the therapy may precede or follow the other agent treatment by intervals ranging from minutes to weeks. In embodiments where the other agents and/or proteins or polynucleotides are administered separately, one would generally ensure that a significant period of time did not expire between the time of each delivery, such that the agent and antigenic composition would still be able to exert an advantageously combined effect on the subject. In such instances, it is contemplated that one may administer both modalities within about 12-24 h of each other or within about 6-12 h of each other. In some situations, it may be desirable to extend the time period for administration significantly, where several days (2, 3, 4, 5, 6 or 7) to several weeks (1, 2, 3, 4, 5, 6, 7 or 8) lapse between the respective administrations.

Various combinations may be employed, for example antibiotic therapy is "A" and the immunogenic molecule given as part of an immune therapy regime, such as an antigen, is "B":  
A/B/A B/A/B B/B/A A/A/B A/B/B B/A/A A/B/B/B B/A/B/B

B/B/B/A B/B/A/B A/A/B/B A/B/A/B A/B/B/A B/B/A/A B/A/B/A B/A/A/B A/A/A/B B/A/A/A A/B/A/A A/A/B/A

Administration of the immunogenic compositions of the present invention to a patient/subject will follow general protocols for the administration of such compounds, taking into account the toxicity, if any, of the SpA composition, or other compositions described herein. It is expected that the treatment cycles would be repeated as necessary. It also is contemplated that various standard therapies, such as hydration, may be applied in combination with the described therapy.

#### D. General Pharmaceutical Compositions

In some embodiments, pharmaceutical compositions are administered to a subject. Different aspects of the present invention involve administering an effective amount of a composition to a subject. In some embodiments of the present invention, staphylococcal antigens, members of the Ess pathway, including polypeptides or peptides of the Esa or Esx class, and/or members of sortase substrates may be administered to the patient to protect against infection by one or more *staphylococcus* pathogens. Alternatively, an expression vector encoding one or more such polypeptides or peptides may be given to a patient as a preventative treatment. Additionally, such compounds can be administered in combination with an

antibiotic or an antibacterial. Such compositions will generally be dissolved or dispersed in a pharmaceutically acceptable carrier or aqueous medium.

In addition to the compounds formulated for parenteral administration, such as those for intravenous or intramuscular injection, other pharmaceutically acceptable forms include, e.g., tablets or other solids for oral administration; time release capsules; and any other form currently used, including creams, lotions, mouthwashes, inhalants and the like.

The active compounds of the present invention can be formulated for parenteral administration, e.g., formulated for injection via the intravenous, intramuscular, subcutaneous, or even intraperitoneal routes. The preparation of an aqueous composition that contains a compound or compounds that increase the expression of an MHC class I molecule will be known to those of skill in the art in light of the present disclosure. Typically, such compositions can be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for use to prepare solutions or suspensions upon the addition of a liquid prior to injection can also be prepared; and, the preparations can also be emulsified.

Solutions of the active compounds as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions; formulations including sesame oil, peanut oil, or aqueous propylene glycol; and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that it may be easily injected. It also should be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi.

The proteinaceous compositions may be formulated into a neutral or salt form. Pharmaceutically acceptable salts, include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like.

The carrier also can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion, and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropri-

ate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques, which yield a powder of the active ingredient, plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Administration of the compositions according to the present invention will typically be via any common route. This includes, but is not limited to oral, nasal, or buccal administration. Alternatively, administration may be by orthotopic, intradermal, subcutaneous, intramuscular, intraperitoneal, intranasal, or intravenous injection. In certain embodiments, a vaccine composition may be inhaled (e.g., U.S. Pat. No. 6,651,655, which is specifically incorporated by reference). Such compositions would normally be administered as pharmaceutically acceptable compositions that include physiologically acceptable carriers, buffers or other excipients. As used herein, the term "pharmaceutically acceptable" refers to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem complications commensurate with a reasonable benefit/risk ratio. The term "pharmaceutically acceptable carrier," means a pharmaceutically acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting a chemical agent.

For parenteral administration in an aqueous solution, for example, the solution should be suitably buffered, if necessary, and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous, and intraperitoneal administration. In this connection, sterile aqueous media which can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage could be dissolved in isotonic NaCl solution and either added to hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, Remington's Pharmaceutical Sciences, 1990). Some variation in dosage will necessarily occur depending on the condition of the subject. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject.

An effective amount of therapeutic or prophylactic composition is determined based on the intended goal. The term "unit dose" or "dosage" refers to physically discrete units suitable for use in a subject, each unit containing a predetermined quantity of the composition calculated to produce the desired responses discussed above in association with its administration, i.e., the appropriate route and regimen. The quantity to be administered, both according to number of treatments and unit dose, depends on the protection desired.

Precise amounts of the composition also depend on the judgment of the practitioner and are peculiar to each individual. Factors affecting dose include physical and clinical state of the subject, route of administration, intended goal of treatment (alleviation of symptoms versus cure), and potency, stability, and toxicity of the particular composition.

Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such

amount as is therapeutically or prophylactically effective. The formulations are easily administered in a variety of dosage forms, such as the type of injectable solutions described above.

#### E. In Vitro, Ex Vivo, or In Vivo Administration

As used herein, the term in vitro administration refers to manipulations performed on cells removed from or outside of a subject, including, but not limited to cells in culture. The term ex vivo administration refers to cells which have been manipulated in vitro, and are subsequently administered to a subject. The term in vivo administration includes all manipulations performed within a subject.

In certain aspects of the present invention, the compositions may be administered either in vitro, ex vivo, or in vivo. In certain in vitro embodiments, autologous B-lymphocyte cell lines are incubated with a virus vector of the instant invention for 24 to 48 hours or with a variant SpA and/or coagulase and/or any other composition described herein for two hours. The transduced cells can then be used for in vitro analysis, or alternatively for ex vivo administration. U.S. Pat. Nos. 4,690,915 and 5,199,942, both incorporated herein by reference, disclose methods for ex vivo manipulation of blood mononuclear cells and bone marrow cells for use in therapeutic applications.

#### F. Antibodies and Passive Immunization

Another aspect of the invention is a method of preparing an immunoglobulin for use in prevention or treatment of staphylococcal infection comprising the steps of immunizing a recipient or donor with the vaccine of the invention and isolating immunoglobulin from the recipient or donor. An immunoglobulin prepared by this method is a further aspect of the invention. A pharmaceutical composition comprising the immunoglobulin of the invention and a pharmaceutically acceptable carrier is a further aspect of the invention which could be used in the manufacture of a medicament for the treatment or prevention of staphylococcal disease. A method for treatment or prevention of staphylococcal infection comprising a step of administering to a patient an effective amount of the pharmaceutical preparation of the invention is a further aspect of the invention.

Inocula for polyclonal antibody production are typically prepared by dispersing the antigenic composition in a physiologically tolerable diluent such as saline or other adjuvants suitable for human use to form an aqueous composition. An immunostimulatory amount of inoculum is administered to a mammal and the inoculated mammal is then maintained for a time sufficient for the antigenic composition to induce protective antibodies.

The antibodies can be isolated to the extent desired by well known techniques such as affinity chromatography (Harlow and Lane, 1988). Antibodies can include antiserum preparations from a variety of commonly used animals, e.g. goats, primates, donkeys, swine, horses, guinea pigs, rats or man.

An immunoglobulin produced in accordance with the present invention can include whole antibodies, antibody fragments or subfragments. Antibodies can be whole immunoglobulins of any class (e.g., IgG, IgM, IgA, IgD or IgE), chimeric antibodies or hybrid antibodies with dual specificity to two or more antigens of the invention. They may also be fragments (e.g., F(ab')<sub>2</sub>, Fab', Fab, Fv and the like) including hybrid fragments. An immunoglobulin also includes natural, synthetic, or genetically engineered proteins that act like an antibody by binding to specific antigens to form a complex.

A vaccine of the present invention can be administered to a recipient who then acts as a source of immunoglobulin, produced in response to challenge from the specific vaccine. A subject thus treated would donate plasma from which hyper-

immune globulin would be obtained via conventional plasma fractionation methodology. The hyperimmune globulin would be administered to another subject in order to impart resistance against or treat staphylococcal infection. Hyperimmune globulins of the invention are particularly useful for treatment or prevention of staphylococcal disease in infants, immune compromised individuals, or where treatment is required and there is no time for the individual to produce antibodies in response to vaccination.

An additional aspect of the invention is a pharmaceutical composition comprising two or more monoclonal antibodies (or fragments thereof; preferably human or humanised) reactive against at least two constituents of the immunogenic composition of the invention, which could be used to treat or prevent infection by Gram positive bacteria, preferably staphylococci, more preferably *S. aureus* or *S. epidermidis*. Such pharmaceutical compositions comprise monoclonal antibodies that can be whole immunoglobulins of any class, chimeric antibodies, or hybrid antibodies with specificity to two or more antigens of the invention. They may also be fragments (e.g., F(ab')<sub>2</sub>, Fab', Fab, Fv and the like) including hybrid fragments.

Methods of making monoclonal antibodies are well known in the art and can include the fusion of splenocytes with myeloma cells (Kohler and Milstein, 1975; Harlow and Lane, 1988). Alternatively, monoclonal Fv fragments can be obtained by screening a suitable phage display library (Vaughan et al., 1998). Monoclonal antibodies may be humanized or part humanized by known methods.

## VI. EXAMPLES

The following examples are given for the purpose of illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion. One skilled in the art will appreciate readily that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those objects, ends and advantages inherent herein. The present examples, along with the methods described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses which are encompassed within the spirit of the invention as defined by the scope of the claims will occur to those skilled in the art.

### Example 1

#### Non-Toxicogenic Protein A Variants as Subunit Vaccines to Prevent *Staphylococcus aureus* Infections

##### A. Results

An animal model for *S. aureus* infection BALB/c mice were infected by intravenous injection with  $1 \times 10^7$  CFU of the human clinical isolate *S. aureus* Newman (Baba et al., 2007). Within 6 hours following infection, 99.999% of staphylococci disappeared from the blood stream and were distributed via the vasculature. Staphylococcal dissemination to peripheral tissues occurred rapidly, as the bacterial load in kidney and other peripheral organ tissues reached  $1 \times 10^5$  CFU g<sup>-1</sup> within the first three hours. The staphylococcal load in kidney tissues increased by 1.5 log CFU within twenty-four hours. Forty-eight hours following infection, mice developed disseminated abscesses in multiple organs, detectable by light microscopy of hematoxylin-eosin stained, thin-sectioned kidney tissue. The initial abscess diameter was 524 μm (±65

μM); lesions were initially marked by an influx of polymorphonuclear leukocytes (PMNs) and harbored no discernable organization of staphylococci, most of which appeared to reside within PMNs. On day 5 of infection, abscesses increased in size and enclosed a central population of staphylococci, surrounded by a layer of eosinophilic, amorphous material and a large cuff of PMNs. Histopathology revealed massive necrosis of PMNs in proximity to the staphylococcal nidus at the center of abscess lesions as well as a mantle of

populations at the abscess center, designated staphylococcal abscess communities (SAC), appeared homogenous and coated by an electron-dense, granular material. The kinetics of the appearance of infectious lesions and the morphological attributes of abscesses formed by *S. aureus* Newman were similar to those observed following mouse infection with *S. aureus* USA300 (LAC), the current epidemic community-acquired methicillin-resistant *S. aureus* (CA-MRSA) clone in the United States (Diep et al., 2006).

TABLE 4

Genotype	Genetic requirements for <i>S. aureus</i> Newman abscess formation in mice					
	Staphylococcal load in kidney tissue			Abscess formation in kidney tissue		
	<sup>a</sup> log <sub>10</sub> CFU g <sup>-1</sup> tissue	<sup>b</sup> Significance (P-value)	<sup>c</sup> Reduction (log <sub>10</sub> CFU g <sup>-1</sup> )	<sup>d</sup> Surface abscesses (%)	<sup>e</sup> Number of abscesses per kidney	<sup>f</sup> Significance (P-value)
wild-type	6.141 ± 0.192	—	—	70	4.364 ± 0.889	—
ΔsrtA	4.095 ± 0.347	6.7 × 10 <sup>-6</sup>	2.046	0	0.000 ± 0.000	0.0216
spa	5.137 ± 0.374	0.0144	1.004	13	0.375 ± 0.374	0.0356

<sup>a</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 5 days following infection in cohorts of fifteen BALB/c mice per challenge strain. Standard error of the means (±SEM) is indicated.

<sup>b</sup>Statistical significance was calculated with the Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

<sup>c</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>d</sup>Abscess formation in kidney tissues five days following infection was measured by macroscopic inspection (% positive)

<sup>e</sup>Histopathology of hematoxylin-eosin stained, thin sectioned kidneys from eight to ten animals; the average number of abscesses per kidney was recorded and averaged again for the final mean (±SEM).

<sup>f</sup>Statistical significance was calculated with the Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

healthy phagocytes. A rim of necrotic PMNs were observed at the periphery of abscess lesions, bordering eosinophilic, amorphous material that separates healthy renal tissue from lesions. Abscesses eventually reached a diameter of ≥1,524 μM on day 15 or 36. At later time intervals, the staphylococcal load was increased to 10<sup>4</sup>-10<sup>6</sup> CFU g<sup>-1</sup> and growing abscess lesions migrated towards the organ capsule. Peripheral lesions were prone to rupture, thereby releasing necrotic material and staphylococci into the peritoneal cavity or the retroperitoneal space. These events resulted in bacteremia as well as a secondary wave of abscesses, eventually precipitating a lethal outcome.

To enumerate staphylococcal load in renal tissue, animals were killed, their kidneys excised and tissue homogenate spread on agar media for colony formation. On day 5 of infection, a mean of 1×10<sup>6</sup> CFU g<sup>-1</sup> renal tissue for *S. aureus* Newman was observed. To quantify abscess formation, kidneys were visually inspected, and each individual organ was given a score of one or zero. The final sum was divided by the total number of kidneys to calculate percent surface abscesses (Table 4). In addition, randomly chosen kidneys were fixed in formalin, embedded, thin sectioned, and stained with hematoxylin-eosin. For each kidney, four sagittal sections at 200 μM intervals were viewed by microscopy. The numbers of lesions were counted for each section and averaged to quantify the number of abscesses within the kidneys. *S. aureus* Newman caused 4.364±0.889 abscesses per kidney, and surface abscesses were observed on 14 out of 20 kidneys (70%) (Table 4).

When examined by scanning electron microscopy, *S. aureus* Newman was located in tightly associated lawns at the center of abscesses. Staphylococci were contained by an amorphous pseudocapsule that separated bacteria from the cuff of abscesses leukocytes. No immune cells were observed in these central nests of staphylococci, however occasional red blood cells were located among the bacteria. Bacterial

### *S. aureus* Protein A (Spa) Mutants are Avirulent and Cannot Form Abscesses

Sortase A is a transpeptidase that immobilizes nineteen surface proteins in the envelope of *S. aureus* strain Newman (Mazmanian et al., 1999; Mazmanian et al., 2000). Earlier work identified sortase A as a virulence factor in multiple animal model systems, however the contributions of this enzyme and its anchored surface proteins to abscess formation or persistence have not yet been revealed (Jonsson et al., 2002; Weiss et al., 2004). Compared to the wild-type parent (Baba et al., 2007), an isogenic srtA variant (ΔsrtA) failed to form abscess lesions on either macroscopic or histopathology examination on days 2, 5, or 15. In mice infected with the srtA mutant, only 1×10<sup>4</sup> CFU g<sup>-1</sup> was recovered from kidney tissue on day 5 of infection, which is a 2.046 log<sub>10</sub> CFU g<sup>-1</sup> reduction compared to the wild-type parent strain (P=6.73×10<sup>-6</sup>). A similar defect was observed for the srtA mutant of MRSA strain USA300 (data not shown). Scanning electron microscopy showed that srtA mutants were highly dispersed and often associated with leukocytes in otherwise healthy renal tissue. On day fifteen following infection, srtA mutants were cleared from renal tissues, a ≥3.5 log<sub>10</sub> CFU g<sup>-1</sup> reduction compared to the wild-type (Table 3). Thus, sortase A anchored surface proteins enable the formation of abscess lesions and the persistence of bacteria in host tissues, wherein staphylococci replicate as communities embedded in an extracellular matrix and shielded from surrounding leukocytes by an amorphous pseudocapsule.

Sortase A anchors a large spectrum of proteins with LPXTG motif sorting signals to the cell wall envelope, thereby providing for the surface display of many virulence factors (Mazmanian et al., 2002). To identify surface proteins required for staphylococcal abscess formation, bursa aurealis insertions were introduced in 5' coding sequences of genes that encode polypeptides with LPXTG motif proteins (Bae et al., 2004) and these mutations were transduced into *S. aureus* Newman. Mutations in the structural gene for Protein A (spa)

reduced the staphylococcal load in infected mouse kidney tissues by  $1.004 \log_{10}$  ( $P=0.0144$ ). When analyzed for their ability to form abscesses in kidney tissues by histopathology, we observed that the spa mutants were unable to form abscesses as compared with the wild-type parent strain *S. aureus* Newman (wild-type *S. aureus* Newman  $4.364 \pm 0.889$  abscesses per kidney vs. the isogenic spa mutant with  $0.375 \pm 0.374$  lesions;  $P=0.0356$ ).

#### Protein A Blocks Innate and Adaptive Immune Responses.

Studies identified Protein A as a critical virulence factor during the pathogenesis of *S. aureus* infections. Earlier work demonstrated that Protein A impedes phagocytosis of staphylococci by binding the Fc component of immunoglobulin (Jensen 1958; Uhlén et al., 1984), activates platelet aggregation via the von Willebrand factor (Hartleib et al., 2000), functions as a B cell superantigen by capturing the F(ab)<sub>2</sub> region of VH3 bearing IgM (Roben et al., 1995), and, through its activation of TNFR1, can initiate staphylococcal pneumonia (Gomez et al., 2004). Due to the fact that Protein A captures immunoglobulin and displays toxic attributes, the possibility that this surface molecule may function as a vaccine in humans has not been rigorously pursued. The inventors demonstrate for the first time that Protein A variants no longer able to bind to immunoglobulins, vWF and TNFR-1 are removed of their toxigenic potential and are able to stimulate humoral immune responses that protect against staphylococcal disease.

#### Molecular Basis of Protein A Surface Display and Function.

Protein A is synthesized as a precursor in the bacterial cytoplasm and secreted via its YSIRK signal peptide at the cross wall, i.e., the cell division septum of staphylococci (FIG. 1). (DeDent et al., 2007; DeDent et al., 2008). Following cleavage of the C-terminal LPXTG sorting signal, Protein A is anchored to bacterial peptidoglycan crossbridges by sortase A (Schneewind et al., 1995; Mazmanian et al., 1999; Mazmanian et al., 2000). Protein A is the most abundant surface protein of staphylococci; the molecule is expressed by virtually all *S. aureus* strains (Said-Salim et al., 2003; Cespedes et al., 2005; Kennedy et al., 2008). Staphylococci turn over 15-20% of their cell wall per division cycle (Navarre and Schneewind 1999). Murine hydrolases cleave the glycan strands and wall peptides of peptidoglycan, thereby releasing Protein A with its attached C-terminal cell wall disaccharide tetrapeptide into the extracellular medium (Ton-That et al., 1999). Thus, by physiological design, Protein A is both anchored to the cell wall and displayed on the bacterial surface but also released into surrounding tissues during host infection (Marraffini et al., 2006).

Protein A captures immunoglobulins on the bacterial surface and this biochemical activity enables staphylococcal escape from host innate and acquired immune responses (Jensen 1958; Goodyear and Silverman 2004). Interestingly, region X of Protein A (Guss et al., 1984), a repeat domain that tethers the IgG binding domains to the LPXTG sorting signal/cell wall anchor, is perhaps the most variable portion of the staphylococcal genome (Schneewind et al., 1992; Said-Salim et al., 2003). Each of the five immunoglobulin binding domains of Protein A (SpA), formed from three helix bundles and designated E, D, A, B, and C, exerts similar structural and functional properties (Sjödahl 1977; Jansson et al., 1998). The solution and crystal structure of domain D has been solved both with and without the Fc and V<sub>H3</sub> (Fab) ligands, which bind Protein A in a non-competitive manner at distinct sites (Graille et al., 2000).

In the crystal structure complex, the Fab interacts with helix II and helix III of domain D via a surface composed of

four VH region  $\beta$ -strands (Graille et al., 2000). The major axis of helix II of domain D is approximately  $50^\circ$  to the orientation of the strands, and the interhelical portion of domain D is most proximal to the C0 strand. The site of interaction on Fab is remote from the Ig light chain and the heavy chain constant region. The interaction involves the following domain D residues: Asp-36 of helix II as well as Asp-37 and Gln-40 in the loop between helix II and helix III, in addition to several other residues with SpA-D (Graille et al., 2000). Both interacting surfaces are composed predominantly of polar side chains, with three negatively charged residues on domain D and two positively charged residues on the 2A2 Fab buried by the interaction, providing an overall electrostatic attraction between the two molecules. Of the five polar interactions identified between Fab and domain D, three are between side chains. A salt bridge is formed between Arg-H19 and Asp-36 and two hydrogen bonds are made between Tyr-H59 and Asp-37 and between Asn-H82a and Ser-33. Because of the conservation of Asp-36 and Asp-37 in all five IgG binding domains of Protein A, these residues were selected for mutagenesis.

The SpA-D sites responsible for Fab binding are structurally separate from the domain surface that mediates Fc $\gamma$  binding. The interaction of Fc $\gamma$  with domain B primarily involves residues in helix I with lesser involvement of helix II (Deisenhofer 1981; Gouda et al., 1992). With the exception of the Gln-32, a minor contact in both complexes, none of the residues that mediate the Fc $\gamma$  interaction are involved in Fab binding. To examine the spatial relationship between these different Ig-binding sites, the SpA domains in these complexes have been superimposed to construct a model of a complex between Fab, the SpA-domain D, and the Fc $\gamma$  molecule. In this ternary model, Fab and Fc $\gamma$  form a sandwich about opposite faces of the helix II without evidence of steric hindrance of either interaction. These findings illustrate how, despite its small size (i.e., 56-61 aa), a SpA domain can simultaneously display both activities, explaining experimental evidence that the interactions of Fab with an individual domain are noncompetitive. Residues for the interaction between SpA-D and Fc $\gamma$  are Gln-9 and Gln-10.

In contrast, occupancy of the Fc portion of IgG on the domain D blocks its interaction with vWF A1 and probably also TNFR1 (O'Seaghda et al., 2006). Mutations in residues essential for IgG Fc binding (F5, Q9, Q10, S11, F13, Y14, L17, N28, I31 and K35) are also required for vWF A1 and TNFR1 binding (Cedergren et al., 1993; Gomez et al., 2006; O'Seaghda et al. 2006), whereas residues critical for the V<sub>H3</sub> interaction (Q26, G29, F30, S33, D36, D37, Q40, N43, E47) have no impact on the binding activities of IgG Fc, vWF A1 or TNFR1 (Jansson et al., 1998; Graille et al., 2000). The Protein A immunoglobulin Fab binding activity targets a subset of B cells that express VH3 family related IgM on their surface, i.e. these molecules function as VH3 type B cell receptors (Roben et al., 1995). Upon interaction with SpA, these B cells rapidly proliferate and then commit to apoptosis, leading to preferential and prolonged deletion of innate-like B lymphocytes (i.e. marginal zone B cells and follicular B2 cells) (Goodyear and Silverman 2003; Goodyear and Silverman 2004). It is important to note that more than 40% of circulating B cells are targeted by the Protein A interaction and the VH3 family represents the largest family of human B cell receptors to impart protective humoral responses against pathogens (Goodyear and Silverman 2003; Goodyear and Silverman 2004). Thus, Protein A functions analogously to staphylococcal superantigens (Roben et al., 1995), albeit that the latter class of molecules, for example SEB, TSST-1, TSST-2, form complexes with the T cell receptor to inappro-

priately stimulate host immune responses and thereby precipitating characteristic disease features of staphylococcal infections (Roben et al., 1995; Tiedemann et al., 1995). Together these findings document the contributions of Protein A in establishing staphylococcal infections and in modulating host immune responses.

#### Non-Toxicogenic Variant of Protein A.

The inventors have developed a non-toxicogenic variant of staphylococcal Protein A and, with this reagent in hand, aimed for the first time to measure the immune response of animals to Protein A immunization. Further, the inventors address whether immunization of animals with a non-toxicogenic variant of Protein A could generate immune responses that raise protective immunity against staphylococcal infection.

To perturb the IgG Fc, vWF A1 and TNFR1 binding activities of Protein A, glutamine (Q) residues 9 and 10 [the numbering here is derived from that established for the SpA domain D] were modified generating lysine or glycine substitutions for both glutamines with the expectation that these substitutions abolish the ion bonds formed between wild-type Protein A and its ligands. The added effect of the dual lysine substitutions may be that these positively charged residues institute a repellent charge for immunoglobulins. To perturb IgM Fab VH3 binding, the inventors selected the aspartate (D) residues 36 and 37 of SpA-D, each of which is required for the association of Protein A with the B cell receptor. D36 and D37 were both substituted with alanine. The Q9,10K and D36,37A mutations were combined in the recombinant molecule SpA-D<sub>Q9,10K;D36,37A</sub> and examined for the binding attributes of Protein A.

In brief, the Protein A (spa) genomic sequence of *Staphylococcus aureus* N315 was PCR amplified with the primers (GCTGCACATATGGCGCAACACGATGAAGCTCAAC [5' primer] (SEQ ID NO:35) and AGTGGATCCTTATGCTTTGTTAGCATCTGC [3' primer] (SEQ ID NO:36)), cloned into the pET15b vector (pYSJ1, codons 48-486) (Stranger-Jones, et al., 2006) and recombinant plasmid transformed into *E. coli* BL21(DE3) (Studier et al., 1990). The Protein A product derived from pYSJ1 harbors SpA residues 36-265 fused to the N-terminal His tag (MGSSHHHHHHHSSGLVPRGS (SEQ ID NO:37)). Following IPTG inducible expression, recombinant N-terminal His<sub>6</sub>-tagged SpA was purified by affinity chromatography on Ni-NTA resin (Stranger-Jones et al., 2006). The domain D of SpA (SpA-D) was PCR amplified with a pair of specific primers (AATCATATGTTCAACAAAGATCAACAAAGC [5' primer] (SEQ ID NO:38) and AAGGATCCAGATTCGTTTAAATTTTAGC [3' primer] (SEQ ID NO:39)), sub-cloned into the pET15b vector (pHAN1, spa codons 212-261) and recombinant plasmid transformed into *E. coli* BL21(DE3) to express and purify recombinant N-terminal His<sub>6</sub>-tagged protein. To generate mutations in the SpA-D coding sequence, sets of two pairs of primers were synthesized (for D to A substitutions: CTTTCATTCAAAGTCTTAAAGCCGCCCAAGCCAAAGACTACTAAC [5' primer] (SEQ ID NO:40) and GTTAGTGCTTTGGCTTGGGGCGGCTTTAAGACTTTGAATGAAG [3' primer] (SEQ ID NO:41); for Q to K substitutions CATATGTTCAACAAAGATAAAAAAAGCGCCTTCTATGAAATC [5' primer] (SEQ ID NO:42) and GATTTCATAGAAGGCGCTTTTTTATCTTTGTTGAACATATG [3' primer] (SEQ ID NO:43); for Q to G substitutions CATATGTTCAACAAAGATGAGGAAGCGCCTTCTATGAAATC [5' primer] (SEQ ID NO:44) and GATTTCATAGAAGGCGCTTCTCCTCATCTTTGTTGAACATATG [3' primer] (SEQ ID NO:45). Primers were used for quick-change mutagenesis protocols.

Following mutagenesis, DNA sequences were confirmed for each of the recombinant proteins: SpA, SpA-D and SpA-D<sub>Q9,10K;D36,37A</sub> and SpA-D<sub>Q9,10K;D36,37A</sub>. All proteins were purified from lysates of recombinant *E. coli* using Ni-NTA chromatography and subsequently dialyzed against PBS and stored at 4° C.

To measure binding of immunoglobulin to Protein A and its variants, 200 µg of purified protein was diluted into a 1 ml volume using column buffer (50 mM Tris-HCl, 150 mM NaCl, pH7.5) and then loaded onto a pre-equilibrated Ni-NTA column (1 ml bed volume). Columns were washed with 10 ml of column buffer. 200 µg of purified human IgG was diluted in a total volume of 1 ml column buffer and then applied to each of the columns charged with Protein A and its variants. The columns were subsequently washed with 5 ml wash buffer (10 mM imidazole in column buffer) and 5 ml column buffer. Protein samples were eluted with 2 ml elution buffer (500 mM imidazole in column buffer), fractions collected and aliquots subjected to SDS-PAGE gel electrophoresis, followed by Coomassie-Blue staining. As shown in FIG. 3, wild-type Protein A (SpA) and its SpA-domain D both retained immunoglobulin during chromatography. In contrast, the SpA-D<sub>Q9,10K;D36,37A</sub> variant did not bind to immunoglobulin.

To quantify the binding of Protein A and its variants to the Fc portion of immunoglobulin and the VH3 domain of Fab, HRP conjugated human immunoglobulin G [hIgG], the Fc portion of human IgG [hFc] and the F(ab)<sub>2</sub> portion of human IgG [hF(ab)<sub>2</sub>] as well as ELISA assays were used to quantify the relative amount binding to Protein A and its variants. The data in FIG. 4 demonstrate the binding of SpA and SpA-D to hIgG and hFc, whereas SpA-D<sub>Q9,10K;D36,37A</sub> and SpA-D<sub>Q9,10K;D36,37A</sub> displayed only background binding activities. SpA bound similar amounts of hFc and hF(ab)<sub>2</sub>, however the binding of SpA-D to hF(ab)<sub>2</sub> was reduced compared to full length SpA. This result suggests that the presence of multiple IgG binding domains may cooperatively increase the ability of Protein A to bind to the B cell receptor. When compared with the reduced binding power of SpA-D for hF(ab)<sub>2</sub>, of the two variants only SpA-D<sub>Q9,10K;D36,37A</sub> displayed a significant reduction in the ability to bind the VH3 domain of immunoglobulin. To examine the toxicogenic attributes of SpA-D and its variants, purified proteins were injected into mice, which were sacrificed after 4 hours to remove their spleens. Organ tissue was homogenized, capsular material removed and B cells stained with fluorescent CD19 antibodies. Following FACS analysis to quantify the abundance of B cells in splenic tissues, it was observed that SpA-D caused a 5% drop in the B cell count compared to a mock (PBS) control (FIG. 5). In contrast, SpA-D<sub>Q9,10K;D36,37A</sub> did not cause a reduction in B-cell counts, indicating that the mutant molecule had lost its toxicogenic attributes of stimulating B cell proliferation and death (FIG. 5). In summary, amino acid substitutions in the SpA-D residues Q9, Q10, D36, and D37 abolished the ability of Protein A domains to bind immunoglobulins or exert toxicogenic functions in human and animal tissues.

#### Non-Toxicogenic Protein A Variants Elicit Vaccine Protection.

To test whether or not Protein A and its variants can function as vaccine antigens, SpA, SpA-D, SpA-D<sub>Q9,10K;D36,37A</sub>, and SpA-D<sub>Q9,10K;D36,37A</sub> were emulsified with complete or incomplete Freund's adjuvant and immunized 4 week old BALB/c mice on day 1 and day 11 with 50 µg of purified protein. Cohort of animals (n=5) were analyzed for humoral immune responses to immunization by bleeding the animals before (day 0) and after the immunization schedule (day 21). Table 5 indicates that immunized mice generated only a mod-

est humoral immune response directed at wild-type Protein A or its SpA-D module, whereas the amount of antibody raised following immunization with SpA-D<sub>Q9,10K;D36,37A</sub> or SpA-D<sub>Q9,10G;D36,37A</sub> was increased four to five fold. Following intravenous challenge with  $1 \times 10^7$  CFU *S. aureus* Newman, animals were killed on day 4, their kidneys removed and

immune responses as is reported here for the SpA-domain D alone. Further, a likely attribute of Protein A specific antibodies may be that the interaction of antigen binding sites with the microbial surface can neutralize the ability of staphylococci to capture immunoglobulins via their Fc portion or to stimulate the B cell receptor via the VH3 binding activities.

TABLE 5

Antigen	Bacterial load in kidney (n = number of mice)			IgG titer	Abscess formation in mice (n = number of mice)				
	<sup>a</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>b</sup> Reduction	<sup>c</sup> P value		<sup>d</sup> Surface abscess	Reduction	<sup>e</sup> Histopathology	Reduction	<sup>f</sup> p value
Mock	6.46 ± 0.25 (n = 19)	—	—	<100	14/19 (70%)	—	3.7 ± 1.2 (n = 10)	—	—
SpA	3.95 ± 0.56 (n = 20)	2.51	0.0003	1706 ± 370	10/20 (50%)	32%	2.1 ± 1.2 (n = 10)	2.2	0.35
SpA-D	4.43 ± 0.41 (n = 18)	2.03	0.0001	381 ± 27	10/18 (55%)	25%	1.5 ± 0.8 (n = 10)	2.2	0.15
SpA-D1	3.39 ± 0.50 (n = 19)	3.07	<0.0001	5600 ± 801	6/20 (30%)	59%	0.5 ± 0.4 (n = 10)	3.2	0.02
SpA-D2	3.43 ± 0.46 (n = 19)	3.03	<0.0001	3980 ± 676	6/19 (32%)	57%	0.8 ± 0.5 (n = 10)	2.9	0.04

<sup>a</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 4 days following infection in cohorts of 18 to 20 BALB/c mice. Standard error of the means (±SEM) is indicated.

<sup>c</sup>Statistical significance was calculated with the Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

<sup>b</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>d</sup>Abscess formation in kidney tissues four days following infection was measured by macroscopic inspection (% positive)

<sup>e</sup>Histopathology of hematoxylin-eosin stained, thin sectioned kidneys from ten animals; the number of abscesses per kidney was recorded and averaged for the final mean (±SEM).

<sup>f</sup>Statistical significance was calculated with the Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

SpA-D1 and SpA-D2 represent SpA-D<sub>Q9,10K;D36,37A</sub> and SpA-D<sub>Q9,10G;D36,37A</sub>, respectively.

either analyzed for staphylococcal load (by plating tissue homogenate on agar plates and enumerating colony forming units, CFU) or histopathology. As expected, mock (PBS) immunized mice (n=19) harbored 6.46 log<sub>10</sub> (±0.25) CFU in kidney tissue and infectious lesions were organized into 3.7 (±1.2) abscesses per organ (n=10) (Table 5). Immunization of animals with SpA led to a 2.51 log<sub>10</sub> CFU reduction on day 5 (P=0.0003) with 2.1 (±1.2) abscesses per organ. The latter data indicate that there was no significant reduction in abscess formation (P=0.35). Immunization with SpA-D generated similar results: a 2.03 log<sub>10</sub> CFU reduction on day 5 (P=0.0001) with 1.5 (±0.8) abscesses per organ (P=0.15). In contrast, immunization with SpA-D<sub>Q9,10K;D36,37A</sub> or SpA-D<sub>Q9,10G;D36,37A</sub> created increased protection, with 3.07 log<sub>10</sub> and 3.03 log<sub>10</sub> CFU reduction on day 4, respectively (statistical significance P<0.0001 for both observations). Further, immunization with both SpA-D<sub>Q9,10K;D36,37A</sub> and SpA-D<sub>Q9,10G;D36,37A</sub> generated significant protection from staphylococcal abscess formation, as only 0.5 (±0.4) and 0.8 (±0.5) infectious lesions per organ (P=0.02 and P=0.04) were identified. Thus, immunization with non-toxicogenic Protein A variants generates increased humoral immune responses for Protein A and provides protective immunity against staphylococcal challenge. These data indicate that Protein A is an ideal candidate for a human vaccine that prevents *S. aureus* disease.

These exciting results have several implications for the design of a human vaccine. First, the generation of substitution mutations that affect the ability of the immunoglobulin binding domains of Protein A, either alone or in combination of two or more domains, can generate non-toxicogenic variants suitable for vaccine development. It seems likely that a combination of mutant IgG binding domains closely resembling the structure of Protein A can generate even better humoral

Vaccine Protection in Murine Abscess, Murine Lethal Infection, and Murine Pneumonia Models.

Three animal models have been established for the study of *S. aureus* infectious disease. These models are used here to examine the level of protective immunity provided via the generation of Protein A specific antibodies.

Murine Abscess—

BALB/c mice (24-day-old female, 8-10 mice per group, Charles River Laboratories, Wilmington, Mass.) are immunized by intramuscular injection into the hind leg with purified protein (Chang et al., 2003; Schneewind et al., 1992). Purified SpA, SpA-D or SpA-DQ9,10K;D36,37A (50 µg protein) is administered on days 0 (emulsified 1:1 with complete Freund's adjuvant) and 11 (emulsified 1:1 with incomplete Freund's adjuvant). Blood samples are drawn by retroorbital bleeding on days 0, 11, and 20. Sera are examined by ELISA for IgG titers for specific SpA-D and SpA-DQ9,10K;D36,37A binding activity. Immunized animals are challenged on day 21 by retroorbital injection of 100 µl of *S. aureus* Newman or *S. aureus* USA300 suspension ( $1 \times 10^7$  cfu). For this, overnight cultures of *S. aureus* Newman are diluted 1:100 into fresh tryptic soy broth and grown for 3 h at 37° C. Staphylococci are centrifuged, washed twice, and diluted in PBS to yield an A<sub>600</sub> of 0.4 ( $1 \times 10^8$  cfu per ml). Dilutions are verified experimentally by agar plating and colony formation. Mice are anesthetized by intraperitoneal injection of 80-120 mg of ketamine and 3-6 mg of xylazine per kilogram of body weight and infected by retroorbital injection. On day 5 or 15 following challenge, mice are euthanized by compressed CO<sub>2</sub> inhalation. Kidneys are removed and homogenized in 1% Triton X-100. Aliquots are diluted and plated on agar medium for triplicate determination of cfu. For histology, kidney tissue is incubated at room temperature in 10% formalin for 24 h. Tissues are embedded in paraffin, thin-sectioned, stained with hematoxylin/leusin, and examined by microscopy.

## Murine Lethal Infection—

BALB/c mice (24-day-old female, 8-10 mice per group, Charles River Laboratories, Wilmington, Mass.) are immunized by intramuscular injection into the hind leg with purified SpA, SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> (50 µg protein). Vaccine is administered on days 0 (emulsified 1:1 with complete Freund's adjuvant) and 11 (emulsified 1:1 with incomplete Freund's adjuvant). Blood samples are drawn by retro-orbital bleeding on days 0, 11, and 20. Sera are examined by ELISA for IgG titers with specific SpA-D and SpA-D<sub>Q9,10K;D36,37A</sub> binding activity. Immunized animals are challenged on day 21 by retro-orbital injection of 100 µl of *S. aureus* Newman or *S. aureus* USA300 suspension (15×10<sup>7</sup> cfu) (34). For this, overnight cultures of *S. aureus* Newman are diluted 1:100 into fresh tryptic soy broth and grown for 3 h at 37° C. Staphylococci are centrifuged, washed twice, diluted in PBS to yield an A<sub>600</sub> of 0.4 (1×10<sup>8</sup> cfu per ml) and concentrated. Dilutions are verified experimentally by agar plating and colony formation. Mice are anesthetized by intraperitoneal injection of 80-120 mg of ketamine and 3-6 mg of xylazine per kilogram of body weight. Immunized animals are challenged on day 21 by intraperitoneal inject with 2×10<sup>10</sup> cfu of *S. aureus* Newman or 3-10×10<sup>9</sup> cfu of clinical *S. aureus* isolates. Animals are monitored for 14 days, and lethal disease is recorded.

## Murine Pneumonia Model—

*S. aureus* strains Newman or USA300 (LAC) are grown at 37° C. in tryptic soy broth/agar to OD<sub>660</sub> 0.5. 50-ml culture aliquots are centrifuged, washed in PBS, and suspended in 750 µl PBS for mortality studies (3-4×10<sup>8</sup> CFU per 30-µl volume), or 1,250 µl PBS (2×10<sup>8</sup> CFU per 30 µl volume) for bacterial load and histopathology experiments (2, 3). For lung infection, 7-wk-old C57BL/6J mice (The Jackson Laboratory) are anesthetized before inoculation of 30 µl of *S. aureus* suspension into the left nare. Animals are placed into the cage in a supine position for recovery and observed for 14 days. For active immunization, 4-wk-old mice receive 20 µg SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> in CFA on day 0 via the i.m. route, followed by a boost with 20 µg SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> in incomplete Freund's adjuvant (IFA) on day 10. Animals are challenged with *S. aureus* on day 21. Sera are collected before immunization and on day 20 to assess specific antibody production. For passive immunization studies, 7-wk-old mice receive 100 µl of either NRS (normal rabbit serum) or SpA-D-specific rabbit antisera via i.p. injection 24 h before challenge. To assess the pathological correlates of pneumonia, infected animals are killed via forced CO<sub>2</sub> inhalation before removal of both lungs. The right lung is homogenized for enumeration of lung bacterial load. The left lung is placed in 1% formalin and paraffin embedded, thin sectioned, stained with hematoxylin-eosin, and analyzed by microscopy.

## Rabbit Antibodies—

Purified 200 µg SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> is used as an immunogen for the production of rabbit antisera. 200 µg protein is emulsified with CFA for injection at day 0, followed by booster injections with 200 µg protein emulsified with IFA on days 21 and 42. Rabbit antibody titers are determined by ELISA. Purified antibodies are obtained by affinity chromatography of rabbit serum on SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> sepharose. The concentration of eluted antibodies is measured by absorbance at A<sub>280</sub> and specific antibody titers are determined by ELISA.

## Active Immunization with SpA-Domain D Variants.—

To determine vaccine efficacy, animals are actively immunized with purified SpA-D or SpAD<sub>Q9,10K;D36,37A</sub>. As a control, animals are immunized with adjuvant alone. Antibody

titers against Protein A preparations are determined using SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> as antigens; note that the SpA-D<sub>Q9,10K;D36,37A</sub> variant cannot bind the Fc or Fab portion of IgG. Using infectious disease models described above, any reduction in bacterial load (murine abscess and pneumonia), histopathology evidence of staphylococcal disease (murine abscess and pneumonia) and protection from lethal disease (murine lethal challenge and pneumonia) is measured.

Passive Immunization with Affinity Purified Rabbit Polyclonal Antibodies Generated Against SpA-Domain D Variants.

To determine protective immunity of Protein A specific rabbit antibodies, mice are passively immunized with 5 mg/kg of purified SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub> derived rabbit antibodies. Both of these antibody preparations are purified by affinity chromatography using immobilized SpA-D or SpA-D<sub>Q9,10K;D36,37A</sub>. As a control, animals are passively immunized with rV10 antibodies (a plague protective antigen that has no impact on the outcome of staphylococcal infections). Antibody titers against all Protein A preparations are determined using SpA-D<sub>Q9,10K;D36,37A</sub> as an antigen, as this variant cannot bind the Fc or Fab portion of IgG. Using the infectious disease models described above, the reduction in bacterial load (murine abscess and pneumonia), histopathology evidence of staphylococcal disease (murine abscess and pneumonia), and the protection from lethal disease (murine lethal challenge and pneumonia) is measured.

## Example 2

Non-Toxicogenic Protein a Vaccine for Methicillin-Resistant *Staphylococcus aureus* Infection

Clinical isolates of *S. aureus* express protein A (Shopsin et al., 1999, whose primary translational product is comprised of an N-terminal signal peptide (DeDent et al., 2008), five Ig-BDs (designated E, D, A, B and C) (Sjodahl, 1977), region X with variable repeats of an eight residue peptide (Guss et al., 1984), and C-terminal sorting signal for the cell wall anchoring of SpA (Schneewind et al., 1992; Schneewind et al., 1995) (FIG. 6). Guided by amino acid homology (Uhlen et al., 1984), the triple α-helical bundle structure of IgBDs (Deisenhofer et al., 1978; Deisenhofer et al., 1981) and their atomic interactions with Fab V<sub>H</sub>3 (Graille et al., 2000) or Fcγ (Gouda et al., 1998), glutamine 9 and 10 were selected as well as aspartate 36 and 37 as critical for the association of SpA with antibodies or B cell receptor, respectively. Substitutions Gln9Lys, Gln10Lys, Asp36Ala and Asp37Ala were introduced into the D domain to generate SpA-D<sub>KKAA</sub> (FIG. 6). The ability of isolated SpA-D or SpA-D<sub>KKAA</sub> to bind human IgG was analyzed by affinity chromatography (FIG. 6). Polystyrene tagged SpA-D as well as full-length SpA retained human IgG on Ni-NTA, whereas SpA-D<sub>KKAA</sub> and a negative control (SrtA) did not (FIG. 6). A similar result was observed with von Willebrand factor (Hartleib et al., 2000), which, along with tumor necrosis factor receptor 1 (TNFR1) (Gomez et al., 2004), can also bind protein A via glutamine 9 and 10 (FIG. 6). Human immunoglobulin encompasses 60-70% V<sub>H</sub>3-type IgG. The inventors distinguish between Fc domain and B cell receptor activation of Igs and measured association of human Fcγ and F(ab)<sub>2</sub> fragments, both of which bound to full-length SpA or SpA-D, but not to SpA-D<sub>KKAA</sub> (FIG. 6). Injection of SpA-D into the peritoneal cavity of mice resulted in B cell expansion followed by apoptotic collapse of CD19+ lymphocytes in spleen tissue of BALB/c mice (Goodyear and Silverman, 2003) (FIG. 6). B cell superantigen activity was

not observed following injection with SpA-D<sub>KKAA</sub>, and TUNEL-staining of splenic tissue failed to detect the increase in apoptotic cells that follows injection of SpA or SpA-D (FIG. 6).

Naive six week old BALB/c mice were injected with 50 µg each of purified SpA, SpA-D or SpA-D<sub>KKAA</sub> emulsified in CFA and boosted with the same antigen emulsified in IFA. In agreement with the hypothesis that SpA-D promotes the apoptotic collapse of activated clonal B cell populations, the inventors observed a ten-fold higher titer of SpA-D<sub>KKAA</sub> specific antibodies following immunization of mice with the non-toxic variant as compared to the B cell superantigen (SpA-D vs. SpA-D<sub>KKAA</sub> P<0.0001, Table 6). Antibody titers raised by immunization with full-length SpA were higher than those elicited by SpA-D (P=0.0022), which is likely due to the larger size and reiterative domain structure of this antigen (Table 6). Nevertheless, even SpA elicited lower antibody titers than SpA-D<sub>KKAA</sub> (P=0.0003), which encompasses only 50 amino acids of protein A (520 residues, SEQ ID NO:33). Immunized mice were challenged by intravenous inoculation with *S. aureus* Newman and the ability of staphylococci to seed abscesses in renal tissues was examined by necropsy four days after challenge. In homogenized renal tissue of mock (PBS/adjuvant) immunized mice, an average staphylococcal load of 6.46 log<sub>10</sub> CFU g<sup>-1</sup> was enumerated (Table 6). Immunization of mice with SpA or SpA-D led to a reduction in staphylococcal load, however SpA-D<sub>KKAA</sub> vaccinated animals displayed an even greater, 3.07 log<sub>10</sub> CFU g<sup>-1</sup> reduction of *S. aureus* Newman in renal tissues (P<0.0001, Table 6). Abscess formation in kidneys was analyzed by histopathology (FIG. 7). Mock immunized animals harbored an average of 3.7 (±1.2) abscesses per kidney (Table 6). Vaccination with SpA-D<sub>KKAA</sub> reduced the average number of abscesses to 0.5 (±0.4) (P=0.0204), whereas immunization with SpA or SpA-D did not cause a significant reduction in the number of abscess lesions (Table 6). Lesions from SpA-D<sub>KKAA</sub> vaccinated animals were smaller in size, with fewer infiltrating PMNs and characteristically lacked staphylococcal abscess communities (Cheng et al., 2009) (FIG. 7). Abscesses in animals that had been immunized with SpA or SpA-D displayed the same overall structure of lesions in mock immunized animals (FIG. 7).

The inventors examined whether SpA-D<sub>KKAA</sub> immunization can protect mice against MRSA strains and selected the USA300 LAC isolate for animal challenge (Diep et al., 2006). This highly virulent CA-MRSA strain spread rapidly throughout the United States, causing significant human morbidity and mortality (Kennedy et al., 2008). Compared to adjuvant control mice, SpA-D<sub>KKAA</sub> immunized animals harbored a 1.07 log<sub>10</sub> CFU g<sup>-1</sup> reduction in bacterial load of

infected kidney tissues. Histopathology examination of renal tissue following *S. aureus* USA300 challenge revealed that the average number of abscesses was reduced from 4.04 (±0.8) to 1.6 (±0.6) (P=0.02774). In contrast, SpA or SpA-D immunization did not cause a significant reduction in bacterial load or abscess formation (Table 6).

Rabbits were immunized with SpA-D<sub>KKAA</sub> and specific antibodies were purified on SpA-D<sub>KKAA</sub> affinity column followed by SDS-PAGE (FIG. 8). SpA-D<sub>KKAA</sub> specific IgG was cleaved with pepsin to generate Fcγ and F(ab)<sub>2</sub> fragments, the latter of which were purified by chromatography on SpA-D<sub>KKAA</sub> column (FIG. 8). Binding of human IgG or vWF to SpA or SpA-D was perturbed by SpA-D<sub>KKAA</sub> specific F(ab)<sub>2</sub>, indicating that SpA-D<sub>KKAA</sub> derived antibodies neutralize the B cell superantigen function of protein A as well as its interactions with Ig (FIG. 8).

To further improve the vaccine properties for non-toxic protein A, the inventors generated SpA<sub>KKAA</sub>, which includes all five IgBDs with four amino acid substitutions—substitutions corresponding to Gln9Lys, Gln10Lys, Asp36Ala and Asp37Ala of domain D—in each of its five domains (E, D, A, B and C). Polyhistidine tagged SpA<sub>KKAA</sub> was purified by affinity chromatography and analyzed by Coomassie Blue-stained SDS-PAGE (FIG. 9). Unlike full-length SpA, SpA<sub>KKAA</sub> did not bind human IgG, Fc and F(ab)<sub>2</sub> or vWF (FIG. 9). SpA<sub>KKAA</sub> failed to display B cell superantigen activity, as injection of the variant into BALB/c mice did not cause a depletion of CD19+ B cells in splenic tissue (FIG. 9). SpA<sub>KKAA</sub> vaccination generated higher specific antibody titers than SpA-D<sub>KKAA</sub> immunization and provided mice with elevated protection against *S. aureus* USA300 challenge (Table 6). Four days following challenge, SpA<sub>KKAA</sub> vaccinated animals harbored 3.54 log<sub>10</sub> CFU g<sup>-1</sup> fewer staphylococci in renal tissues (P=0.0001) and also caused a greater reduction in the number of abscess lesions (P=0.0109) (Table 6).

SpA<sub>KKAA</sub> was used to immunize rabbits. Rabbit antibodies specific for SpA-D<sub>KKAA</sub> or SpA<sub>KKAA</sub> were affinity purified on matrices with immobilized cognate antigen and injected at a concentration of 5 mg kg<sup>-1</sup> body weight into the peritoneal cavity of BALB/c mice (Table 7). Twenty-four hours later, specific antibody titers were determined in serum and animals challenged by intravenous inoculation with *S. aureus* Newman. Passive transfer reduced the staphylococcal load in kidney tissues for SpA-D<sub>KKAA</sub> (P=0.0016) or SpA<sub>KKAA</sub> (P=0.0005) specific antibodies. On histopathology examination, both antibodies reduced the abundance of abscess lesions in the kidneys of mice challenged with *S. aureus* Newman (Table 7). Together these data reveal that vaccine protection following immunization with SpA-D<sub>KKAA</sub> or SpA<sub>KKAA</sub> is conferred by antibodies that neutralize protein A.

TABLE 6

Immunization of mice with protein A vaccines.						
Staphylococcal load and abscess formation in renal tissue						
Antigen	<sup>a</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>b</sup> P-value	<sup>c</sup> Reduction (log <sub>10</sub> CFU g <sup>-1</sup> )	<sup>d</sup> IgG Titer	<sup>e</sup> Number of abscesses	<sup>f</sup> P-value
<i>S. aureus</i> Newman challenge						
Mock	6.46 ± 0.25	—	—	<100	3.7 ± 1.2	—
SpA	3.95 ± 0.56	0.0003	2.51	1706 ± 370	2.1 ± 1.2	0.3581
SpA-D	4.43 ± 0.41	0.0001	2.03	381 ± 27	1.5 ± 0.8	0.1480
SpA-D <sub>KKAA</sub>	3.39 ± 0.50	<0.0001	3.07	5600 ± 801	0.5 ± 0.4	0.0204

TABLE 6-continued

Immunization of mice with protein A vaccines.						
Staphylococcal load and abscess formation in renal tissue						
Antigen	<sup>a</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>b</sup> P-value	<sup>c</sup> Reduction (log <sub>10</sub> CFU g <sup>-1</sup> )	<sup>d</sup> IgG Titer	<sup>e</sup> Number of abscesses	<sup>f</sup> P-value
<i>S. aureus</i> USA300 (LAC) challenge						
Mock	7.20 ± 0.24	—	—	<100	4.0 ± 0.8	—
SpA	6.81 ± 0.26	0.2819	0.39	476 ± 60	3.3 ± 1.0	0.5969
SpA-D	6.34 ± 0.52	0.1249	0.86	358 ± 19	2.2 ± 0.6	0.0912
SpA-D <sub>KKAA</sub>	6.00 ± 0.42	0.0189	1.20	3710 ± 1147	1.6 ± 0.6	0.0277
SpA <sub>KKAA</sub>	3.66 ± 0.76	0.0001	3.54	10200 ± 2476	1.2 ± 0.5	0.0109

<sup>a</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 4 days following infection in cohorts of fifteen to twenty BALB/c mice per immunization. Representative of two independent and reproducible animal experiments is shown. Standard error of the means (±SEM) is indicated.

<sup>b</sup>Statistical significance was calculated with the unpaired two-tailed Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

<sup>c</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>d</sup>Means of five randomly chosen serum IgG titers were measured prior to staphylococcal infection by ELISA.

<sup>e</sup>Histopathology of hematoxyline-eosin stained, thin sectioned kidneys from ten animals; the average number of abscesses per kidney was recorded and averaged again for the final mean (±SEM).

TABLE 7

Passive immunization of mice with antibodies against protein A.						
Staphylococcal load and abscess formation in renal tissue						
<sup>a</sup> Antibody	<sup>b</sup> log <sub>10</sub> CFU g <sup>-1</sup>	<sup>c</sup> P-value	<sup>d</sup> Reduction (log <sub>10</sub> CFU g <sup>-1</sup> )	<sup>e</sup> IgG Titer	<sup>f</sup> Number of abscesses	<sup>g</sup> P-value
Mock	7.10 ± 0.14	—	—	<100	4.5 ± 0.8	—
α-SpA-D <sub>KKAA</sub>	5.53 ± 0.43	0.0016	1.57	466 ± 114	1.9 ± 0.7	0.0235
α-SpA <sub>KKAA</sub>	5.69 ± 0.34	0.0005	1.41	1575 ± 152	1.6 ± 0.5	0.0062

<sup>a</sup>Affinity purified antibodies were injected into the peritoneal cavity of BALB/c mice at a concentration of 5 mg · kg<sup>-1</sup> twenty-four hours prior to intravenous challenge with 1 × 10<sup>7</sup> CFU *S. aureus* Newman.

<sup>b</sup>Means of staphylococcal load calculated as log<sub>10</sub> CFU g<sup>-1</sup> in homogenized renal tissues 4 days following infection in cohorts of fifteen BALB/c mice per immunization. Representative of two independent and reproducible animal experiments is shown. Standard error of the means (±SEM) is indicated.

<sup>c</sup>Statistical significance was calculated with the unpaired two-tailed Students t-test and P-values recorded; P-values < 0.05 were deemed significant.

<sup>d</sup>Reduction in bacterial load calculated as log<sub>10</sub> CFU g<sup>-1</sup>.

<sup>e</sup>Means of five randomly chosen serum IgG titers were measured prior to staphylococcal infection by ELISA.

<sup>f</sup>Histopathology of hematoxyline-eosin stained, thin sectioned kidneys from ten animals; the average number of abscesses per kidney was recorded and averaged again for the final mean (±SEM).

40

Following infection with virulent *S. aureus*, mice do not develop protective immunity against subsequent infection with the same strain (Burts et al., 2008) (FIG. 10). The average abundance of SpA-D<sub>KKAA</sub> specific IgG in these animals was determined by dot blot as 0.20 μg ml<sup>-1</sup> (±0.04) and 0.14 μg ml<sup>-1</sup> (±0.01) for strains Newman and USA300 LAC, respectively (FIG. 9). The minimal concentration of protein A-specific IgG required for disease protection in SpA<sub>KKAA</sub> or SpA-D<sub>KKAA</sub> vaccinated animals (P .0.05 log<sub>10</sub> reduction in staphylococcal CFU g<sup>-1</sup> renal tissue) was calculated as 4.05 μg ml<sup>-1</sup> (±0.88). Average serum concentration of SpA-specific IgG in adult healthy human volunteers (n=16) was 0.21 μg ml<sup>-1</sup> (±0.02). Thus, *S. aureus* infections in mice or humans are not associated with immune responses that raise significant levels of neutralizing antibodies directed against protein A, which is likely due to the B cell superantigen attributes of this molecule. In contrast, the average serum concentration of IgG specific for diphtheria toxin in human volunteers, 0.068 μg ml<sup>-1</sup> (±0.20), was within range for protective immunity against diphtheria (Behring, 1890; Lagergard et al., 1992).

Clinical *S. aureus* isolates express protein A, an essential virulence factor whose B cell superantigen activity and evasive attributes towards opsono-phagocytic clearance are absolutely required for staphylococcal abscess formation (Palmqvist et al., 2005; Cheng et al., 2009; Silverman and Goodyear, 2006). Protein A can thus be thought of as a toxin,

essential for pathogenesis, whose molecular attributes must be neutralized in order to achieve protective immunity. By generating non-toxigenic variants unable to bind Igs via Fcγ or VH<sub>3</sub>-Fab domains, the inventors measure here for the first time protein A neutralizing immune responses as a correlate for protective immunity against *S. aureus* infection. In contrast to many methicillin-sensitive strains, CA-MRSA isolate USA300 LAC is significantly more virulent (Cheng et al., 2009). For example, immunization of experimental animals with the surface protein IsdB (Kuklin et al., 2006; Stranger-Jones et al., 2006) raises antibodies that confer protection against *S. aureus* Newman (Stranger-Jones et al., 2009) but not against USA300 challenge.

The methods utilized include:  
Bacterial Strains and Growth.

*Staphylococcus aureus* strains Newman and USA300 were grown in tryptic soy broth (TSB) at 37° C. *Escherichia coli* strains DH5α and BL21 (DE3) were grown in Luria-Bertani (LB) broth with 100 μg ml<sup>-1</sup> ampicillin at 37° C.

Rabbit Antibodies.

The coding sequence for SpA was PCR-amplified with two primers, gctgcacatgatggcgcaacacgatgaagctcaac (SEQ ID NO:35) and agtggatcctatgcttgagctttgtagcatctgc (SEQ ID NO:36) using *S. aureus* Newman template DNA. SpA-D was PCR-amplified with two primers, aacatattgcaacaagaatcaacaaagc (SEQ ID NO:38) and aaggatccagattgcttatttttagc

60

65

(SEQ ID NO:39). The sequence for SpA-D<sub>KKAA</sub> was mutagenized with two sets of primers catatgtcaacaagaataaaaaagcgccttctatgaatc (SEQ ID NO:42) and gatttcatagaagcgcctttttatctttgtgaacatag (SEQ ID NO:43) for Q9K, Q10K as well as cttcattcaagctttaaagccgc-  
 cccaagcgaagcactaac (SEQ ID NO:40) and gttagtgtcttgct-  
 tggggcgcgtttaagacttgaatgaag (SEQ ID NO:41) for D36A, D37A. The sequence of SpA<sub>KKAA</sub> was synthesized by Integrated DNA Technologies, Inc. PCR products were cloned into pET-15b generating N-terminal His<sub>6</sub> tagged recombinant protein. Plasmids were transformed into BL21 (DE3). Overnight cultures of transformants were diluted 1:100 into fresh media and grown at 37° C. to an OD<sub>600</sub> 0.5, at which point cultures were induced with 1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) and grown for an additional three hours. Bacterial cells were sedimented by centrifugation, suspended in column buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl) and disrupted with a French pressure cell at 14,000 psi. Lysates were cleared of membrane and insoluble components by ultracentrifugation at 40,000×g. Proteins in the soluble lysate were subjected to nickel-nitrilotriacetic acid (Ni-NTA, Qiagen) affinity chromatography. Proteins were eluted in column buffer containing successively higher concentrations of imidazole (100-500 mM). Protein concentrations were determined by bicinchonic acid (BCA) assay (Thermo Scientific). For antibody generation, rabbits (6 month old New-Zealand white, female, Charles River Laboratories) were immunized with 500 μg protein emulsified in Complete Freund's Adjuvant (Difco) by subcutaneous injection. For booster immunizations, proteins emulsified in Incomplete Freund's Adjuvant and injected 24 or 48 days following the initial immunization. On day 60, rabbits were bled and serum recovered.

Purified antigen (5 mg protein) was covalently linked to HiTrap NHS-activated HP columns (GE Healthcare). Antigen-matrix was used for affinity chromatography of 10-20 ml of rabbit serum at 4° C. Charged matrix was washed with 50 column volumes of PBS, antibodies eluted with elution buffer (1 M glycine, pH 2.5, 0.5 M NaCl) and immediately neutralized with 1M Tris-HCl, pH 8.5. Purified antibodies were dialyzed overnight against PBS at 4° C.

#### F(ab)<sub>2</sub> Fragments.

Affinity purified antibodies were mixed with 3 mg of pepsin at 37° C. for 30 minutes. The reaction was quenched with 1 M Tris-HCl, pH 8.5 and F(ab)<sub>2</sub> fragments were affinity purified with specific antigen-conjugated HiTrap NHS-activated HP columns. Purified antibodies were dialyzed overnight against PBS at 4° C., loaded onto SDS-PAGE gel and visualized with Coomassie Blue staining.

#### Active and Passive Immunization.

BALB/c mice (3 week old, female, Charles River Laboratories) were immunized with 50 μg protein emulsified in Complete Freund's Adjuvant (Difco) by intramuscular injection. For booster immunizations, proteins were emulsified in Incomplete Freund's Adjuvant and injected 11 days following the initial immunization. On day 20 following immunization, 5 mice were bled to obtain sera for specific antibody titers by enzyme-linked immunosorbent assay (ELISA).

Affinity purified antibodies in PBS were injected at a concentration 5 mg kg<sup>-1</sup> of experimental animal weight into the peritoneal cavity of BALB/c mice (6 week old, female, Charles River Laboratories) 24 hours prior to challenge with *S. aureus*. Animal blood was collected via periorbital vein puncture. Blood cells were removed with heparinized microhematocrit capillary tubes (Fisher) and Z-gel serum separation micro tubes (Sarstedt) were used to collect and measure antigen specific antibody titers by ELISA.

#### Mouse Renal Abscess.

Overnight cultures of *S. aureus* Newman or USA300 (LAC) were diluted 1:100 into fresh TSB and grown for 2 hours at 37° C. Staphylococci were sedimented, washed and suspended PBS at OD<sub>600</sub> of 0.4 (~1×10<sup>8</sup> CFU ml<sup>-1</sup>). Inocula were quantified by spreading sample aliquots on TSA and enumerating colonies formed. BALB/c mice (6 week old, female, Charles River Laboratories) were anesthetized via intraperitoneal injection with 100 mg ml<sup>-1</sup> ketamine and 20 mg ml<sup>-1</sup> xylazine per kilogram of body weight. Mice were infected by retro-orbital injection with 1×10<sup>7</sup> CFU of *S. aureus* Newman or 5×10<sup>6</sup> CFU of *S. aureus* USA300. On day 4 following challenge, mice were killed by CO<sub>2</sub> inhalation. Both kidneys were removed, and the staphylococcal load in one organ was analyzed by homogenizing renal tissue with PBS, 1% Triton X-100. Serial dilutions of homogenate were spread on TSA and incubated for colony formation. The remaining organ was examined by histopathology. Briefly, kidneys were fixed in 10% formalin for 24 hours at room temperature. Tissues were embedded in paraffin, thin-sectioned, stained with hematoxylin-eosin, and inspected by light microscopy to enumerate abscess lesions. All mouse experiments were performed in accordance with the institutional guidelines following experimental protocol review and approval by the Institutional Biosafety Committee (IBC) and the Institutional Animal Care and Use Committee (IACUC) at the University of Chicago.

#### Protein A Binding.

For human IgG binding, Ni-NTA affinity columns were pre-charged with 200 μg of purified proteins (SpA, SpA-D, SpA-D<sub>KKAA</sub>, and SrtA) in column buffer. After washing, 200 μg of human IgG (Sigma) was loaded onto the column. Protein samples were collected from washes and elutions and subjected to SDS-PAGE gel electrophoresis, followed by Coomassie Blue staining. Purified proteins (SpA, SpA<sub>KKAA</sub>, SpA-D and SpA-D<sub>KKAA</sub>) were coated onto MaxiSorp ELISA plates (NUNC) in 0.1M carbonate buffer (pH 9.5) at 1 μg ml<sup>-1</sup> concentration overnight at 4° C. Plates were next blocked with 5% whole milk followed by incubation with serial dilutions of peroxidase-conjugated human IgG, Fc or F(ab)<sub>2</sub> fragments for one hour. Plates were washed and developed using OptEIA ELISA reagents (BD). Reactions were quenched with 1 M phosphoric acid and A<sub>450</sub> readings were used to calculate half maximal titer and percent binding.

#### von Willebrand Factor (vWF) Binding Assays.

Purified proteins (SpA, SpA<sub>KKAA</sub>, SpA D and SpA-D<sub>KKAA</sub>) were coated and blocked as described above. Plates were incubated with human vWF at 1 μg ml<sup>-1</sup> concentration for two hours, then washed and blocked with human IgG for another hour. After washing, plates were incubated with serial dilution of peroxidase-conjugated antibody directed against human vWF for one hour. Plates were washed and developed using OptEIA ELISA reagents (BD). Reactions were quenched with 1 M phosphoric acid and A<sub>450</sub> readings were used to calculate half maximal titer and percent binding. For inhibition assays, plates were incubated with affinity purified F(ab)<sub>2</sub> fragments specific for SpA-D<sub>KKAA</sub> at 10 μg ml<sup>-1</sup> concentration for one hour prior to ligand binding assays.

#### Splenocyte Apoptosis.

Affinity purified proteins (150 μg of SpA, SpA-D, SpA<sub>KKAA</sub>, and SpA-D<sub>KKAA</sub>) were injected into the peritoneal cavity of BALB/c mice (6 week old, female, Charles River Laboratories). Four hours following injection, animals were killed by CO<sub>2</sub> inhalation. Their spleens were removed and homogenized. Cell debris were removed using cell strainer and suspended cells were transferred to ACK lysis buffer (0.15 M NH<sub>4</sub>Cl, 10 mM KHCO<sub>3</sub>, 0.1 mM EDTA) to lyse red

blood cells. White blood cells were sedimented by centrifugation, suspended in PBS and stained with 1:250 diluted R-PE conjugated anti-CD19 monoclonal antibody (Invitrogen) on ice and in the dark for one hour. Cells were washed with 1% FBS and fixed with 4% formalin overnight at 4° C. The following day, cells were diluted in PBS and analyzed by flow cytometry. The remaining organ was examined for histopathology. Briefly, spleens were fixed in 10% formalin for 24 hours at room temperature. Tissues were embedded in paraffin, thin-sectioned, stained with the Apoptosis detection kit (Millipore), and inspected by light microscopy.

#### Antibody Quantification.

Sera were collected from healthy human volunteers or BALB/c mice that had been either infected with *S. aureus* Newman or USA300 for 30 days or that had been immunized with SpA-D<sub>KKAA</sub>/SpA<sub>KKAA</sub> as described above. Human/mouse IgG (Jackson Immunology Laboratory), SpA<sub>KKAA</sub>, and CRM<sub>197</sub> were blotted onto nitrocellulose membrane. Membranes were blocked with 5% whole milk, followed by incubation with either human or mouse sera. IRDye 700DX conjugated affinity purified anti-human/mouse IgG (Rockland) was used to quantify signal intensities using the Odyssey™ infrared imaging system (Li-cor). Experiments with blood from human volunteers involved protocols that were reviewed, approved and performed under regulatory supervision of The University of Chicago's Institutional Review Board (IRB).

#### Statistical Analysis.

Two tailed Student's t tests were performed to analyze the statistical significance of renal abscess, ELISA, and B cell superantigen data.

### REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

U.S. Pat. No. 3,791,932  
 U.S. Pat. No. 3,949,064  
 U.S. Pat. No. 4,174,384  
 U.S. Pat. No. 4,338,298  
 U.S. Pat. No. 4,356,170  
 U.S. Pat. No. 4,367,110  
 U.S. Pat. No. 4,372,945  
 U.S. Pat. No. 4,452,901  
 U.S. Pat. No. 4,474,757  
 U.S. Pat. No. 4,554,101  
 U.S. Pat. No. 4,578,770  
 U.S. Pat. No. 4,596,792  
 U.S. Pat. No. 4,599,230  
 U.S. Pat. No. 4,599,231  
 U.S. Pat. No. 4,601,903  
 U.S. Pat. No. 4,608,251  
 U.S. Pat. No. 4,683,195  
 U.S. Pat. No. 4,683,202  
 U.S. Pat. No. 4,684,611  
 U.S. Pat. No. 4,690,915  
 U.S. Pat. No. 4,690,915  
 U.S. Pat. No. 4,748,018  
 U.S. Pat. No. 4,800,159  
 U.S. Pat. No. 4,879,236  
 U.S. Pat. No. 4,952,500  
 U.S. Pat. No. 5,084,269  
 U.S. Pat. No. 5,199,942  
 U.S. Pat. No. 5,221,605  
 U.S. Pat. No. 5,238,808

U.S. Pat. No. 5,302,523  
 U.S. Pat. No. 5,310,687  
 U.S. Pat. No. 5,322,783  
 U.S. Pat. No. 5,384,253  
 5 U.S. Pat. No. 5,464,765  
 U.S. Pat. No. 5,512,282  
 U.S. Pat. No. 5,512,282  
 U.S. Pat. No. 5,538,877  
 U.S. Pat. No. 5,538,880  
 10 U.S. Pat. No. 5,548,066  
 U.S. Pat. No. 5,550,318  
 U.S. Pat. No. 5,563,055  
 U.S. Pat. No. 5,580,859  
 U.S. Pat. No. 5,589,466  
 15 U.S. Pat. No. 5,591,616  
 U.S. Pat. No. 5,610,042  
 U.S. Pat. No. 5,620,896  
 U.S. Pat. No. 5,648,240  
 20 U.S. Pat. No. 5,656,610  
 U.S. Pat. No. 5,702,932  
 U.S. Pat. No. 5,736,524  
 U.S. Pat. No. 5,780,448  
 U.S. Pat. No. 5,789,215  
 25 U.S. Pat. No. 5,801,234  
 U.S. Pat. No. 5,840,846  
 U.S. Pat. No. 5,843,650  
 U.S. Pat. No. 5,846,709  
 U.S. Pat. No. 5,846,783  
 30 U.S. Pat. No. 5,849,497  
 U.S. Pat. No. 5,849,546  
 U.S. Pat. No. 5,849,547  
 U.S. Pat. No. 5,858,652  
 U.S. Pat. No. 5,866,366  
 35 U.S. Pat. No. 5,871,986  
 U.S. Pat. No. 5,916,776  
 U.S. Pat. No. 5,922,574  
 U.S. Pat. No. 5,925,565  
 U.S. Pat. No. 5,925,565  
 40 U.S. Pat. No. 5,928,905  
 U.S. Pat. No. 5,928,906  
 U.S. Pat. No. 5,932,451  
 U.S. Pat. No. 5,935,819  
 U.S. Pat. No. 5,935,825  
 45 U.S. Pat. No. 5,939,291  
 U.S. Pat. No. 5,942,391  
 U.S. Pat. No. 5,945,100  
 U.S. Pat. No. 5,958,895  
 U.S. Pat. No. 5,981,274  
 50 U.S. Pat. No. 5,994,624  
 U.S. Pat. No. 6,008,341  
 U.S. Pat. No. 6,288,214  
 U.S. Pat. No. 6,294,177  
 U.S. Pat. No. 6,651,655  
 55 U.S. Pat. No. 6,656,462  
 U.S. Pat. No. 6,733,754  
 U.S. Pat. No. 6,756,361  
 U.S. Pat. No. 6,770,278  
 U.S. Pat. No. 6,793,923  
 60 U.S. Pat. No. 6,814,971  
 U.S. Pat. No. 6,936,258  
 U.S. Patent Appln. 2002/0169288  
 U.S. Patent Appln. 2003/0153022  
 Abdallah et al., *Mol. Microbiol.*, 62, 667-679, 2006.  
 65 Abdallah et al., *Nat. Rev. Microbiol.*, 5, 883-891, 2007.  
 Albus et al., *Infect. Immun.*, 59:1008-1014, 1991.  
 An, *J. Virol.*, 71(3):2292-302, 1997.

- Anavi, Sc. thesis from the department of Molecular Microbiology and Biotechnology of the Tel-Aviv University, Israel, 1998.
- Andersen et al., *J. Immunol.*, 154, 3359-3372, 1995.
- Angel et al., *Cell*, 49:729, 1987b.
- Angel et al., *Mol. Cell. Biol.*, 7:2256, 1987a.
- Archer, *Clin. Infect. Dis.*, 26, 1179-1181, 1998.
- Atchison and Perry, *Cell*, 46:253, 1986.
- Atchison and Perry, *Cell*, 48:121, 1987.
- Ausubel et al., *In: Current Protocols in Molecular Biology*, John, Wiley & Sons, Inc, New York, 1996.
- Baba et al., *J. Bacteriol.* 190:300-310, 2007.
- Bae and Schneewind, *Plasmid*, 55:58-63, 2006.
- Bae et al., *Proc. Natl. Acad. Sci. USA*, 101, 12312-12317, 2004.
- Banerji et al., *Cell*, 27(2 Pt 1):299-308, 1981.
- Banerji et al., *Cell*, 33(3):729-740, 1983.
- Barany and Merrifield, In: *The Peptides*, Gross and Meienhofer (Eds.), Academic Press, NY, 1-284, 1979.
- Behring E A. Über das Zustandekommen der Diphtherie—Immunität bei Thieren. Deutsche Medizinische Wochenschrift, 16:1145-8, 1890.
- Bellus, *J. Macromol. Sci. Pure Appl. Chem.*, A31(1): 1355-1376, 1994.
- Berkhout et al., *Cell*, 59:273-282, 1989.
- Birch-Hirschfeld, L. 1934. Über die Agglutination von Staphylokokken durch Bestandteile des Säugetierblutplasmas. Klinische Wochenschrift 13:331.
- Bjerketorp et al., *FEMS Microbiol. Lett.*, 234:309-314, 2004.
- Blanar et al., *EMBO J.*, 8:1139, 1989.
- Bodine and Ley, *EMBO J.*, 6:2997, 1987.
- Borrebaeck, In: *Antibody Engineering—A Practical Guide*, W. H. Freeman and Co., 1992.
- Boshart et al., *Cell*, 41:521, 1985.
- Bosze et al., *EMBO J.*, 5(7):1615-1623, 1986.
- Boucher and Corey. *Clin. Infect. Dis.* 46:S334-S349, 2008.
- Braddock et al., *Cell*, 58:269, 1989.
- Brown et al., *Biochemistry*, 37:4397-4406, 1998.
- Bubeck Wardenburg and Schneewind. *J. Exp. Med.* 205:287-294, 2008.
- Bubeck-Wardenburg et al., *Infect. Immun.* 74:1040-1044, 2007.
- Bubeck-Wardenburg et al., *Proc. Natl. Acad. Sci. USA*, 103: 13831-13836, 2006.
- Bulla and Siddiqui, *J. Virol.*, 62:1437, 1986.
- Burke et al., *J. Inf. Dis.*, 170:1110-1119, 1994.
- Burlak et al., *Cell Microbiol.*, 9:1172-1190, 2007.
- Burts and Missiakas, *Mol. Microbiol.*, 69:736-46, 2008.
- Burts et al., *Proc. Natl. Acad. Sci. USA*, 102:1169-1174, 2005.
- Campbell and Villarreal, *Mol. Cell. Biol.*, 8:1993, 1988.
- Campere and Tilghman, *Genes and Dev.*, 3:537, 1989.
- Campo et al., *Nature*, 303:77, 1983.
- Carbonelli et al., *FEMS Microbiol. Lett.*, 177(1):75-82, 1999.
- Cedergren et al., *Protein Eng.*, 6:441-448, 1993.
- Celander and Haseltine, *J. Virology*, 61:269, 1987.
- Celander et al., *J. Virology*, 62:1314, 1988.
- Cespedes et al., *J. Infect. Dis.*, 191(3):444-52, 2005.
- Champion et al., *Science*, 313:1632-1636, 2006.
- Chandler et al., *Cell*, 33:489, 1983.
- Chandler et al., *Proc. Natl. Acad. Sci. USA*, 94(8):3596-601, 1997.
- Chang et al., *Lancet.*, 362(9381):362-369, 2003.
- Chang et al., *Mol. Cell. Biol.*, 9:2153, 1989.
- Chatterjee et al., *Proc. Natl. Acad. Sci. USA*, 86:9114, 1989.
- Chen and Okayama, *Mol. Cell Biol.*, 7(8):2745-2752, 1987.
- Cheng et al., *FASEB J.*, 23:1-12, 2009.

- Choi et al., *Cell*, 53:519, 1988.
- Cocea, *Biotechniques*, 23(5):814-816, 1997.
- Cohen et al., *J. Cell. Physiol.*, 5:75, 1987.
- Cosgrove et al., *Infect. Control Hosp. Epidemiol.* 26:166-174, 2005.
- Costa et al., *Mol. Cell. Biol.*, 8:81, 1988.
- Cripe et al., *EMBO J.*, 6:3745, 1987.
- Culotta and Hamer, *Mol. Cell. Biol.*, 9:1376, 1989.
- Dalbey and Widmer, *J. Biol. Chem.*, 260:15925-15931, 1985.
- Dandolo et al., *J. Virology*, 47:55-64, 1983.
- De Villiers et al., *Nature*, 312(5991):242-246, 1984.
- DeBord et al., *Infect. Immun.*, 74:4910-4914, 2006.
- DeDent et al., *EMBO J.* 27:2656-2668, 2008.
- DeDent et al., *J. Bacteriol.* 189:4473-4484, 2007.
- Deisenhofer et al., *Hoppe-Seyh Zeitsch. Physiol. Chem.* 359: 975-985, 1978.
- Deisenhofer, *Biochemistry* 20:2361-2370, 1981.
- Deschamps et al., *Science*, 230:1174-1177, 1985.
- Devereux et al., *Nucl. Acid Res.*, 12:387-395, 1984.
- Diep et al., *J. Infect. Dis.*, 193:1495-1503, 2006a.
- Diep et al., *Lancet.*, 367:731-739, 2006b.
- Dinges et al., *Clin. Microbiol. Rev.*, 13:16-34, 2000.
- Duthie and Lorenz, *J. Gen. Microbiol.*, 6:95-107, 1952.
- Edbrooke et al., *Mol. Cell. Biol.*, 9:1908, 1989.
- Edlund et al., *Science*, 230:912-916, 1985.
- Ekstedt and Yotis, *Ann. N.Y. Acad. Sci.*, 80:496-500, 1960.
- Emorl and Gaynes, *Clin. Microbiol. Rev.*, 6:428-442, 1993. EP 0786519 EP 497524 EP 497525
- Epitope Mapping Protocols In: *Methods in Molecular Biology*, Vol. 66, Morris (Ed.), 1996.
- Fechheimer, et al., *Proc Natl. Acad. Sci. USA*, 84:8463-8467, 1987.
- Feng and Holland, *Nature*, 334:6178, 1988.
- Field and Smith, *J. Comp. Pathol.*, 55:63, 1945.
- Firak and Subramanian, *Mol. Cell. Biol.*, 6:3667, 1986.
- Foecking and Hofstetter, *Gene*, 45(1):101-105, 1986.
- Fortune et al., *Proc Natl. Acad. Sci. USA*, 102:10676-10681, 2005.
- Foster, *Nat. Rev. Microbiol.*, 3:948-958, 2005.
- Fournier et al., *Infect. Immun.*, 45:87-93, 1984.
- Fraley et al., *Proc. Natl. Acad. Sci. USA*, 76:3348-3352, 1979.
- Friedrich et al., *Nature*, 425:535-539, 2003.
- Fujita et al., *Cell*, 49:357, 1987.
- GB Appln. 2 202 328
- Gilles et al., *Cell*, 33:717, 1983.
- Gloss et al., *EMBO J.*, 6:3735, 1987.
- Godbout et al., *Mol. Cell. Biol.*, 8:1169, 1988.
- Gomez et al., *EMBO J.* 26:701-709, 2007.
- Gomez et al., *J. Biol. Chem.* 281:20190-20196, 2006.
- Gomez et al., *Nature Med.* 10:842-8, 2004.
- Goodbourn and Maniatis, *Proc. Natl. Acad. Sci. USA*, 85:1447, 1988.
- Goodbourn et al., *Cell*, 45:601, 1986.
- Goodyear and Silverman, *J. Exp. Med.*, 197:1125-1139, 2003.
- Goodyear and Silverman, *Proc. Nat. Acad. Sci. USA*, 101: 11392-11397, 2004.
- Gopal, *Mol. Cell Biol.*, 5:1188-1190, 1985.
- Gouda et al., *Biochemistry*, 31(40):9665-72, 1992.
- Gouda et al., *Biochemistry*, 37:129-36, 1998.
- Graham and Van Der Eb, *Virology*, 52:456-467, 1973.
- Graille et al., *Proc. Nat. Acad. Sci. USA* 97:5399-5404, 2000.
- Greene et al., *Immunology Today*, 10:272, 1989
- Grosschedl and Baltimore, *Cell*, 41:885, 1985.
- Guinn et al., *Mol. Microbiol.*, 51:359-370, 2004.

- Guss et al., *Eur. J. Biochem.* 138:413-420, 1984.
- Harland and Weintraub, *J. Cell Biol.*, 101(3):1094-1099, 1985.
- Harlow et al., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., Chapter 8, 1988.
- Hartleib et al., *Blood* 96:2149-2156, 2000.
- Harvey et al., *Proc. Natl. Acad. Sci. USA*, 83:1084-1088, 1986.
- Haslinger and Karin, *Proc. Natl. Acad. Sci. USA*, 82:8572, 1985.
- Hauber and Cullen, *J Virology*, 62:673, 1988.
- Hen et al., *Nature*, 321:249, 1986.
- Hensel et al., *Lymphokine Res.*, 8:347, 1989.
- Herr and Clarke, *Cell*, 45:461, 1986.
- Hirochika et al., *J. Virol.*, 61:2599, 1987.
- Hirsch et al., *Mol. Cell. Biol.*, 10:1959, 1990.
- Holbrook et al., *Virology*, 157:211, 1987.
- Horlick and Benfield, *Mol. Cell. Biol.*, 9:2396, 1989.
- Hsu et al., *Proc. Natl. Acad. Sci. USA*, 100:12420-12425, 2003.
- Huang et al., *Cell*, 27:245, 1981.
- Hug et al., *Mol. Cell. Biol.*, 8:3065, 1988.
- Huston et al., In: *Methods in Enzymology*, Langone (Ed.), Academic Press, NY, 203:46-88, 1991.
- Hwang et al., *Mol. Cell. Biol.*, 10:585, 1990.
- Imagawa et al., *Cell*, 51:251, 1987.
- Imbra and Karin, *Nature*, 323:555, 1986.
- Imler et al., *Mol. Cell. Biol.*, 7:2558, 1987.
- Imperiale and Nevins, *Mol. Cell. Biol.*, 4:875, 1984.
- Innis et al., *Proc Natl Acad Sci USA*, 85(24):9436-9440, 1988.
- Inouye and Inouye, *Nucleic Acids Res.*, 13: 3101-3109, 1985.
- Jakobovits et al., *Mol. Cell. Biol.*, 8:2555, 1988.
- Jameel and Siddiqui, *Mol. Cell. Biol.*, 6:710, 1986.
- Jansson et al., *FEMS Immunol. Med. Microbiol.* 20:69-78 1998.
- Jaynes et al., *Mol. Cell. Biol.*, 8:62, 1988.
- Jensen, *Acta Path. Microbiol. Scand.* 44:421-428, 1958.
- Johnson et al., *Methods in Enzymol.*, 203:88-99, 1991.
- Johnson et al., *Mol. Cell. Biol.*, 9:3393, 1989.
- Jones, *Carb. Research*, 340:1097-1106, 2005.
- Jonsson et al., *Oral Dis.*, 8(3):130-140, 2002.
- Joyce et al., *Carbohydrate Research* 338:903-922 (2003)
- Kadesch and Berg, *Mol. Cell. Biol.*, 6:2593, 1986.
- Kaeppler et al., *Plant Cell Rep.*, 8:415-418, 1990.
- Kaneda et al., *Science*, 243:375-378, 1989.
- Karin et al., *Mol. Cell. Biol.*, 7:606, 1987.
- Katinka et al., *Cell*, 20:393, 1980.
- Kato et al., *J. Biol. Chem.*, 266:3361-3364, 1991.
- Kawamoto et al., *Mol. Cell. Biol.*, 8:267, 1988.
- Kennedy et al., *Proc. Natl. Acad. Sci. USA* 105:1327-1332, 2008.
- Kiledjian et al., *Mol. Cell. Biol.*, 8:145, 1988.
- Kinoshita, M., N. Kobayashi, S. Nagashima, M. Ishino, S. Otokozawa, K. Mise, A. Sumi, H. Tsutsumi, N. Uehara, N. Watanabe, and M. Endo. 2008. Diversity of staphylococagulase and identification of novel variants of staphylococagulase gene in *Staphylococcus aureus*. *Microbiol. Immunol.* 52:334-348.
- Klamut et al., *Mol. Cell. Biol.*, 10:193, 1990.
- Klevens et al., *Clin. Infect. Dis.*, 2008; 47:927-30, 2008.
- Klevens et al., *JAMA*, 298:1763-1771, 2007.
- Koch et al., *Mol. Cell. Biol.*, 9:303, 1989.
- Kohler and Milstein, *Nature* 256:495-497 (1975

- Kriegler and Botchan, In: *Eukaryotic Viral Vectors*, Gluzman (Ed.), Cold Spring Harbor: Cold Spring Harbor Laboratory, NY, 1982.
- Kriegler and Botchan, *Mol. Cell. Biol.*, 3:325, 1983.
- Kriegler et al., *Cell*, 38:483, 1984a.
- Kriegler et al., *Cell*, 53:45, 1988.
- Kriegler et al., In: *Cancer Cells 2/Oncogenes and Viral Genes*, Van de Woude et al. eds, Cold Spring Harbor, Cold Spring Harbor Laboratory, 1984b.
- Kroh et al., *Proc. Natl. Acad. Sci. USA*, 106:7786-7791, 2009.
- Kuhl et al., *Cell*, 50:1057, 1987.
- Kuklin et al., *Infect. Immun.*, 74:2215-23, 2006.
- Kunz et al., *Nucl. Acids Res.*, 17:1121, 1989.
- Kuroda et al., *Lancet.*, 357:1225-1240, 2001.
- Kyte and Doolittle, *J. Mol. Biol.*, 157(1):105-132, 1982.
- Lagergard et al., *Eur. J. Clin. Microbiol. Infect. Dis.*, 11:341-5, 1992.
- Lam et al., *J. Bacteriol.*, 86:87-91, 1963.
- Larsen et al., *Proc Natl. Acad. Sci. USA.*, 83:8283, 1986, 1963.
- Laspia et al., *Cell*, 59:283, 1989.
- Latimer et al., *Mol. Cell. Biol.*, 10:760, 1990.
- Lee et al., *Nature*, 294:228, 1981.
- Lee et al., *Nucleic Acids Res.*, 12:4191-206, 1984.
- Lee, *Trends Microbiol.* 4(4):162-166, 1996.
- Levenson et al., *Hum. Gene Ther.*, 9(8):1233-1236, 1998.
- Levinson et al., *Nature*, 295:79, 1982.
- Lin et al., *Mol. Cell. Biol.*, 10:850, 1990.
- Lowy, *New Engl. J. Med.*, 339:520-532, 1998.
- Luria et al., *EMBO J.*, 6:3307, 1987.
- Lusky and Botchan, *Proc. Natl. Acad. Sci. USA*, 83:3609, 1986.
- Lusky et al., *Mol. Cell. Biol.*, 3:1108, 1983.
- Macejak and Sarnow, *Nature*, 353:90-94, 1991.
- MacGurn et al., *Mol. Microbiol.*, 57:1653-1663, 2005.
- Maira-Litran et al., *Infect. Immun.*, 70:4433-4440, 2002.
- Maira-Litran et al., *Vaccine*, 22:872-879, 2004.
- Majors and Varmus, *Proc. Natl. Acad. Sci. USA*, 80:5866, 1983.
- Markwardt, *Untersuchungen über Hirudin. Naturwissenschaften*, 41:537-538, 1955.
- Mazmanian et al., *Mol. Microbiol.* 40, 1049-1057, 2001.
- Mazmanian et al., *Mol. Microbiol.*, 40(5):1049-1057, 2001.
- Mazmanian et al., *Proc. Natl. Acad. Sci. USA*, 97:5510-5515, 2000.
- Mazmanian et al., *Science*, 285(5428):760-3, 1999.
- McLaughlin et al., *PLoS Pathog.*, 3:e105, 2007.
- McNeall et al., *Gene*, 76:81, 1989.
- Mernaugh et al., In: *Molecular Methods in Plant Pathology*, Singh et al. (Eds.), CRC Press Inc., Boca Raton, Fla., 359-365, 1995.
- Merrifield, *Science*, 232(4748):341-347, 1986.
- Miksicek et al., *Cell*, 46:203, 1986.
- Mordacq and Linzer, *Genes and Dev.*, 3:760, 1989.
- Moreau et al., *Carbohydrate Res.*, 201:285-297, 1990.
- Moreau et al., *Nucl. Acids Res.*, 9:6047, 1981.
- Moreillon et al., *Infect. Immun.*, 63:4738-4743, 1995.
- Moreillon et al., *Infect. Immun.*, 63:4738-4743, 1995.
- Mosmann and Coffman, *Ann. Rev. Immunol.*, 7:145-173, 1989.
- Muesing et al., *Cell*, 48:691, 1987.
- Musher et al., *Medicine (Baltimore)*, 73:186-208, 1994.
- Navarre and Schneewind, *J. Biol. Chem.*, 274:15847-15856, 1999.
- Needleman & Wunsch, *J. Mol. Biol.*, 48:443, 1970.
- Ng et al., *Nucl. Acids Res.*, 17:601, 1989.

Nicolau and Sene, *Biochim. Biophys. Acta*, 721:185-190, 1982.

Nicolau et al., *Methods Enzymol.*, 149:157-176, 1987.

Novick, *Mol. Microbiol.*, 48:1429-1449, 2003.

O'Brien et al., *Mol. Microbiol.* 44:1033-1044, 2002.

O'Seaghdha et al., *FEBS J.* 273:4831-4841, 2006.

Omirulleh et al., *Plant Mol. Biol.*, 21(3):415-28, 1993.

Ondek et al., *EMBO J.*, 6:1017, 1987.

Ornitz et al., *Mol. Cell. Biol.*, 7:3466, 1987.

Pallen, *Trends Microbiol.*, 10:209-212, 2002.

Palmiter et al., *Nature*, 300:611, 1982.

Palmqvist et al., *Microbes. Infect.*, 7:1501-11, 2005.

Panizzi et al., *J. Biol. Chem.*, 281:1179-1187, 2006.

PCT Appln. PCT/US89/01025

PCT Appln. WO 00/02523

PCT Appln. WO 00/12132

PCT Appln. WO 00/12689

PCT Appln. WO 00/15238

PCT Appln. WO 01/34809

PCT Appln. WO 01/60852

PCT Appln. WO 01/98499

PCT Appln. WO 02/059148

PCT Appln. WO 02/094868

PCT Appln. WO 03/53462

PCT Appln. WO 04/43407

PCT Appln. WO 06/032472

PCT Appln. WO 06/032475

PCT Appln. WO 06/032500

PCT Appln. WO 07/113,222

PCT Appln. WO 07/113,223

PCT Appln. WO 94/09699

PCT Appln. WO 95/06128

PCT Appln. WO 95/08348

PCT Appln. WO 98/57994

Pearson & Lipman, *Proc. Natl. Acad. Sci. USA*, 85:2444, 1988.

Pech et al., *Mol. Cell. Biol.*, 9:396, 1989.

Pelletier and Sonenberg, *Nature*, 334(6180):320-325, 1988.

Perez-Stable and Constantini, *Mol. Cell. Biol.*, 10:1116, 1990.

Phonimdaeng et al., *Mol. Microbiol.*, 4:393-404, 1990.

Picard and Schaffner, *Nature*, 307:83, 1984.

Pinkert et al., *Genes and Dev.*, 1:268, 1987.

Ponta et al., *Proc. Natl. Acad. Sci. USA*, 82:1020, 1985.

Porton et al., *Mol. Cell. Biol.*, 10:1076, 1990.

Potrykus et al., *Mol. Gen. Genet.*, 199(2):169-177, 1985.

Pugsley, *Microbiol. Rev.*, 57:50-108, 1993.

Pym et al., *Mol. Microbiol.*, 46: 709-717, 2002.

Pym et al., *Nat. Med.*, 9:533-539, 2003.

Queen and Baltimore, *Cell*, 35:741, 1983.

Quinn et al., *Mol. Cell. Biol.*, 9:4713, 1989.

Redondo et al., *Science*, 247:1225, 1990.

Reisman and Rotter, *Mol. Cell. Biol.*, 9:3571, 1989.

Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1289-1329, 1990.

Resendez Jr. et al., *Mol. Cell. Biol.*, 8:4579, 1988.

Ripe et al., *Mol. Cell. Biol.*, 9:2224, 1989.

Rippe, et al., *Mol. Cell. Biol.*, 10:689-695, 1990.

Rittling et al., *Nuc. Acids Res.*, 17:1619, 1989.

Roben et al., *J. Immunol.* 154:6437-6445, 1995.

Rosen et al., *Cell*, 41:813, 1988.

Sakai et al., *Genes and Dev.*, 2:1144, 1988.

Salid-Salim et al., *Infect. Control Hosp. Epidemiol.* 24:451-455, 2003.

Sambrook et al., *In: Molecular cloning*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 2001.

Schaffner et al., *J. Mol. Biol.*, 201:81, 1988.

Schneewind et al., *Cell* 70:267-281, 1992.

Schneewind et al., *EMBO*, 12:4803-4811, 1993.

Schneewind et al., *Science*, 268:103-6, 1995.

Searle et al., *Mol. Cell. Biol.*, 5:1480, 1985.

5 Sharp and Marciniak, *Cell*, 59:229, 1989.

Shaul and Ben-Levy, *EMBO J.*, 6:1913, 1987.

Shaw et al., *Microbiology*, 150:217-228, 2004.

Sheagren, *N. Engl. J. Med.* 310:1368-1373, 1984.

Sherman et al., *Mol. Cell. Biol.*, 9:50, 1989.

10 Shopsin et al., *J. Clin. Microbiol.*, 37:3556-63, 1999.

Sibbald et al., *Microbiol. Mol. Biol. Rev.*, 70:755-788, 2006.

Silverman and Goodyear. *Nat. Rev. Immunol.*, 6:465-75, 2006.

15 Sjodahl, *Eur. J. Biochem.* 73:343-351, 1977.

Sjoquist et al., *Eur. J. Biochem.* 30:190-194, 1972.

Sleigh and Lockett, *J. EMBO*, 4:3831, 1985.

Smith & Waterman, *Adv. Appl. Math.*, 2:482, 1981.

Smith et al., *Brit. J. Exp. Pathol.*, 28:57, 1947.

20 Sorensen et al., *Infect. Immun.*, 63:1710-1717, 1995.

Spalholz et al., *Cell*, 42:183, 1985.

Spandau and Lee, *J. Virology*, 62:427, 1988.

Spandidos and Wilkie, *EMBO J.*, 2:1193, 1983.

Stanley et al., *Proc. Natl. Acad. Sci. USA*, 100:13001-13006, 2003.

25 Stephens and Hentschel, *Biochem. J.*, 248:1, 1987.

Stewart and Young, In: *Solid Phase Peptide Synthesis*, 2d. ed., Pierce Chemical Co., 1984.

Stranger-Jones et al., *Proc. Nat. Acad. Sci. USA*, 103:16942-16947, 2006.

30 Stuart et al., *Nature*, 317:828, 1985.

Studier et al., *Methods Enzymol.* 185:60-89 1990.

Sullivan and Peterlin, *Mol. Cell. Biol.*, 7:3315, 1987.

Swartzendruber and Lehman, *J. Cell. Physiology*, 85:179, 1975.

Takebe et al., *Mol. Cell. Biol.*, 8:466, 1988.

Tam et al., *J. Am. Chem. Soc.*, 105:6442, 1983.

Tavernier et al., *Nature*, 301:634, 1983.

Taylor and Kingston, *Mol. Cell. Biol.*, 10:165, 1990a.

40 Taylor and Kingston, *Mol. Cell. Biol.*, 10:176, 1990b.

Taylor et al., *J. Biol. Chem.*, 264:15160, 1989.

Thiesen et al., *J. Virology*, 62:614, 1988.

Thomson et al., *J. Immunol.*, 157(2):822-826, 1996.

Tigges et al., *J. Immunol.*, 156(10):3901-3910, 1996.

45 Tigges et al., *J. Immunol.*, 156(10):3901-3910, 1996.

Ton-That et al., *Proc. Natl. Acad. Sci. USA*, 96(22):12424-9, 1999.

Treisman, *Cell*, 42:889, 1985.

Tronche et al., *Mol. Biol. Med.*, 7:173, 1990.

50 Trudel and Constantini, *Genes and Dev.*, 6:954, 1987.

Tyndell et al., *Nuc. Acids. Res.*, 9:6231, 1981.

Uhlen et al., *J. Biol. Chem.* 259:1695-1702 and 13628 (Corr.) 1984.

van den Ent and Lowe, *FEBS Lett.*, 579:3837-3841, 2005.

55 van Wely et al., *FEMS Microbiol. Rev.*, 25:437-454, 2001.

Vannice and Levinson, *J. Virology*, 62:1305, 1988.

Vasseur et al., *Proc Natl. Acad. Sci. USA*, 77:1068, 1980.

Vaughan, et al., *Nat. Biotech.* 16: 535-539, 1998.

Wang and Calame, *Cell*, 47:241, 1986.

60 Weber et al., *Cell*, 36:983, 1984.

Weinberger et al. *Mol. Cell. Biol.*, 8:988, 1984.

Weiss et al., *J. Antimicrob. Chemother.*, 53(3):480-6, 2004.

Winoto and Baltimore, *Cell*, 59:649, 1989.

Wong et al., *Gene*, 10:87-94, 1980.

Xu et al., *J. Infect. Dis.*, 189:2323-2333, 2004.

Xu et al., *Mol. Microbiol.*, 66(3):787-800, 2007.

Yutzey et al. *Mol. Cell. Biol.*, 9:1397, 1989.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 64

<210> SEQ ID NO 1  
 <211> LENGTH: 150  
 <212> TYPE: DNA  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 1

```
ttaacaaaag atcaacaaaag cgccttctat gaaatcttga acatgcctaa cttaaacgaa    60
gcgcaacgta acggcttcat tcaaagtctt aaagacgacc caagccaaag cactaatggt    120
ttaggtgaag ctaaaaaatt aaacgaatct                                     150
```

<210> SEQ ID NO 2  
 <211> LENGTH: 54  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 2

```
Gln Gln Asn Asn Phe Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile
1           5           10           15
Leu Asn Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln
20           25           30
Ser Leu Lys Asp Asp Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala
35           40           45
Lys Lys Leu Asn Glu Ser
50
```

<210> SEQ ID NO 3  
 <211> LENGTH: 51  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 3

```
Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr Gln Val Leu Asn Met
1           5           10           15
Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys
20           25           30
Asp Asp Pro Ser Gln Ser Ala Asn Val Leu Gly Glu Ala Gln Lys Leu
35           40           45
Asn Asp Ser
50
```

<210> SEQ ID NO 4  
 <211> LENGTH: 52  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 4

```
Asn Asn Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu Asn
1           5           10           15
Met Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu
20           25           30
Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ser Glu Ala Lys Lys
35           40           45
Leu Asn Glu Ser
50
```

<210> SEQ ID NO 5  
 <211> LENGTH: 52

-continued

<212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 5

```

Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His
1           5           10           15
Leu Pro Asn Leu Thr Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu
           20           25           30
Lys Asp Asp Pro Ser Val Ser Lys Glu Ile Leu Ala Glu Ala Lys Lys
           35           40           45
Leu Asn Asp Ala
           50
  
```

<210> SEQ ID NO 6  
 <211> LENGTH: 52  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 6

```

Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His
1           5           10           15
Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu
           20           25           30
Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys Lys
           35           40           45
Leu Asn Asp Ala
           50
  
```

<210> SEQ ID NO 7  
 <211> LENGTH: 52  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (7)..(8)  
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (34)..(35)  
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<400> SEQUENCE: 7

```

Asn Asn Phe Asn Lys Asp Xaa Xaa Ser Ala Phe Tyr Glu Ile Leu Asn
1           5           10           15
Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser Leu
           20           25           30
Lys Xaa Xaa Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys Lys
           35           40           45
Leu Asn Glu Ser
           50
  
```

<210> SEQ ID NO 8  
 <211> LENGTH: 52  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE  
 <222> LOCATION: (7)..(8)  
 <223> OTHER INFORMATION: where X is any amino acid other than Q  
 <220> FEATURE:  
 <221> NAME/KEY: MISC\_FEATURE  
 <222> LOCATION: (12)..(35)  
 <223> OTHER INFORMATION: where Y is any amion acid other than D

-continued

&lt;400&gt; SEQUENCE: 8

```

Asn Asn Phe Asn Lys Asp Xaa Xaa Ser Ala Phe Tyr Glu Ile Leu Asn
1           5           10           15
Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser Leu
20           25           30
Lys Tyr Tyr Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys Lys
35           40           45
Leu Asn Glu Ser
50

```

&lt;210&gt; SEQ ID NO 9

&lt;211&gt; LENGTH: 450

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 9

```

Met Lys Lys Lys Asn Ile Tyr Ser Ile Arg Lys Leu Gly Val Gly Ile
1           5           10           15
Ala Ser Val Thr Leu Gly Thr Leu Leu Ile Ser Gly Gly Val Thr Pro
20           25           30
Ala Ala Asn Ala Ala Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr
35           40           45
Gln Val Leu Asn Met Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe
50           55           60
Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Val Leu Gly
65           70           75           80
Glu Ala Gln Lys Leu Asn Asp Ser Gln Ala Pro Lys Ala Asp Ala Gln
85           90           95
Gln Asn Asn Phe Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile Leu
100          105          110
Asn Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser
115          120          125
Leu Lys Asp Asp Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys
130          135          140
Lys Leu Asn Glu Ser Gln Ala Pro Lys Ala Asp Asn Asn Phe Asn Lys
145          150          155          160
Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu Asn
165          170          175
Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser
180          185          190
Gln Ser Ala Asn Leu Leu Ser Glu Ala Lys Lys Leu Asn Glu Ser Gln
195          200          205
Ala Pro Lys Ala Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe
210          215          220
Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly
225          230          235          240
Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Val Ser Lys Glu Ile Leu
245          250          255
Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys Glu Glu Asp
260          265          270
Asn Lys Lys Pro Gly Lys Glu Asp Gly Asn Lys Pro Gly Lys Glu Asp
275          280          285
Gly Asn Lys Pro Gly Lys Glu Asp Asn Lys Lys Pro Gly Lys Glu Asp
290          295          300

```

-continued

---

Gly Asn Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp  
 305 310 315 320  
 Gly Asn Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp  
 325 330 335  
 Gly Asn Lys Pro Gly Lys Glu Asp Gly Asn Lys Pro Gly Lys Glu Asp  
 340 345 350  
 Gly Asn Gly Val His Val Val Lys Pro Gly Asp Thr Val Asn Asp Ile  
 355 360 365  
 Ala Lys Ala Asn Gly Thr Thr Ala Asp Lys Ile Ala Ala Asp Asn Lys  
 370 375 380  
 Leu Ala Asp Lys Asn Met Ile Lys Pro Gly Gln Glu Leu Val Val Asp  
 385 390 395 400  
 Lys Lys Gln Pro Ala Asn His Ala Asp Ala Asn Lys Ala Gln Ala Leu  
 405 410 415  
 Pro Glu Thr Gly Glu Glu Asn Pro Phe Ile Gly Thr Thr Val Phe Gly  
 420 425 430  
 Gly Leu Ser Leu Ala Leu Gly Ala Ala Leu Leu Ala Gly Arg Arg Arg  
 435 440 445  
 Glu Leu  
 450

<210> SEQ ID NO 10  
 <211> LENGTH: 450  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 10

Met Lys Lys Lys Asn Ile Tyr Ser Ile Arg Lys Leu Gly Val Gly Ile  
 1 5 10 15  
 Ala Ser Val Thr Leu Gly Thr Leu Leu Ile Ser Gly Gly Val Thr Pro  
 20 25 30  
 Ala Ala Asn Ala Ala Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr  
 35 40 45  
 Gln Val Leu Asn Met Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe  
 50 55 60  
 Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Val Leu Gly  
 65 70 75 80  
 Glu Ala Gln Lys Leu Asn Asp Ser Gln Ala Pro Lys Ala Asp Ala Gln  
 85 90 95  
 Gln Asn Asn Phe Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile Leu  
 100 105 110  
 Asn Met Pro Asn Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser  
 115 120 125  
 Leu Lys Asp Asp Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys  
 130 135 140  
 Lys Leu Asn Glu Ser Gln Ala Pro Lys Ala Asp Asn Asn Phe Asn Lys  
 145 150 155 160  
 Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu Asn  
 165 170 175  
 Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser  
 180 185 190  
 Gln Ser Ala Asn Leu Leu Ser Glu Ala Lys Lys Leu Asn Glu Ser Gln  
 195 200 205  
 Ala Pro Lys Ala Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe  
 210 215 220

-continued

Tyr Glu Ile Leu His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly  
 225 230 235 240  
 Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Val Ser Lys Glu Ile Leu  
 245 250 255  
 Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln Ala Pro Lys Glu Glu Asp  
 260 265 270  
 Asn Lys Lys Pro Gly Lys Glu Asp Gly Asn Lys Pro Gly Lys Glu Asp  
 275 280 285  
 Gly Asn Lys Pro Gly Lys Glu Asp Asn Lys Lys Pro Gly Lys Glu Asp  
 290 295 300  
 Gly Asn Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp  
 305 310 315 320  
 Gly Asn Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp  
 325 330 335  
 Gly Asn Lys Pro Gly Lys Glu Asp Gly Asn Lys Pro Gly Lys Glu Asp  
 340 345 350  
 Gly Asn Gly Val His Val Val Lys Pro Gly Asp Thr Val Asn Asp Ile  
 355 360 365  
 Ala Lys Ala Asn Gly Thr Thr Ala Asp Lys Ile Ala Ala Asp Asn Lys  
 370 375 380  
 Leu Ala Asp Lys Asn Met Ile Lys Pro Gly Gln Glu Leu Val Val Asp  
 385 390 395 400  
 Lys Lys Gln Pro Ala Asn His Ala Asp Ala Asn Lys Ala Gln Ala Leu  
 405 410 415  
 Pro Glu Thr Gly Glu Glu Asn Pro Phe Ile Gly Thr Thr Val Phe Gly  
 420 425 430  
 Gly Leu Ser Leu Ala Leu Gly Ala Ala Leu Leu Ala Gly Arg Arg Arg  
 435 440 445  
 Glu Leu  
 450

<210> SEQ ID NO 11  
 <211> LENGTH: 97  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 11

Met Ala Met Ile Lys Met Ser Pro Glu Glu Ile Arg Ala Lys Ser Gln  
 1 5 10 15  
 Ser Tyr Gly Gln Gly Ser Asp Gln Ile Arg Gln Ile Leu Ser Asp Leu  
 20 25 30  
 Thr Arg Ala Gln Gly Glu Ile Ala Ala Asn Trp Glu Gly Gln Ala Phe  
 35 40 45  
 Ser Arg Phe Glu Glu Gln Phe Gln Gln Leu Ser Pro Lys Val Glu Lys  
 50 55 60  
 Phe Ala Gln Leu Leu Glu Glu Ile Lys Gln Gln Leu Asn Ser Thr Ala  
 65 70 75 80  
 Asp Ala Val Gln Glu Gln Asp Gln Gln Leu Ser Asn Asn Phe Gly Leu  
 85 90 95

Gln

<210> SEQ ID NO 12  
 <211> LENGTH: 102  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

-continued

&lt;400&gt; SEQUENCE: 12

Met Gly Gly Tyr Lys Gly Ile Lys Ala Asp Gly Gly Lys Val Asn Gln  
 1 5 10 15  
 Ala Lys Gln Leu Ala Ala Lys Ile Ala Lys Asp Ile Glu Ala Cys Gln  
 20 25 30  
 Lys Gln Thr Gln Gln Leu Ala Glu Tyr Ile Glu Gly Ser Asp Trp Glu  
 35 40 45  
 Gly Gln Phe Ala Asn Lys Val Lys Asp Val Leu Leu Ile Met Ala Lys  
 50 55 60  
 Phe Gln Glu Glu Leu Val Gln Pro Met Ala Asp His Gln Lys Ala Ile  
 65 70 75 80  
 Asp Asn Leu Ser Gln Asn Leu Ala Lys Tyr Asp Thr Leu Ser Ile Lys  
 85 90 95  
 Gln Gly Leu Asp Arg Val  
 100

&lt;210&gt; SEQ ID NO 13

&lt;211&gt; LENGTH: 1385

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 13

Met Leu Asn Arg Glu Asn Lys Thr Ala Ile Thr Arg Lys Gly Met Val  
 1 5 10 15  
 Ser Asn Arg Leu Asn Lys Phe Ser Ile Arg Lys Tyr Thr Val Gly Thr  
 20 25 30  
 Ala Ser Ile Leu Val Gly Thr Thr Leu Ile Phe Gly Leu Gly Asn Gln  
 35 40 45  
 Glu Ala Lys Ala Ala Glu Ser Thr Asn Lys Glu Leu Asn Glu Ala Thr  
 50 55 60  
 Thr Ser Ala Ser Asp Asn Gln Ser Ser Asp Lys Val Asp Met Gln Gln  
 65 70 75 80  
 Leu Asn Gln Glu Asp Asn Thr Lys Asn Asp Asn Gln Lys Glu Met Val  
 85 90 95  
 Ser Ser Gln Gly Asn Glu Thr Thr Ser Asn Gly Asn Lys Ser Ile Glu  
 100 105 110  
 Lys Glu Ser Val Gln Ser Thr Thr Gly Asn Lys Val Glu Val Ser Thr  
 115 120 125  
 Ala Lys Ser Asp Glu Gln Ala Ser Pro Lys Ser Thr Asn Glu Asp Leu  
 130 135 140  
 Asn Thr Lys Gln Thr Ile Ser Asn Gln Glu Gly Leu Gln Pro Asp Leu  
 145 150 155 160  
 Leu Glu Asn Lys Ser Val Val Asn Val Gln Pro Thr Asn Glu Glu Asn  
 165 170 175  
 Lys Lys Val Asp Ala Lys Thr Glu Ser Thr Thr Leu Asn Val Lys Ser  
 180 185 190  
 Asp Ala Ile Lys Ser Asn Ala Glu Thr Leu Val Asp Asn Asn Ser Asn  
 195 200 205  
 Ser Asn Asn Glu Asn Asn Ala Asp Ile Ile Leu Pro Lys Ser Thr Ala  
 210 215 220  
 Pro Lys Ser Leu Asn Thr Arg Met Arg Met Ala Ala Ile Gln Pro Asn  
 225 230 235 240  
 Ser Thr Asp Ser Lys Asn Val Asn Asp Leu Ile Thr Ser Asn Thr Thr  
 245 250 255

-continued

---

Leu Thr Val Val Asp Ala Asp Asn Ser Lys Thr Ile Val Pro Ala Gln  
                   260                                  265                                  270

Asp Tyr Leu Ser Leu Lys Ser Gln Ile Thr Val Asp Asp Lys Val Lys  
                   275                                  280                                  285

Ser Gly Asp Tyr Phe Thr Ile Lys Tyr Ser Asp Thr Val Gln Val Tyr  
                   290                                  295                                  300

Gly Leu Asn Pro Glu Asp Ile Lys Asn Ile Gly Asp Ile Lys Asp Pro  
 305                                  310                                  315                                  320

Asn Asn Gly Glu Thr Ile Ala Thr Ala Lys His Asp Thr Ala Asn Asn  
                                   325                                  330                                  335

Leu Ile Thr Tyr Thr Phe Thr Asp Tyr Val Asp Arg Phe Asn Ser Val  
                                   340                                  345                                  350

Lys Met Gly Ile Asn Tyr Ser Ile Tyr Met Asp Ala Asp Thr Ile Pro  
                   355                                  360                                  365

Val Asp Lys Lys Asp Val Pro Phe Ser Val Thr Ile Gly Asn Gln Ile  
                   370                                  375                                  380

Thr Thr Thr Thr Ala Asp Ile Thr Tyr Pro Ala Tyr Lys Glu Ala Asp  
 385                                  390                                  395                                  400

Asn Asn Ser Ile Gly Ser Ala Phe Thr Glu Thr Val Ser His Val Gly  
                                   405                                  410                                  415

Asn Val Glu Asp Pro Gly Tyr Tyr Asn Gln Val Val Tyr Val Asn Pro  
                                   420                                  425                                  430

Met Asp Lys Asp Leu Lys Gly Ala Lys Leu Lys Val Glu Ala Tyr His  
                   435                                  440                                  445

Pro Lys Tyr Pro Thr Asn Ile Gly Gln Ile Asn Gln Asn Val Thr Asn  
                   450                                  455                                  460

Ile Lys Ile Tyr Arg Val Pro Glu Gly Tyr Thr Leu Asn Lys Gly Tyr  
 465                                  470                                  475                                  480

Asp Val Asn Thr Asn Asp Leu Val Asp Val Thr Asp Glu Phe Lys Asn  
                                   485                                  490                                  495

Lys Met Thr Tyr Gly Ser Asn Gln Ser Val Asn Leu Asp Phe Gly Asp  
                                   500                                  505                                  510

Ile Thr Ser Ala Tyr Val Val Met Val Asn Thr Lys Phe Gln Tyr Thr  
                                   515                                  520                                  525

Asn Ser Glu Ser Pro Thr Leu Val Gln Met Ala Thr Leu Ser Ser Thr  
                   530                                  535                                  540

Gly Asn Lys Ser Val Ser Thr Gly Asn Ala Leu Gly Phe Thr Asn Asn  
 545                                  550                                  555                                  560

Gln Ser Gly Gly Ala Gly Gln Glu Val Tyr Lys Ile Gly Asn Tyr Val  
                                   565                                  570                                  575

Trp Glu Asp Thr Asn Lys Asn Gly Val Gln Glu Leu Gly Glu Lys Gly  
                                   580                                  585                                  590

Val Gly Asn Val Thr Val Thr Val Phe Asp Asn Asn Thr Asn Thr Lys  
                                   595                                  600                                  605

Val Gly Glu Ala Val Thr Lys Glu Asp Gly Ser Tyr Leu Ile Pro Asn  
                   610                                  615                                  620

Leu Pro Asn Gly Asp Tyr Arg Val Glu Phe Ser Asn Leu Pro Lys Gly  
 625                                  630                                  635                                  640

Tyr Glu Val Thr Pro Ser Lys Gln Gly Asn Asn Glu Glu Leu Asp Ser  
                                   645                                  650                                  655

Asn Gly Leu Ser Ser Val Ile Thr Val Asn Gly Lys Asp Asn Leu Ser  
                   660                                  665                                  670

-continued

---

Ala Asp Leu Gly Ile Tyr Lys Pro Lys Tyr Asn Leu Gly Asp Tyr Val  
675 680 685

Trp Glu Asp Thr Asn Lys Asn Gly Ile Gln Asp Gln Asp Glu Lys Gly  
690 695 700

Ile Ser Gly Val Thr Val Thr Leu Lys Asp Glu Asn Gly Asn Val Leu  
705 710 715 720

Lys Thr Val Thr Thr Asp Ala Asp Gly Lys Tyr Lys Phe Thr Asp Leu  
725 730 735

Asp Asn Gly Asn Tyr Lys Val Glu Phe Thr Thr Pro Glu Gly Tyr Thr  
740 745 750

Pro Thr Thr Val Thr Ser Gly Ser Asp Ile Glu Lys Asp Ser Asn Gly  
755 760 765

Leu Thr Thr Thr Gly Val Ile Asn Gly Ala Asp Asn Met Thr Leu Asp  
770 775 780

Ser Gly Phe Tyr Lys Thr Pro Lys Tyr Asn Leu Gly Asn Tyr Val Trp  
785 790 795 800

Glu Asp Thr Asn Lys Asp Gly Lys Gln Asp Ser Thr Glu Lys Gly Ile  
805 810 815

Ser Gly Val Thr Val Thr Leu Lys Asn Glu Asn Gly Glu Val Leu Gln  
820 825 830

Thr Thr Lys Thr Asp Lys Asp Gly Lys Tyr Gln Phe Thr Gly Leu Glu  
835 840 845

Asn Gly Thr Tyr Lys Val Glu Phe Glu Thr Pro Ser Gly Tyr Thr Pro  
850 855 860

Thr Gln Val Gly Ser Gly Thr Asp Glu Gly Ile Asp Ser Asn Gly Thr  
865 870 875 880

Ser Thr Thr Gly Val Ile Lys Asp Lys Asp Asn Asp Thr Ile Asp Ser  
885 890 895

Gly Phe Tyr Lys Pro Thr Tyr Asn Leu Gly Asp Tyr Val Trp Glu Asp  
900 905 910

Thr Asn Lys Asn Gly Val Gln Asp Lys Asp Glu Lys Gly Ile Ser Gly  
915 920 925

Val Thr Val Thr Leu Lys Asp Glu Asn Asp Lys Val Leu Lys Thr Val  
930 935 940

Thr Thr Asp Glu Asn Gly Lys Tyr Gln Phe Thr Asp Leu Asn Asn Gly  
945 950 955 960

Thr Tyr Lys Val Glu Phe Glu Thr Pro Ser Gly Tyr Thr Pro Thr Ser  
965 970 975

Val Thr Ser Gly Asn Asp Thr Glu Lys Asp Ser Asn Gly Leu Thr Thr  
980 985 990

Thr Gly Val Ile Lys Asp Ala Asp Asn Met Thr Leu Asp Ser Gly Phe  
995 1000 1005

Tyr Lys Thr Pro Lys Tyr Ser Leu Gly Asp Tyr Val Trp Tyr Asp  
1010 1015 1020

Ser Asn Lys Asp Gly Lys Gln Asp Ser Thr Glu Lys Gly Ile Lys  
1025 1030 1035

Asp Val Lys Val Ile Leu Leu Asn Glu Lys Gly Glu Val Ile Gly  
1040 1045 1050

Thr Thr Lys Thr Asp Glu Asn Gly Lys Tyr Arg Phe Asp Asn Leu  
1055 1060 1065

Asp Ser Gly Lys Tyr Lys Val Ile Phe Glu Lys Pro Thr Gly Leu  
1070 1075 1080

Thr Gln Thr Gly Thr Asn Thr Thr Glu Asp Asp Lys Asp Ala Asp

-continued

1085	1090	1095
Gly Gly Glu Val Asp Val Thr Ile Thr Asp His Asp Asp Phe Thr 1100	1105	1110
Leu Asp Asn Gly Tyr Tyr Glu Glu Glu Thr Ser Asp Ser Asp Ser 1115	1120	1125
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp 1130	1135	1140
Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser 1145	1150	1155
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp 1160	1165	1170
Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser 1175	1180	1185
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp 1190	1195	1200
Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser 1205	1210	1215
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp 1220	1225	1230
Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser 1235	1240	1245
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp 1250	1255	1260
Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser 1265	1270	1275
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp 1280	1285	1290
Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser 1295	1300	1305
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp 1310	1315	1320
Ser Asp Ala Gly Lys His Thr Pro Val Lys Pro Met Ser Thr Thr 1325	1330	1335
Lys Asp His His Asn Lys Ala Lys Ala Leu Pro Glu Thr Gly Asn 1340	1345	1350
Glu Asn Ser Gly Ser Asn Asn Ala Thr Leu Phe Gly Gly Leu Phe 1355	1360	1365
Ala Ala Leu Gly Ser Leu Leu Leu Phe Gly Arg Arg Lys Lys Gln 1370	1375	1380
Asn Lys 1385		

&lt;210&gt; SEQ ID NO 14

&lt;211&gt; LENGTH: 1141

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 14

Met Ile Asn Arg Asp Asn Lys Lys Ala Ile Thr Lys Lys Gly Met Ile  
1 5 10 15

Ser Asn Arg Leu Asn Lys Phe Ser Ile Arg Lys Tyr Thr Val Gly Thr  
20 25 30

Ala Ser Ile Leu Val Gly Thr Thr Leu Ile Phe Gly Leu Gly Asn Gln  
35 40 45

-continued

---

Glu Ala Lys Ala Ala Glu Asn Thr Ser Thr Glu Asn Ala Lys Gln Asp  
 50 55 60  
 Asp Ala Thr Thr Ser Asp Asn Lys Glu Val Val Ser Glu Thr Glu Asn  
 65 70 75 80  
 Asn Ser Thr Thr Glu Asn Asp Ser Thr Asn Pro Ile Lys Lys Glu Thr  
 85 90 95  
 Asn Thr Asp Ser Gln Pro Glu Ala Lys Glu Glu Ser Thr Thr Ser Ser  
 100 105 110  
 Thr Gln Gln Gln Gln Asn Asn Val Thr Ala Thr Thr Glu Thr Lys Pro  
 115 120 125  
 Gln Asn Ile Glu Lys Glu Asn Val Lys Pro Ser Thr Asp Lys Thr Ala  
 130 135 140  
 Thr Glu Asp Thr Ser Val Ile Leu Glu Glu Lys Lys Ala Pro Asn Tyr  
 145 150 155 160  
 Thr Asn Asn Asp Val Thr Thr Lys Pro Ser Thr Ser Glu Ile Gln Thr  
 165 170 175  
 Lys Pro Thr Thr Pro Gln Glu Ser Thr Asn Ile Glu Asn Ser Gln Pro  
 180 185 190  
 Gln Pro Thr Pro Ser Lys Val Asp Asn Gln Val Thr Asp Ala Thr Asn  
 195 200 205  
 Pro Lys Glu Pro Val Asn Val Ser Lys Glu Glu Leu Lys Asn Asn Pro  
 210 215 220  
 Glu Lys Leu Lys Glu Leu Val Arg Asn Asp Asn Asn Thr Asp Arg Ser  
 225 230 235 240  
 Thr Lys Pro Val Ala Thr Ala Pro Thr Ser Val Ala Pro Lys Arg Leu  
 245 250 255  
 Asn Ala Lys Met Arg Phe Ala Val Ala Gln Pro Ala Ala Val Ala Ser  
 260 265 270  
 Asn Asn Val Asn Asp Leu Ile Thr Val Thr Lys Gln Thr Ile Lys Val  
 275 280 285  
 Gly Asp Gly Lys Asp Asn Val Ala Ala Ala His Asp Gly Lys Asp Ile  
 290 295 300  
 Glu Tyr Asp Thr Glu Phe Thr Ile Asp Asn Lys Val Lys Lys Gly Asp  
 305 310 315 320  
 Thr Met Thr Ile Asn Tyr Asp Lys Asn Val Ile Pro Ser Asp Leu Thr  
 325 330 335  
 Asp Lys Asn Asp Pro Ile Asp Ile Thr Asp Pro Ser Gly Glu Val Ile  
 340 345 350  
 Ala Lys Gly Thr Phe Asp Lys Ala Thr Lys Gln Ile Thr Tyr Thr Phe  
 355 360 365  
 Thr Asp Tyr Val Asp Lys Tyr Glu Asp Ile Lys Ala Arg Leu Thr Leu  
 370 375 380  
 Tyr Ser Tyr Ile Asp Lys Gln Ala Val Pro Asn Glu Thr Ser Leu Asn  
 385 390 395 400  
 Leu Thr Phe Ala Thr Ala Gly Lys Glu Thr Ser Gln Asn Val Ser Val  
 405 410 415  
 Asp Tyr Gln Asp Pro Met Val His Gly Asp Ser Asn Ile Gln Ser Ile  
 420 425 430  
 Phe Thr Lys Leu Asp Glu Asn Lys Gln Thr Ile Glu Gln Gln Ile Tyr  
 435 440 445  
 Val Asn Pro Leu Lys Lys Thr Ala Thr Asn Thr Lys Val Asp Ile Ala  
 450 455 460  
 Gly Ser Gln Val Asp Asp Tyr Gly Asn Ile Lys Leu Gly Asn Gly Ser



-continued

---

Gln Thr Val Thr Asn Thr Thr Glu Asp Asp Lys Asp Ala Asp Gly Gly  
                   900                                  905                                  910  
 Glu Val Asp Val Thr Ile Thr Asp His Asp Asp Phe Thr Leu Asp Asn  
                   915                                  920                                  925  
 Gly Tyr Phe Glu Glu Asp Thr Ser Asp Ser Asp Ser Asp Ser Asp Ser  
                   930                                  935                                  940  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
                   945                                  950                                  955                                  960  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
                   965                                  970                                  975  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
                   980                                  985                                  990  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
                   995                                  1000                                  1005  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp  
                   1010                                  1015                                  1020  
 Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
                   1025                                  1030                                  1035  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp  
                   1040                                  1045                                  1050  
 Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
                   1055                                  1060                                  1065  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ala Gly  
                   1070                                  1075                                  1080  
 Lys His Thr Pro Val Lys Pro Met Ser Thr Thr Lys Asp His His  
                   1085                                  1090                                  1095  
 Asn Lys Ala Lys Ala Leu Pro Glu Thr Gly Ser Glu Asn Asn Gly  
                   1100                                  1105                                  1110  
 Ser Asn Asn Ala Thr Leu Phe Gly Gly Leu Phe Ala Ala Leu Gly  
                   1115                                  1120                                  1125  
 Ser Leu Leu Leu Phe Gly Arg Arg Lys Lys Gln Asn Lys  
                   1130                                  1135                                  1140

&lt;210&gt; SEQ ID NO 15

&lt;211&gt; LENGTH: 350

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 15

Met Thr Lys His Tyr Leu Asn Ser Lys Tyr Gln Ser Glu Gln Arg Ser  
 1                  5                                  10                                  15  
 Ser Ala Met Lys Lys Ile Thr Met Gly Thr Ala Ser Ile Ile Leu Gly  
                   20                                  25                                  30  
 Ser Leu Val Tyr Ile Gly Ala Asp Ser Gln Gln Val Asn Ala Ala Thr  
                   35                                  40                                  45  
 Glu Ala Thr Asn Ala Thr Asn Asn Gln Ser Thr Gln Val Ser Gln Ala  
                   50                                  55                                  60  
 Thr Ser Gln Pro Ile Asn Phe Gln Val Gln Lys Asp Gly Ser Ser Glu  
 65                  70                                  75                                  80  
 Lys Ser His Met Asp Asp Tyr Met Gln His Pro Gly Lys Val Ile Lys  
                   85                                  90                                  95  
 Gln Asn Asn Lys Tyr Tyr Phe Gln Thr Val Leu Asn Asn Ala Ser Phe  
                   100                                  105                                  110  
 Trp Lys Glu Tyr Lys Phe Tyr Asn Ala Asn Asn Gln Glu Leu Ala Thr

-continued

115					120					125					
Thr	Val	Val	Asn	Asp	Asn	Lys	Lys	Ala	Asp	Thr	Arg	Thr	Ile	Asn	Val
130					135						140				
Ala	Val	Glu	Pro	Gly	Tyr	Lys	Ser	Leu	Thr	Thr	Lys	Val	His	Ile	Val
145				150						155					160
Val	Pro	Gln	Ile	Asn	Tyr	Asn	His	Arg	Tyr	Thr	Thr	His	Leu	Glu	Phe
				165					170						175
Glu	Lys	Ala	Ile	Pro	Thr	Leu	Ala	Asp	Ala	Ala	Lys	Pro	Asn	Asn	Val
			180					185						190	
Lys	Pro	Val	Gln	Pro	Lys	Pro	Ala	Gln	Pro	Lys	Thr	Pro	Thr	Glu	Gln
		195					200						205		
Thr	Lys	Pro	Val	Gln	Pro	Lys	Val	Glu	Lys	Val	Lys	Pro	Thr	Val	Thr
	210					215					220				
Thr	Thr	Ser	Lys	Val	Glu	Asp	Asn	His	Ser	Thr	Lys	Val	Val	Ser	Thr
	225				230					235					240
Asp	Thr	Thr	Lys	Asp	Gln	Thr	Lys	Thr	Gln	Thr	Ala	His	Thr	Val	Lys
			245						250						255
Thr	Ala	Gln	Thr	Ala	Gln	Glu	Gln	Asn	Lys	Val	Gln	Thr	Pro	Val	Lys
		260						265					270		
Asp	Val	Ala	Thr	Ala	Lys	Ser	Glu	Ser	Asn	Asn	Gln	Ala	Val	Ser	Asp
		275					280						285		
Asn	Lys	Ser	Gln	Gln	Thr	Asn	Lys	Val	Thr	Lys	His	Asn	Glu	Thr	Pro
	290					295					300				
Lys	Gln	Ala	Ser	Lys	Ala	Lys	Glu	Leu	Pro	Lys	Thr	Gly	Leu	Thr	Ser
	305				310					315					320
Val	Asp	Asn	Phe	Ile	Ser	Thr	Val	Ala	Phe	Ala	Thr	Leu	Ala	Leu	Leu
			325						330						335
Gly	Ser	Leu	Ser	Leu	Leu	Leu	Phe	Lys	Arg	Lys	Glu	Ser	Lys		
		340						345					350		

&lt;210&gt; SEQ ID NO 16

&lt;211&gt; LENGTH: 645

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 16

Met	Asn	Lys	Gln	Gln	Lys	Glu	Phe	Lys	Ser	Phe	Tyr	Ser	Ile	Arg	Lys
1			5						10					15	
Ser	Ser	Leu	Gly	Val	Ala	Ser	Val	Ala	Ile	Ser	Thr	Leu	Leu	Leu	Leu
		20						25					30		
Met	Ser	Asn	Gly	Glu	Ala	Gln	Ala	Ala	Ala	Glu	Glu	Thr	Gly	Gly	Thr
	35					40						45			
Asn	Thr	Glu	Ala	Gln	Pro	Lys	Thr	Glu	Ala	Val	Ala	Ser	Pro	Thr	Thr
	50					55					60				
Thr	Ser	Glu	Lys	Ala	Pro	Glu	Thr	Lys	Pro	Val	Ala	Asn	Ala	Val	Ser
	65				70					75					80
Val	Ser	Asn	Lys	Glu	Val	Glu	Ala	Pro	Thr	Ser	Glu	Thr	Lys	Glu	Ala
			85						90						95
Lys	Glu	Val	Lys	Glu	Val	Lys	Ala	Pro	Lys	Glu	Thr	Lys	Ala	Val	Lys
		100						105					110		
Pro	Ala	Ala	Lys	Ala	Thr	Asn	Asn	Thr	Tyr	Pro	Ile	Leu	Asn	Gln	Glu
		115					120							125	
Leu	Arg	Glu	Ala	Ile	Lys	Asn	Pro	Ala	Ile	Lys	Asp	Lys	Asp	His	Ser
	130					135							140		

-continued

---

Ala Pro Asn Ser Arg Pro Ile Asp Phe Glu Met Lys Lys Glu Asn Gly  
145 150 155 160

Glu Gln Gln Phe Tyr His Tyr Ala Ser Ser Val Lys Pro Ala Arg Val  
165 170 175

Ile Phe Thr Asp Ser Lys Pro Glu Ile Glu Leu Gly Leu Gln Ser Gly  
180 185 190

Gln Phe Trp Arg Lys Phe Glu Val Tyr Glu Gly Asp Lys Lys Leu Pro  
195 200 205

Ile Lys Leu Val Ser Tyr Asp Thr Val Lys Asp Tyr Ala Tyr Ile Arg  
210 215 220

Phe Ser Val Ser Asn Gly Thr Lys Ala Val Lys Ile Val Ser Ser Thr  
225 230 235 240

His Phe Asn Asn Lys Glu Glu Lys Tyr Asp Tyr Thr Leu Met Glu Phe  
245 250 255

Ala Gln Pro Ile Tyr Asn Ser Ala Asp Lys Phe Lys Thr Glu Glu Asp  
260 265 270

Tyr Lys Ala Glu Lys Leu Leu Ala Pro Tyr Lys Lys Ala Lys Thr Leu  
275 280 285

Glu Arg Gln Val Tyr Glu Leu Asn Lys Ile Gln Asp Lys Leu Pro Glu  
290 295 300

Lys Leu Lys Ala Glu Tyr Lys Lys Lys Leu Glu Asp Thr Lys Lys Ala  
305 310 315 320

Leu Asp Glu Gln Val Lys Ser Ala Ile Thr Glu Phe Gln Asn Val Gln  
325 330 335

Pro Thr Asn Glu Lys Met Thr Asp Leu Gln Asp Thr Lys Tyr Val Val  
340 345 350

Tyr Glu Ser Val Glu Asn Asn Glu Ser Met Met Asp Thr Phe Val Lys  
355 360 365

His Pro Ile Lys Thr Gly Met Leu Asn Gly Lys Lys Tyr Met Val Met  
370 375 380

Glu Thr Thr Asn Asp Asp Tyr Trp Lys Asp Phe Met Val Glu Gly Gln  
385 390 395 400

Arg Val Arg Thr Ile Ser Lys Asp Ala Lys Asn Asn Thr Arg Thr Ile  
405 410 415

Ile Phe Pro Tyr Val Glu Gly Lys Thr Leu Tyr Asp Ala Ile Val Lys  
420 425 430

Val His Val Lys Thr Ile Asp Tyr Asp Gly Gln Tyr His Val Arg Ile  
435 440 445

Val Asp Lys Glu Ala Phe Thr Lys Ala Asn Thr Asp Lys Ser Asn Lys  
450 455 460

Lys Glu Gln Gln Asp Asn Ser Ala Lys Lys Glu Ala Thr Pro Ala Thr  
465 470 475 480

Pro Ser Lys Pro Thr Pro Ser Pro Val Glu Lys Glu Ser Gln Lys Gln  
485 490 495

Asp Ser Gln Lys Asp Asp Asn Lys Gln Leu Pro Ser Val Glu Lys Glu  
500 505 510

Asn Asp Ala Ser Ser Glu Ser Gly Lys Asp Lys Thr Pro Ala Thr Lys  
515 520 525

Pro Thr Lys Gly Glu Val Glu Ser Ser Ser Thr Thr Pro Thr Lys Val  
530 535 540

Val Ser Thr Thr Gln Asn Val Ala Lys Pro Thr Thr Ala Ser Ser Lys  
545 550 555 560

Thr Thr Lys Asp Val Val Gln Thr Ser Ala Gly Ser Ser Glu Ala Lys





-continued

Gly Ser Asp Ser Asp Ser Gly Ser Asp Ser Asp Ser Glu Ser Asp Ser  
 595 600 605  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Glu Ser  
 610 615 620  
 Asp Ser Asp Ser Glu Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 625 630 635 640  
 Asp Ser Asp Ser Asp Ser Glu Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 645 650 655  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Glu Ser Asp Ser Asp Ser Glu Ser  
 660 665 670  
 Asp Ser Glu Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 675 680 685  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 690 695 700  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Glu Ser Asp Ser Asp Ser Asp Ser  
 705 710 715 720  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 725 730 735  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 740 745 750  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 755 760 765  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 770 775 780  
 Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser  
 785 790 795 800  
 Asp Ser Asp Ser Arg Val Thr Pro Pro Asn Asn Glu Gln Lys Ala Pro  
 805 810 815  
 Ser Asn Pro Lys Gly Glu Val Asn His Ser Asn Lys Val Ser Lys Gln  
 820 825 830  
 His Lys Thr Asp Ala Leu Pro Glu Thr Gly Asp Lys Ser Glu Asn Thr  
 835 840 845  
 Asn Ala Thr Leu Phe Gly Ala Met Met Ala Leu Leu Gly Ser Leu Leu  
 850 855 860  
 Leu Phe Arg Lys Arg Lys Gln Asp His Lys Glu Lys Ala  
 865 870 875

&lt;210&gt; SEQ ID NO 19

&lt;211&gt; LENGTH: 227

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 19

Met Lys Asn Ile Leu Lys Val Phe Asn Thr Thr Ile Leu Ala Leu Ile  
 1 5 10 15  
 Ile Ile Ile Ala Thr Phe Ser Asn Ser Ala Asn Ala Ala Asp Ser Gly  
 20 25 30  
 Thr Leu Asn Tyr Glu Val Tyr Lys Tyr Asn Thr Asn Asp Thr Ser Ile  
 35 40 45  
 Ala Asn Asp Tyr Phe Asn Lys Pro Ala Lys Tyr Ile Lys Lys Asn Gly  
 50 55 60  
 Lys Leu Tyr Val Gln Ile Thr Val Asn His Ser His Trp Ile Thr Gly  
 65 70 75 80  
 Met Ser Ile Glu Gly His Lys Glu Asn Ile Ile Ser Lys Asn Thr Ala



-continued

---

Ser Gly His Val Val Leu Asn Lys Phe Leu Ser Asn Glu Glu Asn Gln  
 225 230 235 240  
 Ser His Ser Asn Gln Leu Thr Asp Lys Leu Gln Gly Ser Asp Lys Ile  
 245 250 255  
 Asn His Ala Met Ile Glu Lys Leu Ala Lys Ser Asn Ala Ser Thr Gln  
 260 265 270  
 His Tyr Thr Tyr His Lys Leu Asn Thr Leu Gln Ser Leu Asp Gln Arg  
 275 280 285  
 Ile Ala Asn Thr Gln Leu Pro Lys Asn Gln Lys Ser Asp Leu Met Ser  
 290 295 300  
 Glu Val Asn Lys Thr Lys Glu Arg Ile Lys Ser Gln Arg Asn Ile Ile  
 305 310 315 320  
 Leu Glu Glu Leu Ala Arg Thr Asp Asp Lys Lys Tyr Ala Thr Gln Ser  
 325 330 335  
 Ile Leu Glu Ser Ile Phe Asn Lys Asp Glu Ala Asp Lys Ile Leu Lys  
 340 345 350  
 Asp Ile Arg Val Asp Gly Lys Thr Asp Gln Gln Ile Ala Asp Gln Ile  
 355 360 365  
 Thr Arg His Ile Asp Gln Leu Ser Leu Thr Thr Ser Asp Asp Leu Leu  
 370 375 380  
 Thr Ser Leu Ile Asp Gln Ser Gln Asp Lys Ser Leu Leu Ile Ser Gln  
 385 390 395 400  
 Ile Leu Gln Thr Lys Leu Gly Lys Ala Glu Ala Asp Lys Leu Ala Lys  
 405 410 415  
 Asp Trp Thr Asn Lys Gly Leu Ser Asn Arg Gln Ile Val Asp Gln Leu  
 420 425 430  
 Lys Lys His Phe Ala Ser Thr Gly Asp Thr Ser Ser Asp Asp Ile Leu  
 435 440 445  
 Lys Ala Ile Leu Asn Asn Ala Lys Asp Lys Lys Gln Ala Ile Glu Thr  
 450 455 460  
 Ile Leu Ala Thr Arg Ile Glu Arg Gln Lys Ala Lys Leu Leu Ala Asp  
 465 470 475 480  
 Leu Ile Thr Lys Ile Glu Thr Asp Gln Asn Lys Ile Phe Asn Leu Val  
 485 490 495  
 Lys Ser Ala Leu Asn Gly Lys Ala Asp Asp Leu Leu Asn Leu Gln Lys  
 500 505 510  
 Arg Leu Asn Gln Thr Lys Lys Asp Ile Asp Tyr Ile Leu Ser Pro Ile  
 515 520 525  
 Val Asn Arg Pro Ser Leu Leu Asp Arg Leu Asn Lys Asn Gly Lys Thr  
 530 535 540  
 Thr Asp Leu Asn Lys Leu Ala Asn Leu Met Asn Gln Gly Ser Asn Leu  
 545 550 555 560  
 Leu Asp Ser Ile Pro Asp Ile Pro Thr Pro Lys Pro Glu Lys Thr Leu  
 565 570 575  
 Thr Leu Gly Lys Gly Asn Gly Leu Leu Ser Gly Leu Leu Asn Ala Asp  
 580 585 590  
 Gly Asn Val Ser Leu Pro Lys Ala Gly Glu Thr Ile Lys Glu His Trp  
 595 600 605  
 Leu Pro Ile Ser Val Ile Val Gly Ala Met Gly Val Leu Met Ile Trp  
 610 615 620  
 Leu Ser Arg Arg Asn Lys Leu Lys Asn Lys Ala  
 625 630 635

-continued

---

```

<210> SEQ ID NO 21
<211> LENGTH: 953
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 21

Met Asn Asn Lys Lys Thr Ala Thr Asn Arg Lys Gly Met Ile Pro Asn
 1          5          10          15

Arg Leu Asn Lys Phe Ser Ile Arg Lys Tyr Ser Val Gly Thr Ala Ser
          20          25          30

Ile Leu Val Gly Thr Thr Leu Ile Phe Gly Leu Ser Gly His Glu Ala
          35          40          45

Lys Ala Ala Glu His Thr Asn Gly Glu Leu Asn Gln Ser Lys Asn Glu
          50          55          60

Thr Thr Ala Pro Ser Glu Asn Lys Thr Thr Glu Lys Val Asp Ser Arg
65          70          75          80

Gln Leu Lys Asp Asn Thr Gln Thr Ala Thr Ala Asp Gln Pro Lys Val
          85          90          95

Thr Met Ser Asp Ser Ala Thr Val Lys Glu Thr Ser Ser Asn Met Gln
          100          105          110

Ser Pro Gln Asn Ala Thr Ala Ser Gln Ser Thr Thr Gln Thr Ser Asn
          115          120          125

Val Thr Thr Asn Asp Lys Ser Ser Thr Thr Tyr Ser Asn Glu Thr Asp
          130          135          140

Lys Ser Asn Leu Thr Gln Ala Lys Asn Val Ser Thr Thr Pro Lys Thr
          145          150          155          160

Thr Thr Ile Lys Gln Arg Ala Leu Asn Arg Met Ala Val Asn Thr Val
          165          170          175

Ala Ala Pro Gln Gln Gly Thr Asn Val Asn Asp Lys Val His Phe Thr
          180          185          190

Asn Ile Asp Ile Ala Ile Asp Lys Gly His Val Asn Lys Thr Thr Gly
          195          200          205

Asn Thr Glu Phe Trp Ala Thr Ser Ser Asp Val Leu Lys Leu Lys Ala
          210          215          220

Asn Tyr Thr Ile Asp Asp Ser Val Lys Glu Gly Asp Thr Phe Thr Phe
          225          230          235          240

Lys Tyr Gly Gln Tyr Phe Arg Pro Gly Ser Val Arg Leu Pro Ser Gln
          245          250          255

Thr Gln Asn Leu Tyr Asn Ala Gln Gly Asn Ile Ile Ala Lys Gly Ile
          260          265          270

Tyr Asp Ser Lys Thr Asn Thr Thr Thr Tyr Thr Phe Thr Asn Tyr Val
          275          280          285

Asp Gln Tyr Thr Asn Val Ser Gly Ser Phe Glu Gln Val Ala Phe Ala
          290          295          300

Lys Arg Glu Asn Ala Thr Thr Asp Lys Thr Ala Tyr Lys Met Glu Val
          305          310          315          320

Thr Leu Gly Asn Asp Thr Tyr Ser Lys Asp Val Ile Val Asp Tyr Gly
          325          330          335

Asn Gln Lys Gly Gln Gln Leu Ile Ser Ser Thr Asn Tyr Ile Asn Asn
          340          345          350

Glu Asp Leu Ser Arg Asn Met Thr Val Tyr Val Asn Gln Pro Lys Lys
          355          360          365

Thr Tyr Thr Lys Glu Thr Phe Val Thr Asn Leu Thr Gly Tyr Lys Phe
          370          375          380

```

-continued

Asn	Pro	Asp	Ala	Lys	Asn	Phe	Lys	Ile	Tyr	Glu	Val	Thr	Asp	Gln	Asn	385	390	395	400
Gln	Phe	Val	Asp	Ser	Phe	Thr	Pro	Asp	Thr	Ser	Lys	Leu	Lys	Asp	Val	405	410	415	
Thr	Gly	Gln	Phe	Asp	Val	Ile	Tyr	Ser	Asn	Asp	Asn	Lys	Thr	Ala	Thr	420	425	430	
Val	Asp	Leu	Leu	Asn	Gly	Gln	Ser	Ser	Ser	Asp	Lys	Gln	Tyr	Ile	Ile	435	440	445	
Gln	Gln	Val	Ala	Tyr	Pro	Asp	Asn	Ser	Ser	Thr	Asp	Asn	Gly	Lys	Ile	450	455	460	
Asp	Tyr	Thr	Leu	Glu	Thr	Gln	Asn	Gly	Lys	Ser	Ser	Trp	Ser	Asn	Ser	465	470	475	480
Tyr	Ser	Asn	Val	Asn	Gly	Ser	Ser	Thr	Ala	Asn	Gly	Asp	Gln	Lys	Lys	485	490	495	
Tyr	Asn	Leu	Gly	Asp	Tyr	Val	Trp	Glu	Asp	Thr	Asn	Lys	Asp	Gly	Lys	500	505	510	
Gln	Asp	Ala	Asn	Glu	Lys	Gly	Ile	Lys	Gly	Val	Tyr	Val	Ile	Leu	Lys	515	520	525	
Asp	Ser	Asn	Gly	Lys	Glu	Leu	Asp	Arg	Thr	Thr	Thr	Asp	Glu	Asn	Gly	530	535	540	
Lys	Tyr	Gln	Phe	Thr	Gly	Leu	Ser	Asn	Gly	Thr	Tyr	Ser	Val	Glu	Phe	545	550	555	560
Ser	Thr	Pro	Ala	Gly	Tyr	Thr	Pro	Thr	Thr	Ala	Asn	Ala	Gly	Thr	Asp	565	570	575	
Asp	Ala	Val	Asp	Ser	Asp	Gly	Leu	Thr	Thr	Thr	Gly	Val	Ile	Lys	Asp	580	585	590	
Ala	Asp	Asn	Met	Thr	Leu	Asp	Ser	Gly	Phe	Tyr	Lys	Thr	Pro	Lys	Tyr	595	600	605	
Ser	Leu	Gly	Asp	Tyr	Val	Trp	Tyr	Asp	Ser	Asn	Lys	Asp	Gly	Lys	Gln	610	615	620	
Asp	Ser	Thr	Glu	Lys	Gly	Ile	Lys	Gly	Val	Lys	Val	Thr	Leu	Gln	Asn	625	630	635	640
Glu	Lys	Gly	Glu	Val	Ile	Gly	Thr	Thr	Glu	Thr	Asp	Glu	Asn	Gly	Lys	645	650	655	
Tyr	Arg	Phe	Asp	Asn	Leu	Asp	Ser	Gly	Lys	Tyr	Lys	Val	Ile	Phe	Glu	660	665	670	
Lys	Pro	Ala	Gly	Leu	Thr	Gln	Thr	Gly	Thr	Asn	Thr	Thr	Glu	Asp	Asp	675	680	685	
Lys	Asp	Ala	Asp	Gly	Gly	Glu	Val	Asp	Val	Thr	Ile	Thr	Asp	His	Asp	690	695	700	
Asp	Phe	Thr	Leu	Asp	Asn	Gly	Tyr	Tyr	Glu	Glu	Glu	Thr	Ser	Asp	Ser	705	710	715	720
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	725	730	735	
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	740	745	750	
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	755	760	765	
Asp	Ser	Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	770	775	780	
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	785	790	795	800
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser				

-continued

---

	805		810		815
Asp Ser Asp Ser Asp Ser Asp Asn Asp Ser Asp Ser Asp Ser Asp Ser	820		825		830
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser	835		840		845
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser	850		855		860
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser	865		870		875
Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ala Gly Lys	885		890		895
His Thr Pro Thr Lys Pro Met Ser Thr Val Lys Asp Gln His Lys Thr	900		905		910
Ala Lys Ala Leu Pro Glu Thr Gly Ser Glu Asn Asn Asn Ser Asn Asn	915		920		925
Gly Thr Leu Phe Gly Gly Leu Phe Ala Ala Leu Gly Ser Leu Leu Leu	930		935		940
Phe Gly Arg Arg Lys Lys Gln Asn Lys	945		950		

&lt;210&gt; SEQ ID NO 22

&lt;211&gt; LENGTH: 989

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 22

Met Asn Met Lys Lys Lys Glu Lys His Ala Ile Arg Lys Lys Ser Ile	1		5		10		15
Gly Val Ala Ser Val Leu Val Gly Thr Leu Ile Gly Phe Gly Leu Leu		20		25		30	
Ser Ser Lys Glu Ala Asp Ala Ser Glu Asn Ser Val Thr Gln Ser Asp		35		40		45	
Ser Ala Ser Asn Glu Ser Lys Ser Asn Asp Ser Ser Ser Val Ser Ala		50		55		60	
Ala Pro Lys Thr Asp Asp Thr Asn Val Ser Asp Thr Lys Thr Ser Ser		65		70		75	
Asn Thr Asn Asn Gly Glu Thr Ser Val Ala Gln Asn Pro Ala Gln Gln		85		90		95	
Glu Thr Thr Gln Ser Ser Ser Thr Asn Ala Thr Thr Glu Glu Thr Pro		100		105		110	
Val Thr Gly Glu Ala Thr Thr Thr Thr Thr Asn Gln Ala Asn Thr Pro		115		120		125	
Ala Thr Thr Gln Ser Ser Asn Thr Asn Ala Glu Glu Leu Val Asn Gln		130		135		140	
Thr Ser Asn Glu Thr Thr Ser Asn Asp Thr Asn Thr Val Ser Ser Val		145		150		155	
Asn Ser Pro Gln Asn Ser Thr Asn Ala Glu Asn Val Ser Thr Thr Gln		165		170		175	
Asp Thr Ser Thr Glu Ala Thr Pro Ser Asn Asn Glu Ser Ala Pro Gln		180		185		190	
Asn Thr Asp Ala Ser Asn Lys Asp Val Val Ser Gln Ala Val Asn Pro		195		200		205	
Ser Thr Pro Arg Met Arg Ala Phe Ser Leu Ala Ala Val Ala Ala Asp		210		215		220	

-continued

---

Ala Pro Ala Ala Gly Thr Asp Ile Thr Asn Gln Leu Thr Asp Val Lys  
225 230 235 240

Val Thr Ile Asp Ser Gly Thr Thr Val Tyr Pro His Gln Ala Gly Tyr  
245 250 255

Val Lys Leu Asn Tyr Gly Phe Ser Val Pro Asn Ser Ala Val Lys Gly  
260 265 270

Asp Thr Phe Lys Ile Thr Val Pro Lys Glu Leu Asn Leu Asn Gly Val  
275 280 285

Thr Ser Thr Ala Lys Val Pro Pro Ile Met Ala Gly Asp Gln Val Leu  
290 295 300

Ala Asn Gly Val Ile Asp Ser Asp Gly Asn Val Ile Tyr Thr Phe Thr  
305 310 315 320

Asp Tyr Val Asp Asn Lys Glu Asn Val Thr Ala Asn Ile Thr Met Pro  
325 330 335

Ala Tyr Ile Asp Pro Glu Asn Val Thr Lys Thr Gly Asn Val Thr Leu  
340 345 350

Thr Thr Gly Ile Gly Thr Asn Thr Ala Ser Lys Thr Val Leu Ile Asp  
355 360 365

Tyr Glu Lys Tyr Gly Gln Phe His Asn Leu Ser Ile Lys Gly Thr Ile  
370 375 380

Asp Gln Ile Asp Lys Thr Asn Asn Thr Tyr Arg Gln Thr Ile Tyr Val  
385 390 395 400

Asn Pro Ser Gly Asp Asn Val Val Leu Pro Ala Leu Thr Gly Asn Leu  
405 410 415

Ile Pro Asn Thr Lys Ser Asn Ala Leu Ile Asp Ala Lys Asn Thr Asp  
420 425 430

Ile Lys Val Tyr Arg Val Asp Asn Ala Asn Asp Leu Ser Glu Ser Tyr  
435 440 445

Tyr Val Asn Pro Ser Asp Phe Glu Asp Val Thr Asn Gln Val Arg Ile  
450 455 460

Ser Phe Pro Asn Ala Asn Gln Tyr Lys Val Glu Phe Pro Thr Asp Asp  
465 470 475 480

Asp Gln Ile Thr Thr Pro Tyr Ile Val Val Val Asn Gly His Ile Asp  
485 490 495

Pro Ala Ser Thr Gly Asp Leu Ala Leu Arg Ser Thr Phe Tyr Gly Tyr  
500 505 510

Asp Ser Asn Phe Ile Trp Arg Ser Met Ser Trp Asp Asn Glu Val Ala  
515 520 525

Phe Asn Asn Gly Ser Gly Ser Gly Asp Gly Ile Asp Lys Pro Val Val  
530 535 540

Pro Glu Gln Pro Asp Glu Pro Gly Glu Ile Glu Pro Ile Pro Glu Asp  
545 550 555 560

Ser Asp Ser Asp Pro Gly Ser Asp Ser Gly Ser Asp Ser Asn Ser Asp  
565 570 575

Ser Gly Ser Asp Ser Gly Ser Asp Ser Thr Ser Asp Ser Gly Ser Asp  
580 585 590

Ser Ala Ser Asp Ser Asp Ser Ala Ser Asp Ser Asp Ser Ala Ser Asp  
595 600 605

Ser Asp Ser Ala Ser Asp Ser Asp Ser Ala Ser Asp Ser Asp Ser Ala  
610 615 620

Ser Asp Ser Asp Ser Ala Ser Asp Ser Asp Ser Ala Ser Asp Ser Asp  
625 630 635 640

Ser Ala Ser Asp Ser Asp Ser Ala Ser Asp Ser Asp Ser Ala Ser Asp

-continued

645				650				655							
Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Asp
		660					665						670		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
		675				680							685		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
		690				695					700				
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
		705				710					715				720
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			725								730				735
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			740				745						750		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			755				760						765		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			770				775						780		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			785				790						795		800
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			805								810				815
Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Glu
			820								825				830
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			835				840						845		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			850				855						860		
Ser	Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			865				870						875		880
Ser	Ala	Ser	Asp	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Asp	Ser	Ser	Ser	Asp
			885										890		895
Ser	Asp	Ser	Asp	Ser	Thr	Ser	Asp	Thr	Gly	Ser	Asp	Asn	Asp	Ser	Asp
			900										905		910
Ser	Asp	Ser	Asn	Ser	Asp	Ser	Glu	Ser	Gly	Ser	Asn	Asn	Asn	Val	Val
			915				920						925		
Pro	Pro	Asn	Ser	Pro	Lys	Asn	Gly	Thr	Asn	Ala	Ser	Asn	Lys	Asn	Glu
			930				935						940		
Ala	Lys	Asp	Ser	Lys	Glu	Pro	Leu	Pro	Asp	Thr	Gly	Ser	Glu	Asp	Glu
			945				950						955		960
Ala	Asn	Thr	Ser	Leu	Ile	Trp	Gly	Leu	Leu	Ala	Ser	Leu	Gly	Ser	Leu
			965										970		975
Leu	Leu	Phe	Arg	Arg	Lys	Lys	Glu	Asn	Lys	Asp	Lys	Lys			
			980										985		

&lt;210&gt; SEQ ID NO 23

&lt;211&gt; LENGTH: 584

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 23

Met Lys Phe Lys Ser Leu Ile Thr Thr Thr Leu Ala Leu Gly Val Leu  
1 5 10 15

Ala Ser Thr Gly Ala Asn Phe Asn Asn Asn Glu Ala Ser Ala Ala Ala  
20 25 30





-continued

---

Gln Ala Val Ala Leu Ile Arg Thr Met Pro Gln Arg Gln Gln Thr Ser  
 245 250 255  
 Arg Arg Ser Asn Arg Ile Gln Thr Arg Ser Val Glu Ser Arg Ala Ala  
 260 265 270  
 Glu Pro Arg Ser Val Ser Asp Tyr Gln Asn Ala Asn Ser Ser Tyr Tyr  
 275 280 285  
 Val Glu Asn Ala Asn Asp Gly Ser Gly Tyr Pro Val Gly Thr Tyr Ile  
 290 295 300  
 Asn Ala Ser Ser Lys Gly Ala Pro Tyr Asn Leu Pro Thr Thr Pro Trp  
 305 310 315 320  
 Asn Thr Leu Lys Ala Ser Asp Ser Lys Glu Ile Ala Leu Met Thr Ala  
 325 330 335  
 Lys Gln Thr Gly Asp Gly Tyr Gln Trp Val Ile Lys Phe Asn Lys Gly  
 340 345 350  
 His Ala Pro His Gln Asn Met Ile Phe Trp Phe Ala Leu Pro Ala Asp  
 355 360 365  
 Gln Val Pro Val Gly Arg Thr Asp Phe Val Thr Val Asn Ser Asp Gly  
 370 375 380  
 Thr Asn Val Gln Trp Ser His Gly Ala Gly Ala Asn Lys Pro  
 385 390 395 400  
 Leu Gln Gln Met Trp Glu Tyr Gly Val Asn Asp Pro His Arg Ser His  
 405 410 415  
 Asp Phe Lys Ile Arg Asn Arg Ser Gly Gln Val Ile Tyr Asp Trp Pro  
 420 425 430  
 Thr Val His Ile Tyr Ser Leu Glu Asp Leu Ser Arg Ala Ser Asp Tyr  
 435 440 445  
 Phe Ser Glu Ala Gly Ala Thr Pro Ala Thr Lys Ala Phe Gly Arg Gln  
 450 455 460  
 Asn Phe Glu Tyr Ile Asn Gly Gln Lys Pro Ala Glu Ser Pro Gly Val  
 465 470 475 480  
 Pro Lys Val Tyr Thr Phe Ile Gly Gln Gly Asp Ala Ser Tyr Thr Ile  
 485 490 495  
 Ser Phe Lys Thr Gln Gly Pro Thr Val Asn Lys Leu Tyr Tyr Ala Ala  
 500 505 510  
 Gly Gly Arg Ala Leu Glu Tyr Asn Gln Leu Phe Met Tyr Ser Gln Leu  
 515 520 525  
 Tyr Val Glu Ser Thr Gln Asp His Gln Gln Arg Leu Asn Gly Leu Arg  
 530 535 540  
 Gln Val Val Asn Arg Thr Tyr Arg Ile Gly Thr Thr Lys Arg Val Glu  
 545 550 555 560  
 Val Ser Gln Gly Asn Val Gln Thr Lys Lys Val Leu Glu Ser Thr Asn  
 565 570 575  
 Leu Asn Ile Asp Asp Phe Val Asp Asp Pro Leu Ser Tyr Val Lys Thr  
 580 585 590  
 Pro Ser Asn Lys Val Leu Gly Phe Tyr Ser Asn Asn Ala Asn Thr Asn  
 595 600 605  
 Ala Phe Arg Pro Gly Gly Ala Gln Gln Leu Asn Glu Tyr Gln Leu Ser  
 610 615 620  
 Gln Leu Phe Thr Asp Gln Lys Leu Gln Glu Ala Ala Arg Thr Arg Asn  
 625 630 635 640  
 Pro Ile Arg Leu Met Ile Gly Phe Asp Tyr Pro Asp Ala Tyr Gly Asn  
 645 650 655  
 Ser Glu Thr Leu Val Pro Val Asn Leu Thr Val Leu Pro Glu Ile Gln

-continued

660				665				670							
His	Asn	Ile	Lys	Phe	Phe	Lys	Asn	Asp	Asp	Thr	Gln	Asn	Ile	Ala	Glu
	675						680						685		
Lys	Pro	Phe	Ser	Lys	Gln	Ala	Gly	His	Pro	Val	Phe	Tyr	Val	Tyr	Ala
	690					695					700				
Gly	Asn	Gln	Gly	Asn	Ala	Ser	Val	Asn	Leu	Gly	Gly	Ser	Val	Thr	Ser
	705				710					715					720
Ile	Gln	Pro	Leu	Arg	Ile	Asn	Leu	Thr	Ser	Asn	Glu	Asn	Phe	Thr	Asp
				725					730					735	
Lys	Asp	Trp	Gln	Ile	Thr	Gly	Ile	Pro	Arg	Thr	Leu	His	Ile	Glu	Asn
			740						745				750		
Ser	Thr	Asn	Arg	Pro	Asn	Asn	Ala	Arg	Glu	Arg	Asn	Ile	Glu	Leu	Val
		755					760						765		
Gly	Asn	Leu	Leu	Pro	Gly	Asp	Tyr	Phe	Gly	Thr	Ile	Arg	Phe	Gly	Arg
	770					775					780				
Lys	Glu	Gln	Leu	Phe	Glu	Ile	Arg	Val	Lys	Pro	His	Thr	Pro	Thr	Ile
	785				790					795					800
Thr	Thr	Thr	Ala	Glu	Gln	Leu	Arg	Gly	Thr	Ala	Leu	Gln	Lys	Val	Pro
			805						810					815	
Val	Asn	Ile	Ser	Gly	Ile	Pro	Leu	Asp	Pro	Ser	Ala	Leu	Val	Tyr	Leu
			820						825					830	
Val	Ala	Pro	Thr	Asn	Gln	Thr	Thr	Asn	Gly	Gly	Ser	Glu	Ala	Asp	Gln
		835					840						845		
Ile	Pro	Ser	Gly	Tyr	Thr	Ile	Leu	Ala	Thr	Gly	Thr	Pro	Asp	Gly	Val
	850					855					860				
His	Asn	Thr	Ile	Thr	Ile	Arg	Pro	Gln	Asp	Tyr	Val	Val	Phe	Ile	Pro
	865				870					875					880
Pro	Val	Gly	Lys	Gln	Ile	Arg	Ala	Val	Val	Tyr	Tyr	Asn	Lys	Val	Val
			885							890				895	
Ala	Ser	Asn	Met	Ser	Asn	Ala	Val	Thr	Ile	Leu	Pro	Asp	Asp	Ile	Pro
		900							905					910	
Pro	Thr	Ile	Asn	Asn	Pro	Val	Gly	Ile	Asn	Ala	Lys	Tyr	Tyr	Arg	Gly
		915				920							925		
Asp	Glu	Val	Asn	Phe	Thr	Met	Gly	Val	Ser	Asp	Arg	His	Ser	Gly	Ile
	930					935					940				
Lys	Asn	Thr	Thr	Ile	Thr	Thr	Leu	Pro	Asn	Gly	Trp	Thr	Ser	Asn	Leu
	945				950					955					960
Thr	Lys	Ala	Asp	Lys	Asn	Asn	Gly	Ser	Leu	Ser	Ile	Thr	Gly	Arg	Val
			965						970					975	
Ser	Met	Asn	Gln	Ala	Phe	Asn	Ser	Asp	Ile	Thr	Phe	Lys	Val	Ser	Ala
		980							985					990	
Thr	Asp	Asn	Val	Asn	Asn	Thr	Thr	Asn	Asp	Ser	Gln	Ser	Lys	His	Val
		995				1000							1005		
Ser	Ile	His	Val	Gly	Lys	Ile	Ser	Glu	Asp	Ala	His	Pro	Ile	Val	
	1010					1015							1020		
Leu	Gly	Asn	Thr	Glu	Lys	Val	Val	Val	Val	Asn	Pro	Thr	Ala	Val	
	1025					1030							1035		
Ser	Asn	Asp	Glu	Lys	Gln	Ser	Ile	Ile	Thr	Ala	Phe	Met	Asn	Lys	
	1040					1045							1050		
Asn	Gln	Asn	Ile	Arg	Gly	Tyr	Leu	Ala	Ser	Thr	Asp	Pro	Val	Thr	
	1055					1060							1065		
Val	Asp	Asn	Asn	Gly	Asn	Val	Thr	Leu	His	Tyr	Arg	Asp	Gly	Ser	
	1070					1075							1080		

-continued

---

Ser Thr	Thr Leu Asp Ala Thr	Asn Val Met Thr Tyr	Glu Pro Val
1085	1090	1095	
Val Lys	Pro Glu Tyr Gln Thr	Val Asn Ala Ala Lys	Thr Ala Thr
1100	1105	1110	
Val Thr	Ile Ala Lys Gly Gln	Ser Phe Ser Ile Gly	Asp Ile Lys
1115	1120	1125	
Gln Tyr	Phe Thr Leu Ser Asn	Gly Gln Pro Ile Pro	Ser Gly Thr
1130	1135	1140	
Phe Thr	Asn Ile Thr Ser Asp	Arg Thr Ile Pro Thr	Ala Gln Glu
1145	1150	1155	
Val Ser	Gln Met Asn Ala Gly	Thr Gln Leu Tyr His	Ile Thr Ala
1160	1165	1170	
Thr Asn	Ala Tyr His Lys Asp	Ser Glu Asp Phe Tyr	Ile Ser Leu
1175	1180	1185	
Lys Ile	Ile Asp Val Lys Gln	Pro Glu Gly Asp Gln	Arg Val Tyr
1190	1195	1200	
Arg Thr	Ser Thr Tyr Asp Leu	Thr Thr Asp Glu Ile	Ser Lys Val
1205	1210	1215	
Lys Gln	Ala Phe Ile Asn Ala	Asn Arg Asp Val Ile	Thr Leu Ala
1220	1225	1230	
Glu Gly	Asp Ile Ser Val Thr	Asn Thr Pro Asn Gly	Ala Asn Val
1235	1240	1245	
Ser Thr	Ile Thr Val Asn Ile	Asn Lys Gly Arg Leu	Thr Lys Ser
1250	1255	1260	
Phe Ala	Ser Asn Leu Ala Asn	Met Asn Phe Leu Arg	Trp Val Asn
1265	1270	1275	
Phe Pro	Gln Asp Tyr Thr Val	Thr Trp Thr Asn Ala	Lys Ile Ala
1280	1285	1290	
Asn Arg	Pro Thr Asp Gly Gly	Leu Ser Trp Ser Asp	Asp His Lys
1295	1300	1305	
Ser Leu	Ile Tyr Arg Tyr Asp	Ala Thr Leu Gly Thr	Gln Ile Thr
1310	1315	1320	
Thr Asn	Asp Ile Leu Thr Met	Leu Lys Ala Thr Thr	Thr Val Pro
1325	1330	1335	
Gly Leu	Arg Asn Asn Ile Thr	Gly Asn Glu Lys Ser	Gln Ala Glu
1340	1345	1350	
Ala Gly	Gly Arg Pro Asn Phe	Arg Thr Thr Gly Tyr	Ser Gln Ser
1355	1360	1365	
Asn Ala	Thr Thr Asp Gly Gln	Arg Gln Phe Thr Leu	Asn Gly Gln
1370	1375	1380	
Val Ile	Gln Val Leu Asp Ile	Ile Asn Pro Ser Asn	Gly Tyr Gly
1385	1390	1395	
Gly Gln	Pro Val Thr Asn Ser	Asn Thr Arg Ala Asn	His Ser Asn
1400	1405	1410	
Ser Thr	Val Val Asn Val Asn	Glu Pro Ala Ala Asn	Gly Ala Gly
1415	1420	1425	
Ala Phe	Thr Ile Asp His Val	Val Lys Ser Asn Ser	Thr His Asn
1430	1435	1440	
Ala Ser	Asp Ala Val Tyr Lys	Ala Gln Leu Tyr Leu	Thr Pro Tyr
1445	1450	1455	
Gly Pro	Lys Gln Tyr Val Glu	His Leu Asn Gln Asn	Thr Gly Asn
1460	1465	1470	

-continued

Thr	Thr	Asp	Ala	Ile	Asn	Ile	Tyr	Phe	Val	Pro	Ser	Asp	Leu	Val
1475						1480					1485			
Asn	Pro	Thr	Ile	Ser	Val	Gly	Asn	Tyr	Thr	Asn	His	Gln	Val	Phe
1490						1495					1500			
Ser	Gly	Glu	Thr	Phe	Thr	Asn	Thr	Ile	Thr	Ala	Asn	Asp	Asn	Phe
1505						1510					1515			
Gly	Val	Gln	Ser	Val	Thr	Val	Pro	Asn	Thr	Ser	Gln	Ile	Thr	Gly
1520						1525					1530			
Thr	Val	Asp	Asn	Asn	His	Gln	His	Val	Ser	Ala	Thr	Ala	Pro	Asn
1535						1540					1545			
Val	Thr	Ser	Ala	Thr	Asn	Lys	Thr	Ile	Asn	Leu	Leu	Ala	Thr	Asp
1550						1555					1560			
Thr	Ser	Gly	Asn	Thr	Ala	Thr	Thr	Ser	Phe	Asn	Val	Thr	Val	Lys
1565						1570					1575			
Pro	Leu	Arg	Asp	Lys	Tyr	Arg	Val	Gly	Thr	Ser	Ser	Thr	Ala	Ala
1580						1585					1590			
Asn	Pro	Val	Arg	Ile	Ala	Asn	Ile	Ser	Asn	Asn	Ala	Thr	Val	Ser
1595						1600					1605			
Gln	Ala	Asp	Gln	Thr	Thr	Ile	Ile	Asn	Ser	Leu	Thr	Phe	Thr	Glu
1610						1615					1620			
Thr	Val	Pro	Asn	Arg	Ser	Tyr	Ala	Arg	Ala	Ser	Ala	Asn	Glu	Ile
1625						1630					1635			
Thr	Ser	Lys	Thr	Val	Ser	Asn	Val	Ser	Arg	Thr	Gly	Asn	Asn	Ala
1640						1645					1650			
Asn	Val	Thr	Val	Thr	Val	Thr	Tyr	Gln	Asp	Gly	Thr	Thr	Ser	Thr
1655						1660					1665			
Val	Thr	Val	Pro	Val	Lys	His	Val	Ile	Pro	Glu	Ile	Val	Ala	His
1670						1675					1680			
Ser	His	Tyr	Thr	Val	Gln	Gly	Gln	Asp	Phe	Pro	Ala	Gly	Asn	Gly
1685						1690					1695			
Ser	Ser	Ala	Ser	Asp	Tyr	Phe	Lys	Leu	Ser	Asn	Gly	Ser	Asp	Ile
1700						1705					1710			
Ala	Asp	Ala	Thr	Ile	Thr	Trp	Val	Ser	Gly	Gln	Ala	Pro	Asn	Lys
1715						1720					1725			
Asp	Asn	Thr	Arg	Ile	Gly	Glu	Asp	Ile	Thr	Val	Thr	Ala	His	Ile
1730						1735					1740			
Leu	Ile	Asp	Gly	Glu	Thr	Thr	Pro	Ile	Thr	Lys	Thr	Ala	Thr	Tyr
1745						1750					1755			
Lys	Val	Val	Arg	Thr	Val	Pro	Lys	His	Val	Phe	Glu	Thr	Ala	Arg
1760						1765					1770			
Gly	Val	Leu	Tyr	Pro	Gly	Val	Ser	Asp	Met	Tyr	Asp	Ala	Lys	Gln
1775						1780					1785			
Tyr	Val	Lys	Pro	Val	Asn	Asn	Ser	Trp	Ser	Thr	Asn	Ala	Gln	His
1790						1795					1800			
Met	Asn	Phe	Gln	Phe	Val	Gly	Thr	Tyr	Gly	Pro	Asn	Lys	Asp	Val
1805						1810					1815			
Val	Gly	Ile	Ser	Thr	Arg	Leu	Ile	Arg	Val	Thr	Tyr	Asp	Asn	Arg
1820						1825					1830			
Gln	Thr	Glu	Asp	Leu	Thr	Ile	Leu	Ser	Lys	Val	Lys	Pro	Asp	Pro
1835						1840					1845			
Pro	Arg	Ile	Asp	Ala	Asn	Ser	Val	Thr	Tyr	Lys	Ala	Gly	Leu	Thr
1850						1855					1860			
Asn	Gln	Glu	Ile	Lys	Val	Asn	Asn	Val	Leu	Asn	Asn	Ser	Ser	Val

-continued

---

1865	1870	1875
Lys Leu Phe Lys Ala Asp Asn Thr Pro Leu Asn Val Thr Asn Ile		
1880	1885	1890
Thr His Gly Ser Gly Phe Ser Ser Val Val Thr Val Ser Asp Ala		
1895	1900	1905
Leu Pro Asn Gly Gly Ile Lys Ala Lys Ser Ser Ile Ser Met Asn		
1910	1915	1920
Asn Val Thr Tyr Thr Thr Gln Asp Glu His Gly Gln Val Val Thr		
1925	1930	1935
Val Thr Arg Asn Glu Ser Val Asp Ser Asn Asp Ser Ala Thr Val		
1940	1945	1950
Thr Val Thr Pro Gln Leu Gln Ala Thr Thr Glu Gly Ala Val Phe		
1955	1960	1965
Ile Lys Gly Gly Asp Gly Phe Asp Phe Gly His Val Glu Arg Phe		
1970	1975	1980
Ile Gln Asn Pro Pro His Gly Ala Thr Val Ala Trp His Asp Ser		
1985	1990	1995
Pro Asp Thr Trp Lys Asn Thr Val Gly Asn Thr His Lys Thr Ala		
2000	2005	2010
Val Val Thr Leu Pro Asn Gly Gln Gly Thr Arg Asn Val Glu Val		
2015	2020	2025
Pro Val Lys Val Tyr Pro Val Ala Asn Ala Lys Ala Pro Ser Arg		
2030	2035	2040
Asp Val Lys Gly Gln Asn Leu Thr Asn Gly Thr Asp Ala Met Asn		
2045	2050	2055
Tyr Ile Thr Phe Asp Pro Asn Thr Asn Thr Asn Gly Ile Thr Ala		
2060	2065	2070
Ala Trp Ala Asn Arg Gln Gln Pro Asn Asn Gln Gln Ala Gly Val		
2075	2080	2085
Gln His Leu Asn Val Asp Val Thr Tyr Pro Gly Ile Ser Ala Ala		
2090	2095	2100
Lys Arg Val Pro Val Thr Val Asn Val Tyr Gln Phe Glu Phe Pro		
2105	2110	2115
Gln Thr Thr Tyr Thr Thr Thr Val Gly Gly Thr Leu Ala Ser Gly		
2120	2125	2130
Thr Gln Ala Ser Gly Tyr Ala His Met Gln Asn Ala Thr Gly Leu		
2135	2140	2145
Pro Thr Asp Gly Phe Thr Tyr Lys Trp Asn Arg Asp Thr Thr Gly		
2150	2155	2160
Thr Asn Asp Ala Asn Trp Ser Ala Met Asn Lys Pro Asn Val Ala		
2165	2170	2175
Lys Val Val Asn Ala Lys Tyr Asp Val Ile Tyr Asn Gly His Thr		
2180	2185	2190
Phe Ala Thr Ser Leu Pro Ala Lys Phe Val Val Lys Asp Val Gln		
2195	2200	2205
Pro Ala Lys Pro Thr Val Thr Glu Thr Ala Ala Gly Ala Ile Thr		
2210	2215	2220
Ile Ala Pro Gly Ala Asn Gln Thr Val Asn Thr His Ala Gly Asn		
2225	2230	2235
Val Thr Thr Tyr Ala Asp Lys Leu Val Ile Lys Arg Asn Gly Asn		
2240	2245	2250
Val Val Thr Thr Phe Thr Arg Arg Asn Asn Thr Ser Pro Trp Val		
2255	2260	2265

-continued

---

Lys	Glu	Ala	Ser	Ala	Ala	Thr	Val	Ala	Gly	Ile	Ala	Gly	Thr	Asn
2270						2275					2280			
Asn	Gly	Ile	Thr	Val	Ala	Ala	Gly	Thr	Phe	Asn	Pro	Ala	Asp	Thr
2285						2290					2295			
Ile	Gln	Val	Val	Ala	Thr	Gln	Gly	Ser	Gly	Glu	Thr	Val	Ser	Asp
2300						2305					2310			
Glu	Gln	Arg	Ser	Asp	Asp	Phe	Thr	Val	Val	Ala	Pro	Gln	Pro	Asn
2315						2320					2325			
Gln	Ala	Thr	Thr	Lys	Ile	Trp	Gln	Asn	Gly	His	Ile	Asp	Ile	Thr
2330						2335					2340			
Pro	Asn	Asn	Pro	Ser	Gly	His	Leu	Ile	Asn	Pro	Thr	Gln	Ala	Met
2345						2350					2355			
Asp	Ile	Ala	Tyr	Thr	Glu	Lys	Val	Gly	Asn	Gly	Ala	Glu	His	Ser
2360						2365					2370			
Lys	Thr	Ile	Asn	Val	Val	Arg	Gly	Gln	Asn	Asn	Gln	Trp	Thr	Ile
2375						2380					2385			
Ala	Asn	Lys	Pro	Asp	Tyr	Val	Thr	Leu	Asp	Ala	Gln	Thr	Gly	Lys
2390						2395					2400			
Val	Thr	Phe	Asn	Ala	Asn	Thr	Ile	Lys	Pro	Asn	Ser	Ser	Ile	Thr
2405						2410					2415			
Ile	Thr	Pro	Lys	Ala	Gly	Thr	Gly	His	Ser	Val	Ser	Ser	Asn	Pro
2420						2425					2430			
Ser	Thr	Leu	Thr	Ala	Pro	Ala	Ala	His	Thr	Val	Asn	Thr	Thr	Glu
2435						2440					2445			
Ile	Val	Lys	Asp	Tyr	Gly	Ser	Asn	Val	Thr	Ala	Ala	Glu	Ile	Asn
2450						2455					2460			
Asn	Ala	Val	Gln	Val	Ala	Asn	Lys	Arg	Thr	Ala	Thr	Ile	Lys	Asn
2465						2470					2475			
Gly	Thr	Ala	Met	Pro	Thr	Asn	Leu	Ala	Gly	Gly	Ser	Thr	Thr	Thr
2480						2485					2490			
Ile	Pro	Val	Thr	Val	Thr	Tyr	Asn	Asp	Gly	Ser	Thr	Glu	Glu	Val
2495						2500					2505			
Gln	Glu	Ser	Ile	Phe	Thr	Lys	Ala	Asp	Lys	Arg	Glu	Leu	Ile	Thr
2510						2515					2520			
Ala	Lys	Asn	His	Leu	Asp	Asp	Pro	Val	Ser	Thr	Glu	Gly	Lys	Lys
2525						2530					2535			
Pro	Gly	Thr	Ile	Thr	Gln	Tyr	Asn	Asn	Ala	Met	His	Asn	Ala	Gln
2540						2545					2550			
Gln	Gln	Ile	Asn	Thr	Ala	Lys	Thr	Glu	Ala	Gln	Gln	Val	Ile	Asn
2555						2560					2565			
Asn	Glu	Arg	Ala	Thr	Pro	Gln	Gln	Val	Ser	Asp	Ala	Leu	Thr	Lys
2570						2575					2580			
Val	Arg	Ala	Ala	Gln	Thr	Lys	Ile	Asp	Gln	Ala	Lys	Ala	Leu	Leu
2585						2590					2595			
Gln	Asn	Lys	Glu	Asp	Asn	Ser	Gln	Leu	Val	Thr	Ser	Lys	Asn	Asn
2600						2605					2610			
Leu	Gln	Ser	Ser	Val	Asn	Gln	Val	Pro	Ser	Thr	Ala	Gly	Met	Thr
2615						2620					2625			
Gln	Gln	Ser	Ile	Asp	Asn	Tyr	Asn	Ala	Lys	Lys	Arg	Glu	Ala	Glu
2630						2635					2640			
Thr	Glu	Ile	Thr	Ala	Ala	Gln	Arg	Val	Ile	Asp	Asn	Gly	Asp	Ala
2645						2650					2655			



-continued

---

3050	3055	3060
Gln Lys Leu Thr Glu Ile Asn	Gln Val Leu Asn Gly Asn Pro Thr	
3065	3070	3075
Val Gln Asn Ile Asn Asp Lys	Val Thr Glu Ala Asn Gln Ala Lys	
3080	3085	3090
Asp Gln Leu Asn Thr Ala Arg	Gln Gly Leu Thr Leu Asp Arg Gln	
3095	3100	3105
Pro Ala Leu Thr Thr Leu His	Gly Ala Ser Asn Leu Asn Gln Ala	
3110	3115	3120
Gln Gln Asn Asn Phe Thr Gln	Gln Ile Asn Ala Ala Gln Asn His	
3125	3130	3135
Ala Ala Leu Glu Thr Ile Lys	Ser Asn Ile Thr Ala Leu Asn Thr	
3140	3145	3150
Ala Met Thr Lys Leu Lys Asp	Ser Val Ala Asp Asn Asn Thr Ile	
3155	3160	3165
Lys Ser Asp Gln Asn Tyr Thr	Asp Ala Thr Pro Ala Asn Lys Gln	
3170	3175	3180
Ala Tyr Asp Asn Ala Val Asn	Ala Ala Lys Gly Val Ile Gly Glu	
3185	3190	3195
Thr Thr Asn Pro Thr Met Asp	Val Asn Thr Val Asn Gln Lys Ala	
3200	3205	3210
Ala Ser Val Lys Ser Thr Lys	Asp Ala Leu Asp Gly Gln Gln Asn	
3215	3220	3225
Leu Gln Arg Ala Lys Thr Glu	Ala Thr Asn Ala Ile Thr His Ala	
3230	3235	3240
Ser Asp Leu Asn Gln Ala Gln	Lys Asn Ala Leu Thr Gln Gln Val	
3245	3250	3255
Asn Ser Ala Gln Asn Val Gln	Ala Val Asn Asp Ile Lys Gln Thr	
3260	3265	3270
Thr Gln Ser Leu Asn Thr Ala	Met Thr Gly Leu Lys Arg Gly Val	
3275	3280	3285
Ala Asn His Asn Gln Val Val	Gln Ser Asp Asn Tyr Val Asn Ala	
3290	3295	3300
Asp Thr Asn Lys Lys Asn Asp	Tyr Asn Asn Ala Tyr Asn His Ala	
3305	3310	3315
Asn Asp Ile Ile Asn Gly Asn	Ala Gln His Pro Val Ile Thr Pro	
3320	3325	3330
Ser Asp Val Asn Asn Ala Leu	Ser Asn Val Thr Ser Lys Glu His	
3335	3340	3345
Ala Leu Asn Gly Glu Ala Lys	Leu Asn Ala Ala Lys Gln Glu Ala	
3350	3355	3360
Asn Thr Ala Leu Gly His Leu	Asn Asn Leu Asn Asn Ala Gln Arg	
3365	3370	3375
Gln Asn Leu Gln Ser Gln Ile	Asn Gly Ala His Gln Ile Asp Ala	
3380	3385	3390
Val Asn Thr Ile Lys Gln Asn	Ala Thr Asn Leu Asn Ser Ala Met	
3395	3400	3405
Gly Asn Leu Arg Gln Ala Val	Ala Asp Lys Asp Gln Val Lys Arg	
3410	3415	3420
Thr Glu Asp Tyr Ala Asp Ala	Asp Thr Ala Lys Gln Asn Ala Tyr	
3425	3430	3435
Asn Ser Ala Val Ser Ser Ala	Glu Thr Ile Ile Asn Gln Thr Thr	
3440	3445	3450

-continued

---

Asn Pro	Thr Met Ser Val	Asp	Asp Val Asn Arg	Ala	Thr Ser Ala
3455		3460		3465	
Val Thr	Ser Asn Lys Asn	Ala	Leu Asn Gly Tyr	Glu	Lys Leu Ala
3470		3475		3480	
Gln Ser	Lys Thr Asp Ala	Ala	Arg Ala Ile Asp	Ala	Leu Pro His
3485		3490		3495	
Leu Asn	Asn Ala Gln Lys	Ala	Asp Val Lys Ser	Lys	Ile Asn Ala
3500		3505		3510	
Ala Ser	Asn Ile Ala Gly	Val	Asn Thr Val Lys	Gln	Gln Gly Thr
3515		3520		3525	
Asp Leu	Asn Thr Ala Met	Gly	Asn Leu Gln Gly	Ala	Ile Asn Asp
3530		3535		3540	
Glu Gln	Thr Thr Leu Asn	Ser	Gln Asn Tyr Gln	Asp	Ala Thr Pro
3545		3550		3555	
Ser Lys	Lys Thr Ala Tyr	Thr	Asn Ala Val Gln	Ala	Ala Lys Asp
3560		3565		3570	
Ile Leu	Asn Lys Ser Asn	Gly	Gln Asn Lys Thr	Lys	Asp Gln Val
3575		3580		3585	
Thr Glu	Ala Met Asn Gln	Val	Asn Ser Ala Lys	Asn	Asn Leu Asp
3590		3595		3600	
Gly Thr	Arg Leu Leu Asp	Gln	Ala Lys Gln Thr	Ala	Lys Gln Gln
3605		3610		3615	
Leu Asn	Asn Met Thr His	Leu	Thr Thr Ala Gln	Lys	Thr Asn Leu
3620		3625		3630	
Thr Asn	Gln Ile Asn Ser	Gly	Thr Thr Val Ala	Gly	Val Gln Thr
3635		3640		3645	
Val Gln	Ser Asn Ala Asn	Thr	Leu Asp Gln Ala	Met	Asn Thr Leu
3650		3655		3660	
Arg Gln	Ser Ile Ala Asn	Lys	Asp Ala Thr Lys	Ala	Ser Glu Asp
3665		3670		3675	
Tyr Val	Asp Ala Asn Asn	Asp	Lys Gln Thr Ala	Tyr	Asn Asn Ala
3680		3685		3690	
Val Ala	Ala Ala Glu Thr	Ile	Ile Asn Ala Asn	Ser	Asn Pro Glu
3695		3700		3705	
Met Asn	Pro Ser Thr Ile	Thr	Gln Lys Ala Glu	Gln	Val Asn Ser
3710		3715		3720	
Ser Lys	Thr Ala Leu Asn	Gly	Asp Glu Asn Leu	Ala	Ala Ala Lys
3725		3730		3735	
Gln Asn	Ala Lys Thr Tyr	Leu	Asn Thr Leu Thr	Ser	Ile Thr Asp
3740		3745		3750	
Ala Gln	Lys Asn Asn Leu	Ile	Ser Gln Ile Thr	Ser	Ala Thr Arg
3755		3760		3765	
Val Ser	Gly Val Asp Thr	Val	Lys Gln Asn Ala	Gln	His Leu Asp
3770		3775		3780	
Gln Ala	Met Ala Ser Leu	Gln	Asn Gly Ile Asn	Asn	Glu Ser Gln
3785		3790		3795	
Val Lys	Ser Ser Glu Lys	Tyr	Arg Asp Ala Asp	Thr	Asn Lys Gln
3800		3805		3810	
Gln Glu	Tyr Asp Asn Ala	Ile	Thr Ala Ala Lys	Ala	Ile Leu Asn
3815		3820		3825	
Lys Ser	Thr Gly Pro Asn	Thr	Ala Gln Asn Ala	Val	Glu Ala Ala
3830		3835		3840	

-continued

Leu	Gln	Arg	Val	Asn	Asn	Ala	Lys	Asp	Ala	Leu	Asn	Gly	Asp	Ala
3845						3850					3855			
Lys	Leu	Ile	Ala	Ala	Gln	Asn	Ala	Ala	Lys	Gln	His	Leu	Gly	Thr
3860						3865					3870			
Leu	Thr	His	Ile	Thr	Thr	Ala	Gln	Arg	Asn	Asp	Leu	Thr	Asn	Gln
3875						3880					3885			
Ile	Ser	Gln	Ala	Thr	Asn	Leu	Ala	Gly	Val	Glu	Ser	Val	Lys	Gln
3890						3895					3900			
Asn	Ala	Asn	Ser	Leu	Asp	Gly	Ala	Met	Gly	Asn	Leu	Gln	Thr	Ala
3905						3910					3915			
Ile	Asn	Asp	Lys	Ser	Gly	Thr	Leu	Ala	Ser	Gln	Asn	Phe	Leu	Asp
3920						3925					3930			
Ala	Asp	Glu	Gln	Lys	Arg	Asn	Ala	Tyr	Asn	Gln	Ala	Val	Ser	Ala
3935						3940					3945			
Ala	Glu	Thr	Ile	Leu	Asn	Lys	Gln	Thr	Gly	Pro	Asn	Thr	Ala	Lys
3950						3955					3960			
Thr	Ala	Val	Glu	Gln	Ala	Leu	Asn	Asn	Val	Asn	Asn	Ala	Lys	His
3965						3970					3975			
Ala	Leu	Asn	Gly	Thr	Gln	Asn	Leu	Asn	Asn	Ala	Lys	Gln	Ala	Ala
3980						3985					3990			
Ile	Thr	Ala	Ile	Asn	Gly	Ala	Ser	Asp	Leu	Asn	Gln	Lys	Gln	Lys
3995						4000					4005			
Asp	Ala	Leu	Lys	Ala	Gln	Ala	Asn	Gly	Ala	Gln	Arg	Val	Ser	Asn
4010						4015					4020			
Ala	Gln	Asp	Val	Gln	His	Asn	Ala	Thr	Glu	Leu	Asn	Thr	Ala	Met
4025						4030					4035			
Gly	Thr	Leu	Lys	His	Ala	Ile	Ala	Asp	Lys	Thr	Asn	Thr	Leu	Ala
4040						4045					4050			
Ser	Ser	Lys	Tyr	Val	Asn	Ala	Asp	Ser	Thr	Lys	Gln	Asn	Ala	Tyr
4055						4060					4065			
Thr	Thr	Lys	Val	Thr	Asn	Ala	Glu	His	Ile	Ile	Ser	Gly	Thr	Pro
4070						4075					4080			
Thr	Val	Val	Thr	Thr	Pro	Ser	Glu	Val	Thr	Ala	Ala	Ala	Asn	Gln
4085						4090					4095			
Val	Asn	Ser	Ala	Lys	Gln	Glu	Leu	Asn	Gly	Asp	Glu	Arg	Leu	Arg
4100						4105					4110			
Glu	Ala	Lys	Gln	Asn	Ala	Asn	Thr	Ala	Ile	Asp	Ala	Leu	Thr	Gln
4115						4120					4125			
Leu	Asn	Thr	Pro	Gln	Lys	Ala	Lys	Leu	Lys	Glu	Gln	Val	Gly	Gln
4130						4135					4140			
Ala	Asn	Arg	Leu	Glu	Asp	Val	Gln	Thr	Val	Gln	Thr	Asn	Gly	Gln
4145						4150					4155			
Ala	Leu	Asn	Asn	Ala	Met	Lys	Gly	Leu	Arg	Asp	Ser	Ile	Ala	Asn
4160						4165					4170			
Glu	Thr	Thr	Val	Lys	Thr	Ser	Gln	Asn	Tyr	Thr	Asp	Ala	Ser	Pro
4175						4180					4185			
Asn	Asn	Gln	Ser	Thr	Tyr	Asn	Ser	Ala	Val	Ser	Asn	Ala	Lys	Gly
4190						4195					4200			
Ile	Ile	Asn	Gln	Thr	Asn	Asn	Pro	Thr	Met	Asp	Thr	Ser	Ala	Ile
4205						4210					4215			
Thr	Gln	Ala	Thr	Thr	Gln	Val	Asn	Asn	Ala	Lys	Asn	Gly	Leu	Asn
4220						4225					4230			
Gly	Ala	Glu	Asn	Leu	Arg	Asn	Ala	Gln	Asn	Thr	Ala	Lys	Gln	Asn



-continued

---

Asp	Ala	Leu	Lys	Ala	Gln	Val	Thr	Ser	Ala	Gln	Arg	Val	Ala	Asn
4640						4645					4650			
Val	Thr	Ser	Ile	Gln	Gln	Thr	Ala	Asn	Glu	Leu	Asn	Thr	Ala	Met
4655						4660					4665			
Gly	Gln	Leu	Gln	His	Gly	Ile	Asp	Asp	Glu	Asn	Ala	Thr	Lys	Gln
4670						4675					4680			
Thr	Gln	Lys	Tyr	Arg	Asp	Ala	Glu	Gln	Ser	Lys	Lys	Thr	Ala	Tyr
4685						4690					4695			
Asp	Gln	Ala	Val	Ala	Ala	Ala	Lys	Ala	Ile	Leu	Asn	Lys	Gln	Thr
4700						4705					4710			
Gly	Ser	Asn	Ser	Asp	Lys	Ala	Ala	Val	Asp	Arg	Ala	Leu	Gln	Gln
4715						4720					4725			
Val	Thr	Ser	Thr	Lys	Asp	Ala	Leu	Asn	Gly	Asp	Ala	Lys	Leu	Ala
4730						4735					4740			
Glu	Ala	Lys	Ala	Ala	Ala	Lys	Gln	Asn	Leu	Gly	Thr	Leu	Asn	His
4745						4750					4755			
Ile	Thr	Asn	Ala	Gln	Arg	Thr	Asp	Leu	Glu	Gly	Gln	Ile	Asn	Gln
4760						4765					4770			
Ala	Thr	Thr	Val	Asp	Gly	Val	Asn	Thr	Val	Lys	Thr	Asn	Ala	Asn
4775						4780					4785			
Thr	Leu	Asp	Gly	Ala	Met	Asn	Ser	Leu	Gln	Gly	Ser	Ile	Asn	Asp
4790						4795					4800			
Lys	Asp	Ala	Thr	Leu	Arg	Asn	Gln	Asn	Tyr	Leu	Asp	Ala	Asp	Glu
4805						4810					4815			
Ser	Lys	Arg	Asn	Ala	Tyr	Thr	Gln	Ala	Val	Thr	Ala	Ala	Glu	Gly
4820						4825					4830			
Ile	Leu	Asn	Lys	Gln	Thr	Gly	Gly	Asn	Thr	Ser	Lys	Ala	Asp	Val
4835						4840					4845			
Asp	Asn	Ala	Leu	Asn	Ala	Val	Thr	Arg	Ala	Lys	Ala	Ala	Leu	Asn
4850						4855					4860			
Gly	Ala	Asp	Asn	Leu	Arg	Asn	Ala	Lys	Thr	Ser	Ala	Thr	Asn	Thr
4865						4870					4875			
Ile	Asp	Gly	Leu	Pro	Asn	Leu	Thr	Gln	Leu	Gln	Lys	Asp	Asn	Leu
4880						4885					4890			
Lys	His	Gln	Val	Glu	Gln	Ala	Gln	Asn	Val	Ala	Gly	Val	Asn	Gly
4895						4900					4905			
Val	Lys	Asp	Lys	Gly	Asn	Thr	Leu	Asn	Thr	Ala	Met	Gly	Ala	Leu
4910						4915					4920			
Arg	Thr	Ser	Ile	Gln	Asn	Asp	Asn	Thr	Thr	Lys	Thr	Ser	Gln	Asn
4925						4930					4935			
Tyr	Leu	Asp	Ala	Ser	Asp	Ser	Asn	Lys	Asn	Asn	Tyr	Asn	Thr	Ala
4940						4945					4950			
Val	Asn	Asn	Ala	Asn	Gly	Val	Ile	Asn	Ala	Thr	Asn	Asn	Pro	Asn
4955						4960					4965			
Met	Asp	Ala	Asn	Ala	Ile	Asn	Gly	Met	Ala	Asn	Gln	Val	Asn	Thr
4970						4975					4980			
Thr	Lys	Ala	Ala	Leu	Asn	Gly	Ala	Gln	Asn	Leu	Ala	Gln	Ala	Lys
4985						4990					4995			
Thr	Asn	Ala	Thr	Asn	Thr	Ile	Asn	Asn	Ala	His	Asp	Leu	Asn	Gln
5000						5005					5010			
Lys	Gln	Lys	Asp	Ala	Leu	Lys	Thr	Gln	Val	Asn	Asn	Ala	Gln	Arg
5015						5020					5025			

-continued

Val	Ser	Asp	Ala	Asn	Asn	Val	Gln	His	Thr	Ala	Thr	Glu	Leu	Asn
5030						5035						5040		
Ser	Ala	Met	Thr	Ala	Leu	Lys	Ala	Ala	Ile	Ala	Asp	Lys	Glu	Arg
5045						5050					5055			
Thr	Lys	Ala	Ser	Gly	Asn	Tyr	Val	Asn	Ala	Asp	Gln	Glu	Lys	Arg
5060						5065					5070			
Gln	Ala	Tyr	Asp	Ser	Lys	Val	Thr	Asn	Ala	Glu	Asn	Ile	Ile	Ser
5075						5080					5085			
Gly	Thr	Pro	Asn	Ala	Thr	Leu	Thr	Val	Asn	Asp	Val	Asn	Ser	Ala
5090						5095					5100			
Ala	Ser	Gln	Val	Asn	Ala	Ala	Lys	Thr	Ala	Leu	Asn	Gly	Asp	Asn
5105						5110					5115			
Asn	Leu	Arg	Val	Ala	Lys	Glu	His	Ala	Asn	Asn	Thr	Ile	Asp	Gly
5120						5125					5130			
Leu	Ala	Gln	Leu	Asn	Asn	Ala	Gln	Lys	Ala	Lys	Leu	Lys	Glu	Gln
5135						5140					5145			
Val	Gln	Ser	Ala	Thr	Thr	Leu	Asp	Gly	Val	Gln	Thr	Val	Lys	Asn
5150						5155					5160			
Ser	Ser	Gln	Thr	Leu	Asn	Thr	Ala	Met	Lys	Gly	Leu	Arg	Asp	Ser
5165						5170					5175			
Ile	Ala	Asn	Glu	Ala	Thr	Ile	Lys	Ala	Gly	Gln	Asn	Tyr	Thr	Asp
5180						5185					5190			
Ala	Ser	Pro	Asn	Asn	Arg	Asn	Glu	Tyr	Asp	Ser	Ala	Val	Thr	Ala
5195						5200					5205			
Ala	Lys	Ala	Ile	Ile	Asn	Gln	Thr	Ser	Asn	Pro	Thr	Met	Glu	Pro
5210						5215					5220			
Asn	Thr	Ile	Thr	Gln	Val	Thr	Ser	Gln	Val	Thr	Thr	Lys	Glu	Gln
5225						5230					5235			
Ala	Leu	Asn	Gly	Ala	Arg	Asn	Leu	Ala	Gln	Ala	Lys	Thr	Thr	Ala
5240						5245					5250			
Lys	Asn	Asn	Leu	Asn	Asn	Leu	Thr	Ser	Ile	Asn	Asn	Ala	Gln	Lys
5255						5260					5265			
Asp	Ala	Leu	Thr	Arg	Ser	Ile	Asp	Gly	Ala	Thr	Thr	Val	Ala	Gly
5270						5275					5280			
Val	Asn	Gln	Glu	Thr	Ala	Lys	Ala	Thr	Glu	Leu	Asn	Asn	Ala	Met
5285						5290					5295			
His	Ser	Leu	Gln	Asn	Gly	Ile	Asn	Asp	Glu	Thr	Gln	Thr	Lys	Gln
5300						5305					5310			
Thr	Gln	Lys	Tyr	Leu	Asp	Ala	Glu	Pro	Ser	Lys	Lys	Ser	Ala	Tyr
5315						5320					5325			
Asp	Gln	Ala	Val	Asn	Ala	Ala	Lys	Ala	Ile	Leu	Thr	Lys	Ala	Ser
5330						5335					5340			
Gly	Gln	Asn	Val	Asp	Lys	Ala	Ala	Val	Glu	Gln	Ala	Leu	Gln	Asn
5345						5350					5355			
Val	Asn	Ser	Thr	Lys	Thr	Ala	Leu	Asn	Gly	Asp	Ala	Lys	Leu	Asn
5360						5365					5370			
Glu	Ala	Lys	Ala	Ala	Ala	Lys	Gln	Thr	Leu	Gly	Thr	Leu	Thr	His
5375						5380					5385			
Ile	Asn	Asn	Ala	Gln	Arg	Thr	Ala	Leu	Asp	Asn	Glu	Ile	Thr	Gln
5390						5395					5400			
Ala	Thr	Asn	Val	Glu	Gly	Val	Asn	Thr	Val	Lys	Ala	Lys	Ala	Gln
5405						5410					5415			
Gln	Leu	Asp	Gly	Ala	Met	Gly	Gln	Leu	Glu	Thr	Ser	Ile	Arg	Asp

-continued

---

5420	5425	5430
Lys Asp Thr Thr Leu Gln Ser Gln Asn Tyr Gln Asp Ala Asp Asp		
5435	5440	5445
Ala Lys Arg Thr Ala Tyr Ser Gln Ala Val Asn Ala Ala Ala Thr		
5450	5455	5460
Ile Leu Asn Lys Thr Ala Gly Gly Asn Thr Pro Lys Ala Asp Val		
5465	5470	5475
Glu Arg Ala Met Gln Ala Val Thr Gln Ala Asn Thr Ala Leu Asn		
5480	5485	5490
Gly Ile Gln Asn Leu Asp Arg Ala Lys Gln Ala Ala Asn Thr Ala		
5495	5500	5505
Ile Thr Asn Ala Ser Asp Leu Asn Thr Lys Gln Lys Glu Ala Leu		
5510	5515	5520
Lys Ala Gln Val Thr Ser Ala Gly Arg Val Ser Ala Ala Asn Gly		
5525	5530	5535
Val Glu His Thr Ala Thr Glu Leu Asn Thr Ala Met Thr Ala Leu		
5540	5545	5550
Lys Arg Ala Ile Ala Asp Lys Ala Glu Thr Lys Ala Ser Gly Asn		
5555	5560	5565
Tyr Val Asn Ala Asp Ala Asn Lys Arg Gln Ala Tyr Asp Glu Lys		
5570	5575	5580
Val Thr Ala Ala Glu Asn Ile Val Ser Gly Thr Pro Thr Pro Thr		
5585	5590	5595
Leu Thr Pro Ala Asp Val Thr Asn Ala Ala Thr Gln Val Thr Asn		
5600	5605	5610
Ala Lys Thr Gln Leu Asn Gly Asn His Asn Leu Glu Val Ala Lys		
5615	5620	5625
Gln Asn Ala Asn Thr Ala Ile Asp Gly Leu Thr Ser Leu Asn Gly		
5630	5635	5640
Pro Gln Lys Ala Lys Leu Lys Glu Gln Val Gly Gln Ala Thr Thr		
5645	5650	5655
Leu Pro Asn Val Gln Thr Val Arg Asp Asn Ala Gln Thr Leu Asn		
5660	5665	5670
Thr Ala Met Lys Gly Leu Arg Asp Ser Ile Ala Asn Glu Ala Thr		
5675	5680	5685
Ile Lys Ala Gly Gln Asn Tyr Thr Asp Ala Ser Gln Asn Lys Gln		
5690	5695	5700
Thr Asp Tyr Asn Ser Ala Val Thr Ala Ala Lys Ala Ile Ile Gly		
5705	5710	5715
Gln Thr Thr Ser Pro Ser Met Asn Ala Gln Glu Ile Asn Gln Ala		
5720	5725	5730
Lys Asp Gln Val Thr Ala Lys Gln Gln Ala Leu Asn Gly Gln Glu		
5735	5740	5745
Asn Leu Arg Thr Ala Gln Thr Asn Ala Lys Gln His Leu Asn Gly		
5750	5755	5760
Leu Ser Asp Leu Thr Asp Ala Gln Lys Asp Ala Val Lys Arg Gln		
5765	5770	5775
Ile Glu Gly Ala Thr His Val Asn Glu Val Thr Gln Ala Gln Asn		
5780	5785	5790
Asn Ala Asp Ala Leu Asn Thr Ala Met Thr Asn Leu Lys Asn Gly		
5795	5800	5805
Ile Gln Asp Gln Asn Thr Ile Lys Gln Gly Val Asn Phe Thr Asp		
5810	5815	5820

-continued

---

Ala Asp	Glu Ala Lys Arg	Asn	Ala Tyr Thr	Asn	Ala Val Thr Gln
5825		5830			5835
Ala Glu	Gln Ile Leu Asn	Lys	Ala Gln Gly Pro	Asn	Thr Ser Lys
5840		5845			5850
Asp Gly	Val Glu Thr Ala	Leu	Glu Asn Val Gln	Arg	Ala Lys Asn
5855		5860			5865
Glu Leu	Asn Gly Asn Gln	Asn	Val Ala Asn Ala	Lys	Thr Thr Ala
5870		5875			5880
Lys Asn	Ala Leu Asn Asn	Leu	Thr Ser Ile Asn	Asn	Ala Gln Lys
5885		5890			5895
Glu Ala	Leu Lys Ser Gln	Ile	Glu Gly Ala Thr	Thr	Val Ala Gly
5900		5905			5910
Val Asn	Gln Val Ser Thr	Thr	Ala Ser Glu Leu	Asn	Thr Ala Met
5915		5920			5925
Ser Asn	Leu Gln Asn Gly	Ile	Asn Asp Glu Ala	Ala	Thr Lys Ala
5930		5935			5940
Ala Gln	Lys Tyr Thr Asp	Ala	Asp Arg Glu Lys	Gln	Thr Ala Tyr
5945		5950			5955
Asn Asp	Ala Val Thr Ala	Ala	Lys Thr Leu Leu	Asp	Lys Thr Ala
5960		5965			5970
Gly Ser	Asn Asp Asn Lys	Ala	Ala Val Glu Gln	Ala	Leu Gln Arg
5975		5980			5985
Val Asn	Thr Ala Lys Thr	Ala	Leu Asn Gly Asp	Glu	Arg Leu Asn
5990		5995			6000
Glu Ala	Lys Asn Thr Ala	Lys	Gln Gln Val Ala	Thr	Met Ser His
6005		6010			6015
Leu Thr	Asp Ala Gln Lys	Ala	Asn Leu Thr Ser	Gln	Ile Glu Ser
6020		6025			6030
Gly Thr	Thr Val Ala Gly	Val	Gln Gly Ile Gln	Ala	Asn Ala Gly
6035		6040			6045
Thr Leu	Asp Gln Ala Met	Asn	Gln Leu Arg Gln	Ser	Ile Ala Ser
6050		6055			6060
Lys Asp	Ala Thr Lys Ser	Ser	Glu Asp Tyr Gln	Asp	Ala Asn Ala
6065		6070			6075
Asp Leu	Gln Asn Ala Tyr	Asn	Asp Ala Val Thr	Asn	Ala Glu Gly
6080		6085			6090
Ile Ile	Ser Ala Thr Asn	Asn	Pro Glu Met Asn	Pro	Asp Thr Ile
6095		6100			6105
Asn Gln	Lys Ala Ser Gln	Val	Asn Ser Ala Lys	Ser	Ala Leu Asn
6110		6115			6120
Gly Asp	Glu Lys Leu Ala	Ala	Ala Lys Gln Thr	Ala	Lys Ser Asp
6125		6130			6135
Ile Gly	Arg Leu Thr Asp	Leu	Asn Asn Ala Gln	Arg	Thr Ala Ala
6140		6145			6150
Asn Ala	Glu Val Asp Gln	Ala	Pro Asn Leu Ala	Ala	Val Thr Ala
6155		6160			6165
Ala Lys	Asn Lys Ala Thr	Ser	Leu Asn Thr Ala	Met	Gly Asn Leu
6170		6175			6180
Lys His	Ala Leu Ala Glu	Lys	Asp Asn Thr Lys	Arg	Ser Val Asn
6185		6190			6195
Tyr Thr	Asp Ala Asp Gln	Pro	Lys Gln Gln Ala	Tyr	Asp Thr Ala
6200		6205			6210

-continued

---

Val Thr	Gln Ala	Glu Ala	Ile	Thr Asn	Ala Asn	Gly	Ser Asn	Ala		
6215			6220			6225				
Asn Glu	Thr Gln	Val Gln	Ala	Ala Leu	Asn Gln	Leu	Asn Gln	Ala		
6230			6235			6240				
Lys Asn	Asp Leu	Asn Gly	Asp	Asn Lys	Val Ala	Gln	Ala Lys	Glu		
6245			6250			6255				
Ser Ala	Lys Arg	Ala Leu	Ala	Ser Tyr	Ser Asn	Leu	Asn Asn	Ala		
6260			6265			6270				
Gln Ser	Thr Ala	Ala Ile	Ser	Gln Ile	Asp Asn	Ala	Thr Thr	Val		
6275			6280			6285				
Ala Gly	Val Thr	Ala Ala	Gln	Asn Thr	Ala Asn	Glu	Leu Asn	Thr		
6290			6295			6300				
Ala Met	Gly Gln	Leu Gln	Asn	Gly Ile	Asn Asp	Gln	Asn Thr	Val		
6305			6310			6315				
Lys Gln	Gln Val	Asn Phe	Thr	Asp Ala	Asp Gln	Gly	Lys Lys	Asp		
6320			6325			6330				
Ala Tyr	Thr Asn	Ala Val	Thr	Asn Ala	Gln Gly	Ile	Leu Asp	Lys		
6335			6340			6345				
Ala His	Gly Gln	Asn Met	Thr	Lys Ala	Gln Val	Glu	Ala Ala	Leu		
6350			6355			6360				
Asn Gln	Val Thr	Thr Ala	Lys	Asn Ala	Leu Asn	Gly	Asp Ala	Asn		
6365			6370			6375				
Val Arg	Gln Ala	Lys Ser	Asp	Ala Lys	Ala Asn	Leu	Gly Thr	Leu		
6380			6385			6390				
Thr His	Leu Asn	Asn Ala	Gln	Lys Gln	Asp Leu	Thr	Ser Gln	Ile		
6395			6400			6405				
Glu Gly	Ala Thr	Thr Val	Asn	Gly Val	Asn Gly	Val	Lys Thr	Lys		
6410			6415			6420				
Ala Gln	Asp Leu	Asp Gly	Ala	Met Gln	Arg Leu	Gln	Ser Ala	Ile		
6425			6430			6435				
Ala Asn	Lys Asp	Gln Thr	Lys	Ala Ser	Glu Asn	Tyr	Ile Asp	Ala		
6440			6445			6450				
Asp Pro	Thr Lys	Lys Thr	Ala	Phe Asp	Asn Ala	Ile	Thr Gln	Ala		
6455			6460			6465				
Glu Ser	Tyr Leu	Asn Lys	Asp	His Gly	Ala Asn	Lys	Asp Lys	Gln		
6470			6475			6480				
Ala Val	Glu Gln	Ala Ile	Gln	Ser Val	Thr Ser	Thr	Glu Asn	Ala		
6485			6490			6495				
Leu Asn	Gly Asp	Ala Asn	Leu	Gln Arg	Ala Lys	Thr	Glu Ala	Ile		
6500			6505			6510				
Gln Ala	Ile Asp	Asn Leu	Thr	His Leu	Asn Thr	Pro	Gln Lys	Thr		
6515			6520			6525				
Ala Leu	Lys Gln	Gln Val	Asn	Ala Ala	Gln Arg	Val	Ser Gly	Val		
6530			6535			6540				
Thr Asp	Leu Lys	Asn Ser	Ala	Thr Ser	Leu Asn	Asn	Ala Met	Asp		
6545			6550			6555				
Gln Leu	Lys Gln	Ala Ile	Ala	Asp His	Asp Thr	Ile	Val Ala	Ser		
6560			6565			6570				
Gly Asn	Tyr Thr	Asn Ala	Ser	Pro Asp	Lys Gln	Gly	Ala Tyr	Thr		
6575			6580			6585				
Asp Ala	Tyr Asn	Ala Ala	Lys	Asn Ile	Val Asn	Gly	Ser Pro	Asn		
6590			6595			6600				
Val Ile	Thr Asn	Ala Ala	Asp	Val Thr	Ala Ala	Thr	Gln Arg	Val		

-continued

---

6605	6610	6615
Asn Asn Ala Glu Thr Gly Leu Asn Gly Asp Thr Asn Leu Ala Thr 6620 6625 6630		
Ala Lys Gln Gln Ala Lys Asp Ala Leu Arg Gln Met Thr His Leu 6635 6640 6645		
Ser Asp Ala Gln Lys Gln Ser Ile Thr Gly Gln Ile Asp Ser Ala 6650 6655 6660		
Thr Gln Val Thr Gly Val Gln Ser Val Lys Asp Asn Ala Thr Asn 6665 6670 6675		
Leu Asp Asn Ala Met Asn Gln Leu Arg Asn Ser Ile Ala Asn Lys 6680 6685 6690		
Asp Asp Val Lys Ala Ser Gln Pro Tyr Val Asp Ala Asp Arg Asp 6695 6700 6705		
Lys Gln Asn Ala Tyr Asn Thr Ala Val Thr Asn Ala Glu Asn Ile 6710 6715 6720		
Ile Asn Ala Thr Ser Gln Pro Thr Leu Asp Pro Ser Ala Val Thr 6725 6730 6735		
Gln Ala Ala Asn Gln Val Ser Thr Asn Lys Thr Ala Leu Asn Gly 6740 6745 6750		
Ala Gln Asn Leu Ala Asn Lys Lys Gln Glu Thr Thr Ala Asn Ile 6755 6760 6765		
Asn Gln Leu Ser His Leu Asn Asn Ala Gln Lys Gln Asp Leu Asn 6770 6775 6780		
Thr Gln Val Thr Asn Ala Pro Asn Ile Ser Thr Val Asn Gln Val 6785 6790 6795		
Lys Thr Lys Ala Glu Gln Leu Asp Gln Ala Met Glu Arg Leu Ile 6800 6805 6810		
Asn Gly Ile Gln Asp Lys Asp Gln Val Lys Gln Ser Val Asn Phe 6815 6820 6825		
Thr Asp Ala Asp Pro Glu Lys Gln Thr Ala Tyr Asn Asn Ala Val 6830 6835 6840		
Thr Ala Ala Glu Asn Ile Ile Asn Gln Ala Asn Gly Thr Asn Ala 6845 6850 6855		
Asn Gln Ser Gln Val Glu Ala Ala Leu Ser Thr Val Thr Thr Thr 6860 6865 6870		
Lys Gln Ala Leu Asn Gly Asp Arg Lys Val Thr Asp Ala Lys Asn 6875 6880 6885		
Asn Ala Asn Gln Thr Leu Ser Thr Leu Asp Asn Leu Asn Asn Ala 6890 6895 6900		
Gln Lys Gly Ala Val Thr Gly Asn Ile Asn Gln Ala His Thr Val 6905 6910 6915		
Ala Glu Val Thr Gln Ala Ile Gln Thr Ala Gln Glu Leu Asn Thr 6920 6925 6930		
Ala Met Gly Asn Leu Lys Asn Ser Leu Asn Asp Lys Asp Thr Thr 6935 6940 6945		
Leu Gly Ser Gln Asn Phe Ala Asp Ala Asp Pro Glu Lys Lys Asn 6950 6955 6960		
Ala Tyr Asn Glu Ala Val His Asn Ala Glu Asn Ile Leu Asn Lys 6965 6970 6975		
Ser Thr Gly Thr Asn Val Pro Lys Asp Gln Val Glu Ala Ala Met 6980 6985 6990		
Asn Gln Val Asn Ala Thr Lys Ala Ala Leu Asn Gly Thr Gln Asn 6995 7000 7005		

-continued

---

Leu	Glu	Lys	Ala	Lys	Gln	His	Ala	Asn	Thr	Ala	Ile	Asp	Gly	Leu
7010						7015						7020		
Ser	His	Leu	Thr	Asn	Ala	Gln	Lys	Glu	Ala	Leu	Lys	Gln	Leu	Val
7025						7030						7035		
Gln	Gln	Ser	Thr	Thr	Val	Ala	Glu	Ala	Gln	Gly	Asn	Glu	Gln	Lys
7040						7045						7050		
Ala	Asn	Asn	Val	Asp	Ala	Ala	Met	Asp	Lys	Leu	Arg	Gln	Ser	Ile
7055						7060						7065		
Ala	Asp	Asn	Ala	Thr	Thr	Lys	Gln	Asn	Gln	Asn	Tyr	Thr	Asp	Ala
7070						7075						7080		
Ser	Gln	Asn	Lys	Lys	Asp	Ala	Tyr	Asn	Asn	Ala	Val	Thr	Thr	Ala
7085						7090						7095		
Gln	Gly	Ile	Ile	Asp	Gln	Thr	Thr	Ser	Pro	Thr	Leu	Asp	Pro	Thr
7100						7105						7110		
Val	Ile	Asn	Gln	Ala	Ala	Gly	Gln	Val	Ser	Thr	Thr	Lys	Asn	Ala
7115						7120						7125		
Leu	Asn	Gly	Asn	Glu	Asn	Leu	Glu	Ala	Ala	Lys	Gln	Gln	Ala	Ser
7130						7135						7140		
Gln	Ser	Leu	Gly	Ser	Leu	Asp	Asn	Leu	Asn	Asn	Ala	Gln	Lys	Gln
7145						7150						7155		
Thr	Val	Thr	Asp	Gln	Ile	Asn	Gly	Ala	His	Thr	Val	Asp	Glu	Ala
7160						7165						7170		
Asn	Gln	Ile	Lys	Gln	Asn	Ala	Gln	Asn	Leu	Asn	Thr	Ala	Met	Gly
7175						7180						7185		
Asn	Leu	Lys	Gln	Ala	Ile	Ala	Asp	Lys	Asp	Ala	Thr	Lys	Ala	Thr
7190						7195						7200		
Val	Asn	Phe	Thr	Asp	Ala	Asp	Gln	Ala	Lys	Gln	Gln	Ala	Tyr	Asn
7205						7210						7215		
Thr	Ala	Val	Thr	Asn	Ala	Glu	Asn	Ile	Ser	Lys	Ala	Asn	Gly	Asn
7220						7225						7230		
Ala	Thr	Gln	Ala	Glu	Val	Glu	Gln	Ala	Ile	Lys	Gln	Val	Asn	Ala
7235						7240						7245		
Ala	Lys	Gln	Ala	Leu	Asn	Gly	Asn	Ala	Asn	Val	Gln	His	Ala	Lys
7250						7255						7260		
Asp	Glu	Ala	Thr	Ala	Leu	Ile	Asn	Ser	Ser	Asn	Asp	Leu	Asn	Gln
7265						7270						7275		
Ala	Gln	Lys	Asp	Ala	Leu	Lys	Gln	Gln	Val	Gln	Asn	Ala	Thr	Thr
7280						7285						7290		
Val	Ala	Gly	Val	Asn	Asn	Val	Lys	Gln	Thr	Ala	Gln	Glu	Leu	Asn
7295						7300						7305		
Asn	Ala	Met	Thr	Gln	Leu	Lys	Gln	Gly	Ile	Ala	Asp	Lys	Glu	Gln
7310						7315						7320		
Thr	Lys	Ala	Asp	Gly	Asn	Phe	Val	Asn	Ala	Asp	Pro	Asp	Lys	Gln
7325						7330						7335		
Asn	Ala	Tyr	Asn	Gln	Ala	Val	Ala	Lys	Ala	Glu	Ala	Leu	Ile	Ser
7340						7345						7350		
Ala	Thr	Pro	Asp	Val	Val	Val	Thr	Pro	Ser	Glu	Ile	Thr	Ala	Ala
7355						7360						7365		
Leu	Asn	Lys	Val	Thr	Gln	Ala	Lys	Asn	Asp	Leu	Asn	Gly	Asn	Thr
7370						7375						7380		
Asn	Leu	Ala	Thr	Ala	Lys	Gln	Asn	Val	Gln	His	Ala	Ile	Asp	Gln
7385						7390						7395		

-continued

Leu Pro 7400	Asn Leu Asn Gln Ala 7405	Gln Arg Asp Glu Tyr 7410	Ser Lys Gln
Ile Thr 7415	Gln Ala Thr Leu Val 7420	Pro Asn Val Asn Ala 7425	Ile Gln Gln
Ala Ala 7430	Thr Thr Leu Asn Asp 7435	Ala Met Thr Gln Leu 7440	Lys Gln Gly
Ile Ala 7445	Asn Lys Ala Gln Ile 7450	Lys Gly Ser Glu Asn 7455	Tyr His Asp
Ala Asp 7460	Thr Asp Lys Gln Thr 7465	Ala Tyr Asp Asn Ala 7470	Val Thr Lys
Ala Glu 7475	Glu Leu Leu Lys Gln 7480	Thr Thr Asn Pro Thr 7485	Met Asp Pro
Asn Thr 7490	Ile Gln Gln Ala Leu 7495	Thr Lys Val Asn Asp 7500	Thr Asn Gln
Ala Leu 7505	Asn Gly Asn Gln Lys 7510	Leu Ala Asp Ala Lys 7515	Gln Asp Ala
Lys Thr 7520	Thr Leu Gly Thr Leu 7525	Asp His Leu Asn Asp 7530	Ala Gln Lys
Gln Ala 7535	Leu Thr Thr Gln Val 7540	Glu Gln Ala Pro Asp 7545	Ile Ala Thr
Val Asn 7550	Asn Val Lys Gln Asn 7555	Ala Gln Asn Leu Asn 7560	Asn Ala Met
Thr Asn 7565	Leu Asn Asn Ala Leu 7570	Gln Asp Lys Thr Glu 7575	Thr Leu Asn
Ser Ile 7580	Asn Phe Thr Asp Ala 7585	Asp Gln Ala Lys Lys 7590	Asp Ala Tyr
Thr Asn 7595	Ala Val Ser His Ala 7600	Glu Gly Ile Leu Ser 7605	Lys Ala Asn
Gly Ser 7610	Asn Ala Ser Gln Thr 7615	Glu Val Glu Gln Ala 7620	Met Gln Arg
Val Asn 7625	Glu Ala Lys Gln Ala 7630	Leu Asn Gly Asn Asp 7635	Asn Val Gln
Arg Ala 7640	Lys Asp Ala Ala Lys 7645	Gln Val Ile Thr Asn 7650	Ala Asn Asp
Leu Asn 7655	Gln Ala Gln Lys Asp 7660	Ala Leu Lys Gln Gln 7665	Val Asp Ala
Ala Gln 7670	Thr Val Ala Asn Val 7675	Asn Thr Ile Lys Gln 7680	Thr Ala Gln
Asp Leu 7685	Asn Gln Ala Met Thr 7690	Gln Leu Lys Gln Gly 7695	Ile Ala Asp
Lys Asp 7700	Gln Thr Lys Ala Asn 7705	Gly Asn Phe Val Asn 7710	Ala Asp Thr
Asp Lys 7715	Gln Asn Ala Tyr Asn 7720	Asn Ala Val Ala His 7725	Ala Glu Gln
Ile Ile 7730	Ser Gly Thr Pro Asn 7735	Ala Asn Val Asp Pro 7740	Gln Gln Val
Ala Gln 7745	Ala Leu Gln Gln Val 7750	Asn Gln Ala Lys Gly 7755	Asp Leu Asn
Gly Asn 7760	His Asn Leu Gln Val 7765	Ala Lys Asp Asn Ala 7770	Asn Thr Ala
Ile Asp 7775	Gln Leu Pro Asn Leu 7780	Asn Gln Pro Gln Lys 7785	Thr Ala Leu
Lys Asp 7790	Gln Val Ser His Ala 7795	Glu Leu Val Thr Gly 7800	Val Asn Ala

-continued

---

7790	7795	7800
Ile Lys Gln Asn Ala Asp 7805	Ala Leu Asn Asn Ala 7810	Met Gly Thr Leu 7815
Lys Gln Gln Ile Gln Ala 7820	Asn Ser Gln Val Pro 7825	Gln Ser Val Asp 7830
Phe Thr Gln Ala Asp Gln 7835	Asp Lys Gln Gln Ala 7840	Tyr Asn Asn Ala 7845
Ala Asn Gln Ala Gln Gln 7850	Ile Ala Asn Gly Ile 7855	Pro Thr Pro Val 7860
Leu Thr Pro Asp Thr Val 7865	Thr Gln Ala Val Thr 7870	Thr Met Asn Gln 7875
Ala Lys Asp Ala Leu Asn 7880	Gly Asp Glu Lys Leu 7885	Ala Gln Ala Lys 7890
Gln Glu Ala Leu Ala Asn 7895	Leu Asp Thr Leu Arg 7900	Asp Leu Asn Gln 7905
Pro Gln Arg Asp Ala Leu 7910	Arg Asn Gln Ile Asn 7915	Gln Ala Gln Ala 7920
Leu Ala Thr Val Glu Gln 7925	Thr Lys Gln Asn Ala 7930	Gln Asn Val Asn 7935
Thr Ala Met Ser Asn Leu 7940	Lys Gln Gly Ile Ala 7945	Asn Lys Asp Thr 7950
Val Lys Ala Ser Glu Asn 7955	Tyr His Asp Ala Asp 7960	Ala Asp Lys Gln 7965
Thr Ala Tyr Thr Asn Ala 7970	Val Ser Gln Ala Glu 7975	Gly Ile Ile Asn 7980
Gln Thr Thr Asn Pro Thr 7985	Leu Asn Pro Asp Glu 7990	Ile Thr Arg Ala 7995
Leu Thr Gln Val Thr Asp 8000	Ala Lys Asn Gly Leu 8005	Asn Gly Glu Ala 8010
Lys Leu Ala Thr Glu Lys 8015	Gln Asn Ala Lys Asp 8020	Ala Val Ser Gly 8025
Met Thr His Leu Asn Asp 8030	Ala Gln Lys Gln Ala 8035	Leu Lys Gly Gln 8040
Ile Asp Gln Ser Pro Glu 8045	Ile Ala Thr Val Asn 8050	Gln Val Lys Gln 8055
Thr Ala Thr Ser Leu Asp 8060	Gln Ala Met Asp Gln 8065	Leu Ser Gln Ala 8070
Ile Asn Asp Lys Ala Gln 8075	Thr Leu Ala Asp Gly 8080	Asn Tyr Leu Asn 8085
Ala Asp Pro Asp Lys Gln 8090	Asn Ala Tyr Lys Gln 8095	Ala Val Ala Lys 8100
Ala Glu Ala Leu Leu Asn 8105	Lys Gln Ser Gly Thr 8110	Asn Glu Val Gln 8115
Ala Gln Val Glu Ser Ile 8120	Thr Asn Glu Val Asn 8125	Ala Ala Lys Gln 8130
Ala Leu Asn Gly Asn Asp 8135	Asn Leu Ala Asn Ala 8140	Lys Gln Gln Ala 8145
Lys Gln Gln Leu Ala Asn 8150	Leu Thr His Leu Asn 8155	Asp Ala Gln Lys 8160
Gln Ser Phe Glu Ser Gln 8165	Ile Thr Gln Ala Pro 8170	Leu Val Thr Asp 8175
Val Thr Thr Ile Asn Gln 8180	Lys Ala Gln Thr Leu 8185	Asp His Ala Met 8190

-continued

---

Glu Leu	Leu Arg	Asn Ser	Val	Ala Asp	Asn Gln	Thr	Thr Leu	Ala	
8195			8200			8205			
Ser Glu	Asp Tyr	His Asp	Ala	Thr Ala	Gln Arg	Gln	Asn Asp	Tyr	
8210			8215			8220			
Asn Gln	Ala Val	Thr Ala	Ala	Asn Asn	Ile Ile	Asn	Gln Thr	Thr	
8225			8230			8235			
Ser Pro	Thr Met	Asn Pro	Asp	Asp Val	Asn Gly	Ala	Thr Thr	Gln	
8240			8245			8250			
Val Asn	Asn Thr	Lys Val	Ala	Leu Asp	Gly Asp	Glu	Asn Leu	Ala	
8255			8260			8265			
Ala Ala	Lys Gln	Gln Ala	Asn	Asn Arg	Leu Asp	Gln	Leu Asp	His	
8270			8275			8280			
Leu Asn	Asn Ala	Gln Lys	Gln	Gln Leu	Gln Ser	Gln	Ile Thr	Gln	
8285			8290			8295			
Ser Ser	Asp Ile	Ala Ala	Val	Asn Gly	His Lys	Gln	Thr Ala	Glu	
8300			8305			8310			
Ser Leu	Asn Thr	Ala Met	Gly	Asn Leu	Ile Asn	Ala	Ile Ala	Asp	
8315			8320			8325			
His Gln	Ala Val	Glu Gln	Arg	Gly Asn	Phe Ile	Asn	Ala Asp	Thr	
8330			8335			8340			
Asp Lys	Gln Thr	Ala Tyr	Asn	Thr Ala	Val Asn	Glu	Ala Ala	Ala	
8345			8350			8355			
Met Ile	Asn Lys	Gln Thr	Gly	Gln Asn	Ala Asn	Gln	Thr Glu	Val	
8360			8365			8370			
Glu Gln	Ala Ile	Thr Lys	Val	Gln Thr	Thr Leu	Gln	Ala Leu	Asn	
8375			8380			8385			
Gly Asp	His Asn	Leu Gln	Val	Ala Lys	Thr Asn	Ala	Thr Gln	Ala	
8390			8395			8400			
Ile Asp	Ala Leu	Thr Ser	Leu	Asn Asp	Pro Gln	Lys	Thr Ala	Leu	
8405			8410			8415			
Lys Asp	Gln Val	Thr Ala	Ala	Thr Leu	Val Thr	Ala	Val His	Gln	
8420			8425			8430			
Ile Glu	Gln Asn	Ala Asn	Thr	Leu Asn	Gln Ala	Met	His Gly	Leu	
8435			8440			8445			
Arg Gln	Ser Ile	Gln Asp	Asn	Ala Ala	Thr Lys	Ala	Asn Ser	Lys	
8450			8455			8460			
Tyr Ile	Asn Glu	Asp Gln	Pro	Glu Gln	Gln Asn	Tyr	Asp Gln	Ala	
8465			8470			8475			
Val Gln	Ala Ala	Asn Asn	Ile	Ile Asn	Glu Gln	Thr	Ala Thr	Leu	
8480			8485			8490			
Asp Asn	Asn Ala	Ile Asn	Gln	Ala Ala	Thr Thr	Val	Asn Thr	Thr	
8495			8500			8505			
Lys Ala	Ala Leu	His Gly	Asp	Val Lys	Leu Gln	Asn	Asp Lys	Asp	
8510			8515			8520			
His Ala	Lys Gln	Thr Val	Ser	Gln Leu	Ala His	Leu	Asn Asn	Ala	
8525			8530			8535			
Gln Lys	His Met	Glu Asp	Thr	Leu Ile	Asp Ser	Glu	Thr Thr	Arg	
8540			8545			8550			
Thr Ala	Val Lys	Gln Asp	Leu	Thr Glu	Ala Gln	Ala	Leu Asp	Gln	
8555			8560			8565			
Leu Met	Asp Ala	Leu Gln	Gln	Ser Ile	Ala Asp	Lys	Asp Ala	Thr	
8570			8575			8580			

-continued

---

Arg Ala 8585	Ser Ser Ala Tyr Val 8590	Asn Ala Glu Pro Asn 8595	Lys Lys Gln
Ser Tyr 8600	Asp Glu Ala Val Gln 8605	Asn Ala Glu Ser Ile 8610	Ile Ala Gly
Leu Asn 8615	Asn Pro Thr Ile Asn 8620	Lys Gly Asn Val Ser 8625	Ser Ala Thr
Gln Ala 8630	Val Ile Ser Ser Lys 8635	Asn Ala Leu Asp Gly 8640	Val Glu Arg
Leu Ala 8645	Gln Asp Lys Gln Thr 8650	Ala Gly Asn Ser Leu 8655	Asn His Leu
Asp Gln 8660	Leu Thr Pro Ala Gln 8665	Gln Gln Ala Leu Glu 8670	Asn Gln Ile
Asn Asn 8675	Ala Thr Thr Arg Gly 8680	Glu Val Ala Gln Lys 8685	Leu Thr Glu
Ala Gln 8690	Ala Leu Asn Gln Ala 8695	Met Glu Ala Leu Arg 8700	Asn Ser Ile
Gln Asp 8705	Gln Gln Gln Thr Glu 8710	Ala Gly Ser Lys Phe 8715	Ile Asn Glu
Asp Lys 8720	Pro Gln Lys Asp Ala 8725	Tyr Gln Ala Ala Val 8730	Gln Asn Ala
Lys Asp 8735	Leu Ile Asn Gln Thr 8740	Asn Asn Pro Thr Leu 8745	Asp Lys Ala
Gln Val 8750	Glu Gln Leu Thr Gln 8755	Ala Val Asn Gln Ala 8760	Lys Asp Asn
Leu His 8765	Gly Asp Gln Lys Leu 8770	Ala Asp Asp Lys Gln 8775	His Ala Val
Thr Asp 8780	Leu Asn Gln Leu Asn 8785	Gly Leu Asn Asn Pro 8790	Gln Arg Gln
Ala Leu 8795	Glu Ser Gln Ile Asn 8800	Asn Ala Ala Thr Arg 8805	Gly Glu Val
Ala Gln 8810	Lys Leu Ala Glu Ala 8815	Lys Ala Leu Asp Gln 8820	Ala Met Gln
Ala Leu 8825	Arg Asn Ser Ile Gln 8830	Asp Gln Gln Gln Thr 8835	Glu Ser Gly
Ser Lys 8840	Phe Ile Asn Glu Asp 8845	Lys Pro Gln Lys Asp 8850	Ala Tyr Gln
Ala Ala 8855	Val Gln Asn Ala Lys 8860	Asp Leu Ile Asn Gln 8865	Thr Gly Asn
Pro Thr 8870	Leu Asp Lys Ser Gln 8875	Val Glu Gln Leu Thr 8880	Gln Ala Val
Thr Thr 8885	Ala Lys Asp Asn Leu 8890	His Gly Asp Gln Lys 8895	Leu Ala Arg
Asp Gln 8900	Gln Gln Ala Val Thr 8905	Thr Val Asn Ala Leu 8910	Pro Asn Leu
Asn His 8915	Ala Gln Gln Gln Ala 8920	Leu Thr Asp Ala Ile 8925	Asn Ala Ala
Pro Thr 8930	Arg Thr Glu Val Ala 8935	Gln His Val Gln Thr 8940	Ala Thr Glu
Leu Asp 8945	His Ala Met Glu Thr 8950	Leu Lys Asn Lys Val 8955	Asp Gln Val
Asn Thr 8960	Asp Lys Ala Gln Pro 8965	Asn Tyr Thr Glu Ala 8970	Ser Thr Asp
Lys Lys	Glu Ala Val Asp Gln	Ala Leu Gln Ala Ala	Glu Ser Ile

-continued

8975	8980	8985
Thr Asp Pro Thr Asn Gly Ser Asn Ala Asn Lys Asp Ala Val Asp 8990 8995 9000		
Gln Val Leu Thr Lys Leu Gln Glu Lys Glu Asn Glu Leu Asn Gly 9005 9010 9015		
Asn Glu Arg Val Ala Glu Ala Lys Thr Gln Ala Lys Gln Thr Ile 9020 9025 9030		
Asp Gln Leu Thr His Leu Asn Ala Asp Gln Ile Ala Thr Ala Lys 9035 9040 9045		
Gln Asn Ile Asp Gln Ala Thr Lys Leu Gln Pro Ile Ala Glu Leu 9050 9055 9060		
Val Asp Gln Ala Thr Gln Leu Asn Gln Ser Met Asp Gln Leu Gln 9065 9070 9075		
Gln Ala Val Asn Glu His Ala Asn Val Glu Gln Thr Val Asp Tyr 9080 9085 9090		
Thr Gln Ala Asp Ser Asp Lys Gln Asn Ala Tyr Lys Gln Ala Ile 9095 9100 9105		
Ala Asp Ala Glu Asn Val Leu Lys Gln Asn Ala Asn Lys Gln Gln 9110 9115 9120		
Val Asp Gln Ala Leu Gln Asn Ile Leu Asn Ala Lys Gln Ala Leu 9125 9130 9135		
Asn Gly Asp Glu Arg Val Ala Leu Ala Lys Thr Asn Gly Lys His 9140 9145 9150		
Asp Ile Asp Gln Leu Asn Ala Leu Asn Asn Ala Gln Gln Asp Gly 9155 9160 9165		
Phe Lys Gly Arg Ile Asp Gln Ser Asn Asp Leu Asn Gln Ile Gln 9170 9175 9180		
Gln Ile Val Asp Glu Ala Lys Ala Leu Asn Arg Ala Met Asp Gln 9185 9190 9195		
Leu Ser Gln Glu Ile Thr Asp Asn Glu Gly Arg Thr Lys Gly Ser 9200 9205 9210		
Thr Asn Tyr Val Asn Ala Asp Thr Gln Val Lys Gln Val Tyr Asp 9215 9220 9225		
Glu Thr Val Asp Lys Ala Lys Gln Ala Leu Asp Lys Ser Thr Gly 9230 9235 9240		
Gln Asn Leu Thr Ala Lys Gln Val Ile Lys Leu Asn Asp Ala Val 9245 9250 9255		
Thr Ala Ala Lys Lys Ala Leu Asn Gly Glu Glu Arg Leu Asn Asn 9260 9265 9270		
Arg Lys Ala Glu Ala Leu Gln Arg Leu Asp Gln Leu Thr His Leu 9275 9280 9285		
Asn Asn Ala Gln Arg Gln Leu Ala Ile Gln Gln Ile Asn Asn Ala 9290 9295 9300		
Glu Thr Leu Asn Lys Ala Ser Arg Ala Ile Asn Arg Ala Thr Lys 9305 9310 9315		
Leu Asp Asn Ala Met Gly Ala Val Gln Gln Tyr Ile Asp Glu Gln 9320 9325 9330		
His Leu Gly Val Ile Ser Ser Thr Asn Tyr Ile Asn Ala Asp Asp 9335 9340 9345		
Asn Leu Lys Ala Asn Tyr Asp Asn Ala Ile Ala Asn Ala Ala His 9350 9355 9360		
Glu Leu Asp Lys Val Gln Gly Asn Ala Ile Ala Lys Ala Glu Ala 9365 9370 9375		

-continued

---

Glu	Gln	Leu	Lys	Gln	Asn	Ile	Ile	Asp	Ala	Gln	Asn	Ala	Leu	Asn
9380						9385					9390			
Gly	Asp	Gln	Asn	Leu	Ala	Asn	Ala	Lys	Asp	Lys	Ala	Asn	Ala	Phe
9395						9400					9405			
Val	Asn	Ser	Leu	Asn	Gly	Leu	Asn	Gln	Gln	Gln	Gln	Asp	Leu	Ala
9410						9415					9420			
His	Lys	Ala	Ile	Asn	Asn	Ala	Asp	Thr	Val	Ser	Asp	Val	Thr	Asp
9425						9430					9435			
Ile	Val	Asn	Asn	Gln	Ile	Asp	Leu	Asn	Asp	Ala	Met	Glu	Thr	Leu
9440						9445					9450			
Lys	His	Leu	Val	Asp	Asn	Glu	Ile	Pro	Asn	Ala	Glu	Gln	Thr	Val
9455						9460					9465			
Asn	Tyr	Gln	Asn	Ala	Asp	Asp	Asn	Ala	Lys	Thr	Asn	Phe	Asp	Asp
9470						9475					9480			
Ala	Lys	Arg	Leu	Ala	Asn	Thr	Leu	Leu	Asn	Ser	Asp	Asn	Thr	Asn
9485						9490					9495			
Val	Asn	Asp	Ile	Asn	Gly	Ala	Ile	Gln	Ala	Val	Asn	Asp	Ala	Ile
9500						9505					9510			
His	Asn	Leu	Asn	Gly	Asp	Gln	Arg	Leu	Gln	Asp	Ala	Lys	Asp	Lys
9515						9520					9525			
Ala	Ile	Gln	Ser	Ile	Asn	Gln	Ala	Leu	Ala	Asn	Lys	Leu	Lys	Glu
9530						9535					9540			
Ile	Glu	Ala	Ser	Asn	Ala	Thr	Asp	Gln	Asp	Lys	Leu	Ile	Ala	Lys
9545						9550					9555			
Asn	Lys	Ala	Glu	Glu	Leu	Ala	Asn	Ser	Ile	Ile	Asn	Asn	Ile	Asn
9560						9565					9570			
Lys	Ala	Thr	Ser	Asn	Gln	Ala	Val	Ser	Gln	Val	Gln	Thr	Ala	Gly
9575						9580					9585			
Asn	His	Ala	Ile	Glu	Gln	Val	His	Ala	Asn	Glu	Ile	Pro	Lys	Ala
9590						9595					9600			
Lys	Ile	Asp	Ala	Asn	Lys	Asp	Val	Asp	Lys	Gln	Val	Gln	Ala	Leu
9605						9610					9615			
Ile	Asp	Glu	Ile	Asp	Arg	Asn	Pro	Asn	Leu	Thr	Asp	Lys	Glu	Lys
9620						9625					9630			
Gln	Ala	Leu	Lys	Asp	Arg	Ile	Asn	Gln	Ile	Leu	Gln	Gln	Gly	His
9635						9640					9645			
Asn	Gly	Ile	Asn	Asn	Ala	Met	Thr	Lys	Glu	Glu	Ile	Glu	Gln	Ala
9650						9655					9660			
Lys	Ala	Gln	Leu	Ala	Gln	Ala	Leu	Gln	Asp	Ile	Lys	Asp	Leu	Val
9665						9670					9675			
Lys	Ala	Lys	Glu	Asp	Ala	Lys	Gln	Asp	Val	Asp	Lys	Gln	Val	Gln
9680						9685					9690			
Ala	Leu	Ile	Asp	Glu	Ile	Asp	Gln	Asn	Pro	Asn	Leu	Thr	Asp	Lys
9695						9700					9705			
Glu	Lys	Gln	Ala	Leu	Lys	Tyr	Arg	Ile	Asn	Gln	Ile	Leu	Gln	Gln
9710						9715					9720			
Gly	His	Asn	Asp	Ile	Asn	Asn	Ala	Leu	Thr	Lys	Glu	Glu	Ile	Glu
9725						9730					9735			
Gln	Ala	Lys	Ala	Gln	Leu	Ala	Gln	Ala	Leu	Gln	Asp	Ile	Lys	Asp
9740						9745					9750			
Leu	Val	Lys	Ala	Lys	Glu	Asp	Ala	Lys	Asn	Ala	Ile	Lys	Ala	Leu
9755						9760					9765			



-continued

10160	10165	10170
Gly Asn 10175	Glu Ser Asn Ser His 10180	Leu Thr Ile Gly Tyr 10185
Asn His 10190	Pro Phe Asn Ser Ser 10195	Thr Ile Gly His Lys 10200
Asp Glu 10205	Asp Asp Asp Ile Asp 10210	Pro Leu His Met Arg 10215
Asn Asn 10220	Phe Gly Asn Val Ile 10225	Lys Asn Ala Ile Gly 10230
Ile Ser 10235	Gly Leu Leu Ala Ser 10240	Phe Trp Phe Phe Ile 10245
Arg Arg 10250	Lys Glu Asp Glu Glu 10255	Glu Glu Leu Glu Ile 10260
Asn Lys 10265	Asp Ser Ile Lys Glu 10270	Thr Leu Asp Asp Thr 10275
Pro Leu 10280	Leu Phe Ala Lys Arg 10285	Arg Arg Lys Glu Asp 10290
Val Thr 10295	Val Glu Glu Lys Asp 10300	Ser Leu Asn Asn Gly 10305
Asp Lys 10310	Val Lys His Thr Pro 10315	Phe Phe Leu Pro Lys 10320
Lys Glu 10325	Asp Glu Glu Asp Val 10330	Glu Val Thr Asn Glu 10335
Glu Lys 10340	Val Leu Lys Asp Asn 10345	Glu His Ser Pro Leu 10350
Lys Arg 10355	Arg Lys Asp Lys Glu 10360	Glu Asp Val Glu Thr 10365
Ile Glu 10370	Ser Lys Asp Glu Asp 10375	Val Pro Leu Leu Leu 10380
Lys Asn 10385	Gln Lys Asp Asn Gln 10390	Ser Lys Asp Lys Lys 10395
Lys Asn 10400	Thr Ser Lys Lys Val 10405	Ala Ala Lys Lys Lys 10410
Ala Lys 10415	Lys Asn Lys Lys	

&lt;210&gt; SEQ ID NO 25

&lt;211&gt; LENGTH: 340

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 25

Met Lys Lys Lys Leu Leu Val Leu Thr Met Ser Thr Leu Phe Ala Thr 1 5 10 15
Gln Ile Met Asn Ser Asn His Ala Lys Ala Ser Val Thr Glu Ser Val 20 25 30
Asp Lys Lys Phe Val Val Pro Glu Ser Gly Ile Asn Lys Ile Ile Pro 35 40 45
Ala Tyr Asp Glu Phe Lys Asn Ser Pro Lys Val Asn Val Ser Asn Leu 50 55 60
Thr Asp Asn Lys Asn Phe Val Ala Ser Glu Asp Lys Leu Asn Lys Ile 65 70 75 80
Ala Asp Ser Ser Ala Ala Ser Lys Ile Val Asp Lys Asn Phe Val Val 85 90 95

-continued

---

Pro Glu Ser Lys Leu Gly Asn Ile Val Pro Glu Tyr Lys Glu Ile Asn  
 100 105 110  
 Asn Arg Val Asn Val Ala Thr Asn Asn Pro Ala Ser Gln Gln Val Asp  
 115 120 125  
 Lys His Phe Val Ala Lys Gly Pro Glu Val Asn Arg Phe Ile Thr Gln  
 130 135 140  
 Asn Lys Val Asn His His Phe Ile Thr Thr Gln Thr His Tyr Lys Lys  
 145 150 155 160  
 Val Ile Thr Ser Tyr Lys Ser Thr His Val His Lys His Val Asn His  
 165 170 175  
 Ala Lys Asp Ser Ile Asn Lys His Phe Ile Val Lys Pro Ser Glu Ser  
 180 185 190  
 Pro Arg Tyr Thr His Pro Ser Gln Ser Leu Ile Ile Lys His His Phe  
 195 200 205  
 Ala Val Pro Gly Tyr His Ala His Lys Phe Val Thr Pro Gly His Ala  
 210 215 220  
 Ser Ile Lys Ile Asn His Phe Cys Val Val Pro Gln Ile Asn Ser Phe  
 225 230 235 240  
 Lys Val Ile Pro Pro Tyr Gly His Asn Ser His Arg Met His Val Pro  
 245 250 255  
 Ser Phe Gln Asn Asn Thr Thr Ala Thr His Gln Asn Ala Lys Val Asn  
 260 265 270  
 Lys Ala Tyr Asp Tyr Lys Tyr Phe Tyr Ser Tyr Lys Val Val Lys Gly  
 275 280 285  
 Val Lys Lys Tyr Phe Ser Phe Ser Gln Ser Asn Gly Tyr Lys Ile Gly  
 290 295 300  
 Lys Pro Ser Leu Asn Ile Lys Asn Val Asn Tyr Gln Tyr Ala Val Pro  
 305 310 315 320  
 Ser Tyr Ser Pro Thr His Tyr Val Pro Glu Phe Lys Gly Ser Leu Pro  
 325 330 335  
 Ala Pro Arg Val  
 340

&lt;210&gt; SEQ ID NO 26

&lt;211&gt; LENGTH: 130

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 26

Met Asn Phe Asn Asp Ile Glu Thr Met Val Lys Ser Lys Phe Lys Asp  
 1 5 10 15  
 Ile Lys Lys His Ala Glu Glu Ile Ala His Glu Ile Glu Val Arg Ser  
 20 25 30  
 Gly Tyr Leu Arg Lys Ala Glu Gln Tyr Lys Arg Leu Glu Phe Asn Leu  
 35 40 45  
 Ser Phe Ala Leu Asp Asp Ile Glu Ser Thr Ala Lys Asp Val Gln Thr  
 50 55 60  
 Ala Lys Ser Ser Ala Asn Lys Asp Ser Val Thr Val Lys Gly Lys Ala  
 65 70 75 80  
 Pro Asn Thr Leu Tyr Ile Glu Lys Arg Asn Leu Met Lys Gln Lys Leu  
 85 90 95  
 Glu Met Leu Gly Glu Asp Ile Asp Lys Asn Lys Glu Ser Leu Gln Lys  
 100 105 110  
 Ala Lys Glu Ile Ala Gly Glu Lys Ala Ser Glu Tyr Phe Asn Lys Ala  
 115 120 125

-continued

Met Asn  
130

<210> SEQ ID NO 27  
<211> LENGTH: 636  
<212> TYPE: PRT  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 27

Met Lys Lys Gln Ile Ile Ser Leu Gly Ala Leu Ala Val Ala Ser Ser  
1 5 10 15  
Leu Phe Thr Trp Asp Asn Lys Ala Asp Ala Ile Val Thr Lys Asp Tyr  
20 25 30  
Ser Gly Lys Ser Gln Val Asn Ala Gly Ser Lys Asn Gly Thr Leu Ile  
35 40 45  
Asp Ser Arg Tyr Leu Asn Ser Ala Leu Tyr Tyr Leu Glu Asp Tyr Ile  
50 55 60  
Ile Tyr Ala Ile Gly Leu Thr Asn Lys Tyr Glu Tyr Gly Asp Asn Ile  
65 70 75 80  
Tyr Lys Glu Ala Lys Asp Arg Leu Leu Glu Lys Val Leu Arg Glu Asp  
85 90 95  
Gln Tyr Leu Leu Glu Arg Lys Lys Ser Gln Tyr Glu Asp Tyr Lys Gln  
100 105 110  
Trp Tyr Ala Asn Tyr Lys Lys Glu Asn Pro Arg Thr Asp Leu Lys Met  
115 120 125  
Ala Asn Phe His Lys Tyr Asn Leu Glu Glu Leu Ser Met Lys Glu Tyr  
130 135 140  
Asn Glu Leu Gln Asp Ala Leu Lys Arg Ala Leu Asp Asp Phe His Arg  
145 150 155 160  
Glu Val Lys Asp Ile Lys Asp Lys Asn Ser Asp Leu Lys Thr Phe Asn  
165 170 175  
Ala Ala Glu Glu Asp Lys Ala Thr Lys Glu Val Tyr Asp Leu Val Ser  
180 185 190  
Glu Ile Asp Thr Leu Val Val Ser Tyr Tyr Gly Asp Lys Asp Tyr Gly  
195 200 205  
Glu His Ala Lys Glu Leu Arg Ala Lys Leu Asp Leu Ile Leu Gly Asp  
210 215 220  
Thr Asp Asn Pro His Lys Ile Thr Asn Glu Arg Ile Lys Lys Glu Met  
225 230 235 240  
Ile Asp Asp Leu Asn Ser Ile Ile Asp Asp Phe Phe Met Glu Thr Lys  
245 250 255  
Gln Asn Arg Pro Lys Ser Ile Thr Lys Tyr Asn Pro Thr Thr His Asn  
260 265 270  
Tyr Lys Thr Asn Ser Asp Asn Lys Pro Asn Phe Asp Lys Leu Val Glu  
275 280 285  
Glu Thr Lys Lys Ala Val Lys Glu Ala Asp Asp Ser Trp Lys Lys Lys  
290 295 300  
Thr Val Lys Lys Tyr Gly Glu Thr Glu Thr Lys Ser Pro Val Val Lys  
305 310 315 320  
Glu Glu Lys Lys Val Glu Glu Pro Gln Ala Pro Lys Val Asp Asn Gln  
325 330 335  
Gln Glu Val Lys Thr Thr Ala Gly Lys Ala Glu Glu Thr Thr Gln Pro  
340 345 350  
Val Ala Gln Pro Leu Val Lys Ile Pro Gln Gly Thr Ile Thr Gly Glu

-continued

355					360					365					
Ile	Val	Lys	Gly	Pro	Glu	Tyr	Pro	Thr	Met	Glu	Asn	Lys	Thr	Val	Gln
370					375					380					
Gly	Glu	Ile	Val	Gln	Gly	Pro	Asp	Phe	Leu	Thr	Met	Glu	Gln	Ser	Gly
385					390					395					400
Pro	Ser	Leu	Ser	Asn	Asn	Tyr	Thr	Asn	Pro	Pro	Leu	Thr	Asn	Pro	Ile
				405					410					415	
Leu	Glu	Gly	Leu	Glu	Gly	Ser	Ser	Ser	Lys	Leu	Glu	Ile	Lys	Pro	Gln
			420					425					430		
Gly	Thr	Glu	Ser	Thr	Leu	Lys	Gly	Thr	Gln	Gly	Glu	Ser	Ser	Asp	Ile
		435					440					445			
Glu	Val	Lys	Pro	Gln	Ala	Thr	Glu	Thr	Thr	Glu	Ala	Ser	Gln	Tyr	Gly
		450					455					460			
Pro	Arg	Pro	Gln	Phe	Asn	Lys	Thr	Pro	Lys	Tyr	Val	Lys	Tyr	Arg	Asp
465					470					475					480
Ala	Gly	Thr	Gly	Ile	Arg	Glu	Tyr	Asn	Asp	Gly	Thr	Phe	Gly	Tyr	Glu
				485					490					495	
Ala	Arg	Pro	Arg	Phe	Asn	Lys	Pro	Ser	Glu	Thr	Asn	Ala	Tyr	Asn	Val
			500					505					510		
Thr	Thr	His	Ala	Asn	Gly	Gln	Val	Ser	Tyr	Gly	Ala	Arg	Pro	Thr	Tyr
		515					520					525			
Lys	Lys	Pro	Ser	Glu	Thr	Asn	Ala	Tyr	Asn	Val	Thr	Thr	His	Ala	Asn
		530					535					540			
Gly	Gln	Val	Ser	Tyr	Gly	Ala	Arg	Pro	Thr	Gln	Asn	Lys	Pro	Ser	Lys
545					550					555					560
Thr	Asn	Ala	Tyr	Asn	Val	Thr	Thr	His	Gly	Asn	Gly	Gln	Val	Ser	Tyr
				565					570					575	
Gly	Ala	Arg	Pro	Thr	Gln	Asn	Lys	Pro	Ser	Lys	Thr	Asn	Ala	Tyr	Asn
			580					585					590		
Val	Thr	Thr	His	Ala	Asn	Gly	Gln	Val	Ser	Tyr	Gly	Ala	Arg	Pro	Thr
		595					600					605			
Tyr	Lys	Lys	Pro	Ser	Lys	Thr	Asn	Ala	Tyr	Asn	Val	Thr	Thr	His	Ala
		610					615					620			
Asp	Gly	Thr	Ala	Thr	Tyr	Gly	Pro	Arg	Val	Thr	Lys				
625					630					635					

&lt;210&gt; SEQ ID NO 28

&lt;211&gt; LENGTH: 745

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 28

Ala	Glu	Gln	His	Thr	Pro	Met	Lys	Ala	His	Ala	Val	Thr	Thr	Ile	Asp
1				5					10					15	
Lys	Ala	Thr	Thr	Asp	Lys	Gln	Gln	Val	Pro	Pro	Thr	Lys	Glu	Ala	Ala
			20					25					30		
His	His	Ser	Gly	Lys	Glu	Ala	Ala	Thr	Asn	Val	Ser	Ala	Ser	Ala	Gln
			35				40					45			
Gly	Thr	Ala	Asp	Asp	Thr	Asn	Ser	Lys	Val	Thr	Ser	Asn	Ala	Pro	Ser
			50				55					60			
Asn	Lys	Pro	Ser	Thr	Val	Val	Ser	Thr	Lys	Val	Asn	Glu	Thr	Arg	Asp
65					70					75					80
Val	Asp	Thr	Gln	Gln	Ala	Ser	Thr	Gln	Lys	Pro	Thr	His	Thr	Ala	Thr
				85					90					95	

-continued

---

Phe Lys Leu Ser Asn Ala Lys Thr Ala Ser Leu Ser Pro Arg Met Phe  
 100 105 110  
 Ala Ala Asn Ala Pro Gln Thr Thr Thr His Lys Ile Leu His Thr Asn  
 115 120 125  
 Asp Ile His Gly Arg Leu Ala Glu Glu Lys Gly Arg Val Ile Gly Met  
 130 135 140  
 Ala Lys Leu Lys Thr Val Lys Glu Gln Glu Lys Pro Asp Leu Met Leu  
 145 150 155 160  
 Asp Ala Gly Asp Ala Phe Gln Gly Leu Pro Leu Ser Asn Gln Ser Lys  
 165 170 175  
 Gly Glu Glu Met Ala Lys Ala Met Asn Ala Val Gly Tyr Asp Ala Met  
 180 185 190  
 Ala Val Gly Asn His Glu Phe Asp Phe Gly Tyr Asp Gln Leu Lys Lys  
 195 200 205  
 Leu Glu Gly Met Leu Asp Phe Pro Met Leu Ser Thr Asn Val Tyr Lys  
 210 215 220  
 Asp Gly Lys Arg Ala Phe Lys Pro Ser Thr Ile Val Thr Lys Asn Gly  
 225 230 235 240  
 Ile Arg Tyr Gly Ile Ile Gly Val Thr Thr Pro Glu Thr Lys Thr Lys  
 245 250 255  
 Thr Arg Pro Glu Gly Ile Lys Gly Val Glu Phe Arg Asp Pro Leu Gln  
 260 265 270  
 Ser Val Thr Ala Glu Met Met Arg Ile Tyr Lys Asp Val Asp Thr Phe  
 275 280 285  
 Val Val Ile Ser His Leu Gly Ile Asp Pro Ser Thr Gln Glu Thr Trp  
 290 295 300  
 Arg Gly Asp Tyr Leu Val Lys Gln Leu Ser Gln Asn Pro Gln Leu Lys  
 305 310 315 320  
 Lys Arg Ile Thr Val Ile Asp Gly His Ser His Thr Val Leu Gln Asn  
 325 330 335  
 Gly Gln Ile Tyr Asn Asn Asp Ala Leu Ala Gln Thr Gly Thr Ala Leu  
 340 345 350  
 Ala Asn Ile Gly Lys Ile Thr Phe Asn Tyr Arg Asn Gly Glu Val Ser  
 355 360 365  
 Asn Ile Lys Pro Ser Leu Ile Asn Val Lys Asp Val Glu Asn Val Thr  
 370 375 380  
 Pro Asn Lys Ala Leu Ala Glu Gln Ile Asn Gln Ala Asp Gln Thr Phe  
 385 390 395 400  
 Arg Ala Gln Thr Ala Glu Val Ile Ile Pro Asn Asn Thr Ile Asp Phe  
 405 410 415  
 Lys Gly Glu Arg Asp Asp Val Arg Thr Arg Glu Thr Asn Leu Gly Asn  
 420 425 430  
 Ala Ile Ala Asp Ala Met Glu Ala Tyr Gly Val Lys Asn Phe Ser Lys  
 435 440 445  
 Lys Thr Asp Phe Ala Val Thr Asn Gly Gly Gly Ile Arg Ala Ser Ile  
 450 455 460  
 Ala Lys Gly Lys Val Thr Arg Tyr Asp Leu Ile Ser Val Leu Pro Phe  
 465 470 475 480  
 Gly Asn Thr Ile Ala Gln Ile Asp Val Lys Gly Ser Asp Val Trp Thr  
 485 490 495  
 Ala Phe Glu His Ser Leu Gly Ala Pro Thr Thr Gln Lys Asp Gly Lys  
 500 505 510  
 Thr Val Leu Thr Ala Asn Gly Gly Leu Leu His Ile Ser Asp Ser Ile

-continued

515					520					525					
Arg	Val	Tyr	Tyr	Asp	Ile	Asn	Lys	Pro	Ser	Gly	Lys	Arg	Ile	Asn	Ala
530						535					540				
Ile	Gln	Ile	Leu	Asn	Lys	Glu	Thr	Gly	Lys	Phe	Glu	Asn	Ile	Asp	Leu
545				550						555					560
Lys	Arg	Val	Tyr	His	Val	Thr	Met	Asn	Asp	Phe	Thr	Ala	Ser	Gly	Gly
				565					570					575	
Asp	Gly	Tyr	Ser	Met	Phe	Gly	Gly	Pro	Arg	Glu	Glu	Gly	Ile	Ser	Leu
			580					585					590		
Asp	Gln	Val	Leu	Ala	Ser	Tyr	Leu	Lys	Thr	Ala	Asn	Leu	Ala	Lys	Tyr
		595					600					605			
Asp	Thr	Thr	Glu	Pro	Gln	Arg	Met	Leu	Leu	Gly	Lys	Pro	Ala	Val	Ser
	610					615					620				
Glu	Gln	Pro	Ala	Lys	Gly	Gln	Gln	Gly	Ser	Lys	Gly	Ser	Lys	Ser	Gly
625				630						635					640
Lys	Asp	Thr	Gln	Pro	Ile	Gly	Asp	Asp	Lys	Val	Met	Asp	Pro	Ala	Lys
			645						650					655	
Lys	Pro	Ala	Pro	Gly	Lys	Val	Val	Leu	Leu	Leu	Ala	His	Arg	Gly	Thr
		660						665					670		
Val	Ser	Ser	Gly	Thr	Glu	Gly	Ser	Gly	Arg	Thr	Ile	Glu	Gly	Ala	Thr
		675					680					685			
Val	Ser	Ser	Lys	Ser	Gly	Lys	Gln	Leu	Ala	Arg	Met	Ser	Val	Pro	Lys
		690				695					700				
Gly	Ser	Ala	His	Glu	Lys	Gln	Leu	Pro	Lys	Thr	Gly	Thr	Asn	Gln	Ser
705				710						715					720
Ser	Ser	Pro	Glu	Ala	Met	Phe	Val	Leu	Leu	Ala	Gly	Ile	Gly	Leu	Ile
			725					730						735	
Ala	Thr	Val	Arg	Arg	Arg	Lys	Ala	Ser							
		740					745								

&lt;210&gt; SEQ ID NO 29

&lt;211&gt; LENGTH: 628

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 29

Met	Ser	Asp	Arg	Phe	Ile	Lys	Phe	Asn	Asp	Glu	Gln	Leu	Asp	Ala	Lys
1				5					10					15	
Gln	Val	Met	Met	Leu	Gln	Asp	Leu	Ala	Arg	Leu	Leu	Leu	Lys	Asn	Glu
		20					25						30		
Gln	Thr	Gln	Val	Lys	Ile	Gln	Lys	Phe	Pro	Tyr	Tyr	Asn	Pro	Val	Gln
		35					40					45			
Asn	Val	Leu	Ile	Thr	Ser	Trp	Phe	Trp	Ser	His	Arg	Pro	Ser	His	Ile
		50				55					60				
Glu	Met	Ala	Gly	Leu	Lys	Thr	Asp	Val	Met	Leu	Ala	Ala	Tyr	Gly	Tyr
65					70						75				80
His	Met	Met	Asp	Val	Gln	Ile	Val	Asn	Glu	Val	Val	Gln	Asp	Lys	Thr
				85					90					95	
Phe	Lys	His	Pro	Lys	Phe	Tyr	Gln	Gln	Leu	Phe	Lys	Leu	Leu	Glu	Asp
			100					105						110	
Met	Arg	Val	Leu	Asn	Ser	Ile	Lys	Val	Glu	Arg	Pro	Ser	Thr	Ala	Lys
		115					120						125		
Leu	Ile	Asp	Leu	Arg	Leu	Asp	Thr	Arg	Ile	Ser	Tyr	Thr	Glu	Ser	Gln
		130				135							140		

-continued

---

Ile Lys Val Tyr Arg Thr Lys Thr Gln Tyr Thr Asp Leu Leu Phe Leu  
 145 150 155 160  
 Tyr Leu Glu His Ala Phe Leu Ser Gln Asp Phe Phe Asp Ile Pro Ser  
 165 170 175  
 Ile His Ser Asp Leu Asp Asp Ile Leu Val Asn Met Phe Leu Tyr Leu  
 180 185 190  
 Pro Asn Phe Phe Gln Asn Gln Asn Ser Glu Asp Asn Met Tyr Leu Ala  
 195 200 205  
 Gln Arg Ile Met Tyr Gln Val Asp Asp Ile Leu Lys Glu Asp Met Leu  
 210 215 220  
 Asn Glu Tyr Tyr Tyr Leu Pro Lys Thr Leu Tyr Asn Thr Leu Ala Ser  
 225 230 235 240  
 Pro Glu Phe Asp Asp Leu Lys Arg Thr Asp Ala Ser Gln Val Asp Gly  
 245 250 255  
 Gln Asp Asp Thr Ser Glu Asp Asp Asp Asn Glu Ser Glu Lys Ala Asp  
 260 265 270  
 Ser Lys Ser Ala Asp Ser Glu Ser Lys Gly Gly Ala Tyr Leu Glu Met  
 275 280 285  
 Glu Leu His Glu Gly Gln Asn Ser Glu Thr Leu Gly Asn Asp Glu Ala  
 290 295 300  
 Arg Glu Gly Asp Ala Thr Asp Asp Met Thr Asp Met Met Thr Lys Lys  
 305 310 315 320  
 Gly Lys Gly Ser Asn Asp Thr Leu Asn Arg Glu Glu Gly Asp Ala Val  
 325 330 335  
 Gly Gln Ser Gln Ala Phe Gln Leu Asp Gly Val Asn Lys Asn Val Glu  
 340 345 350  
 Ile Lys Trp Gln Ile Pro Glu Ile Glu Pro Gln Tyr Val Leu Glu Tyr  
 355 360 365  
 Gln Glu Ser Lys Gln Asp Val Gln Tyr Glu Ile Lys Asp Leu Ile Gln  
 370 375 380  
 Ile Ile Lys Lys Thr Ile Glu Arg Glu Gln Arg Asp Ala Arg Phe Asn  
 385 390 395 400  
 Leu Thr Lys Gly Arg Leu Gln Lys Asp Leu Ile Asn Trp Phe Ile Asp  
 405 410 415  
 Asp Gln Tyr Lys Leu Phe Tyr Lys Lys Gln Asp Leu Ser Lys Ser Phe  
 420 425 430  
 Asp Ala Thr Phe Thr Leu Leu Ile Asp Ala Ser Ala Ser Met His Asp  
 435 440 445  
 Lys Met Ala Glu Thr Lys Lys Gly Val Val Leu Phe His Glu Thr Leu  
 450 455 460  
 Lys Ala Leu Asn Ile Lys His Glu Ile Leu Ser Phe Ser Glu Asp Ala  
 465 470 475 480  
 Phe Asp Ser Asp Glu His Ala Gln Pro Asn Ile Ile Asn Glu Ile Ile  
 485 490 495  
 Asn Tyr Asp Tyr Ser Thr Phe Glu Lys Asp Gly Pro Arg Ile Met Ala  
 500 505 510  
 Leu Glu Pro Gln Asp Asp Asn Arg Asp Gly Val Ala Ile Arg Val Ala  
 515 520 525  
 Ser Glu Arg Leu Met Arg Arg Asn Gln His Gln Arg Phe Leu Ile Val  
 530 535 540  
 Phe Ser Asp Gly Glu Pro Ser Ala Phe Asn Tyr Ser Gln Asp Gly Ile  
 545 550 555 560  
 Ile Asp Thr Tyr Glu Ala Val Glu Met Ser Arg Lys Phe Gly Ile Glu

-continued

---

```

          565              570              575
Val Phe Asn Val Phe Leu Ser Gln Asp Pro Ile Thr Glu Asp Val Glu
      580              585              590
Gln Thr Ile His Asn Ile Tyr Gly Gln Tyr Ala Ile Phe Val Glu Gly
      595              600              605
Val Ala His Leu Pro Gly His Leu Ser Pro Leu Leu Lys Lys Leu Leu
      610              615              620
Leu Lys Ser Leu
      625

```

```

<210> SEQ ID NO 30
<211> LENGTH: 154
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus sp.

```

```

<400> SEQUENCE: 30

```

```

Ala Glu Ile Asn Lys Gln Thr Thr Ser Gln Gly Val Thr Thr Glu Lys
 1              5              10              15
Asn Asn Gly Ile Ala Val Leu Glu Gln Asp Val Ile Thr Pro Thr Val
 20              25              30
Lys Pro Gln Ala Lys Gln Asp Ile Ile Gln Ala Val Thr Thr Arg Lys
 35              40              45
Gln Gln Ile Lys Lys Ser Asn Ala Ser Leu Gln Asp Glu Lys Asp Val
 50              55              60
Ala Asn Asp Lys Ile Gly Lys Ile Glu Thr Lys Ala Ile Lys Asp Ile
 65              70              75              80
Asp Ala Ala Thr Thr Asn Ala Gln Val Glu Ala Ile Lys Thr Lys Ala
 85              90              95
Ile Asn Asp Ile Asn Gln Thr Thr Pro Ala Thr Thr Ala Lys Ala Ala
 100             105             110
Ala Leu Glu Glu Phe Asp Glu Val Val Gln Ala Gln Ile Asp Gln Ala
 115             120             125
Pro Leu Asn Pro Asp Thr Thr Asn Glu Glu Val Ala Glu Ala Ile Glu
 130             135             140
Arg Ile Asn Ala Ala Lys Val Ser Gly Val
 145             150

```

```

<210> SEQ ID NO 31
<211> LENGTH: 584
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus sp.

```

```

<400> SEQUENCE: 31

```

```

Met Lys Phe Lys Ser Leu Ile Thr Thr Thr Leu Ala Leu Gly Val Leu
 1              5              10              15
Ala Ser Thr Gly Ala Asn Phe Asn Asn Asn Glu Ala Ser Ala Ala Ala
 20              25              30
Lys Pro Leu Asp Lys Ser Ser Ser Ser Leu His His Gly Tyr Ser Lys
 35              40              45
Val His Val Pro Tyr Ala Ile Thr Val Asn Gly Thr Ser Gln Asn Ile
 50              55              60
Leu Ser Ser Leu Thr Phe Asn Lys Asn Gln Asn Ile Ser Tyr Lys Asp
 65              70              75              80
Leu Glu Asp Arg Val Lys Ser Val Leu Lys Ser Asp Arg Gly Ile Ser
 85              90              95
Asp Ile Asp Leu Arg Leu Ser Lys Gln Ala Lys Tyr Thr Val Tyr Phe

```

-continued

100				105				110							
Lys	Asn	Gly	Thr	Lys	Lys	Val	Ile	Asp	Leu	Lys	Ala	Gly	Ile	Tyr	Thr
	115						120					125			
Ala	Asp	Leu	Ile	Asn	Thr	Ser	Glu	Ile	Lys	Ala	Ile	Asn	Ile	Asn	Val
	130					135					140				
Asp	Thr	Lys	Lys	Gln	Val	Glu	Asp	Lys	Lys	Lys	Asp	Lys	Ala	Asn	Tyr
	145				150					155					160
Gln	Val	Pro	Tyr	Thr	Ile	Thr	Val	Asn	Gly	Thr	Ser	Gln	Asn	Ile	Leu
				165					170						175
Ser	Asn	Leu	Thr	Phe	Asn	Lys	Asn	Gln	Asn	Ile	Ser	Tyr	Lys	Asp	Leu
		180							185				190		
Glu	Asp	Lys	Val	Lys	Ser	Val	Leu	Glu	Ser	Asn	Arg	Gly	Ile	Thr	Asp
		195					200					205			
Val	Asp	Leu	Arg	Leu	Ser	Lys	Gln	Ala	Lys	Tyr	Thr	Val	Asn	Phe	Lys
	210					215					220				
Asn	Gly	Thr	Lys	Lys	Val	Ile	Asp	Leu	Lys	Ser	Gly	Ile	Tyr	Thr	Ala
	225				230					235					240
Asn	Leu	Ile	Asn	Ser	Ser	Asp	Ile	Lys	Ser	Ile	Asn	Ile	Asn	Val	Asp
			245						250					255	
Thr	Lys	Lys	His	Ile	Glu	Asn	Lys	Ala	Lys	Arg	Asn	Tyr	Gln	Val	Pro
			260						265					270	
Tyr	Ser	Ile	Asn	Leu	Asn	Gly	Thr	Ser	Thr	Asn	Ile	Leu	Ser	Asn	Leu
		275				280						285			
Ser	Phe	Ser	Asn	Lys	Pro	Trp	Thr	Asn	Tyr	Lys	Asn	Leu	Thr	Ser	Gln
	290					295					300				
Ile	Lys	Ser	Val	Leu	Lys	His	Asp	Arg	Gly	Ile	Ser	Glu	Gln	Asp	Leu
	305				310					315					320
Lys	Tyr	Ala	Lys	Lys	Ala	Tyr	Tyr	Thr	Val	Tyr	Phe	Lys	Asn	Gly	Gly
			325						330					335	
Lys	Arg	Ile	Leu	Gln	Leu	Asn	Ser	Lys	Asn	Tyr	Thr	Ala	Asn	Leu	Val
		340							345				350		
His	Ala	Lys	Asp	Val	Lys	Arg	Ile	Glu	Ile	Thr	Val	Lys	Thr	Gly	Thr
		355					360					365			
Lys	Ala	Lys	Ala	Asp	Arg	Tyr	Val	Pro	Tyr	Thr	Ile	Ala	Val	Asn	Gly
		370				375					380				
Thr	Ser	Thr	Pro	Ile	Leu	Ser	Asp	Leu	Lys	Phe	Thr	Gly	Asp	Pro	Arg
	385				390					395				400	
Val	Gly	Tyr	Lys	Asp	Ile	Ser	Lys	Lys	Val	Lys	Ser	Val	Leu	Lys	His
			405						410					415	
Asp	Arg	Gly	Ile	Gly	Glu	Arg	Glu	Leu	Lys	Tyr	Ala	Lys	Lys	Ala	Thr
		420					425						430		
Tyr	Thr	Val	His	Phe	Lys	Asn	Gly	Thr	Lys	Lys	Val	Ile	Asn	Ile	Asn
		435				440						445			
Ser	Asn	Ile	Ser	Gln	Leu	Asn	Leu	Leu	Tyr	Val	Gln	Asp	Ile	Lys	Lys
	450					455					460				
Ile	Asp	Ile	Asp	Val	Lys	Thr	Gly	Thr	Lys	Ala	Lys	Ala	Asp	Ser	Tyr
	465				470					475				480	
Val	Pro	Tyr	Thr	Ile	Ala	Val	Asn	Gly	Thr	Ser	Thr	Pro	Ile	Leu	Ser
			485						490					495	
Lys	Leu	Lys	Ile	Ser	Asn	Lys	Gln	Leu	Ile	Ser	Tyr	Lys	Tyr	Leu	Asn
		500							505				510		
Asp	Lys	Val	Lys	Ser	Val	Leu	Lys	Ser	Glu	Arg	Gly	Ile	Ser	Asp	Leu
		515					520					525			

-continued

---

Asp Leu Lys Phe Ala Lys Gln Ala Lys Tyr Thr Val Tyr Phe Lys Asn  
 530 535 540

Gly Lys Lys Gln Val Val Asn Leu Lys Ser Asp Ile Phe Thr Pro Asn  
 545 550 555 560

Leu Phe Ser Ala Lys Asp Ile Lys Lys Ile Asp Ile Asp Val Lys Gln  
 565 570 575

Tyr Thr Lys Ser Lys Lys Asn Lys  
 580

<210> SEQ ID NO 32  
 <211> LENGTH: 508  
 <212> TYPE: PRT  
 <213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 32

Met Lys Asn Lys Leu Leu Val Leu Ser Leu Gly Ala Leu Cys Val Ser  
 1 5 10 15

Gln Ile Trp Glu Ser Asn Arg Ala Ser Ala Val Val Ser Gly Glu Lys  
 20 25 30

Asn Pro Tyr Val Ser Glu Ser Leu Lys Leu Thr Asn Asn Lys Asn Lys  
 35 40 45

Ser Arg Thr Val Glu Glu Tyr Lys Lys Ser Leu Asp Asp Leu Ile Trp  
 50 55 60

Ser Phe Pro Asn Leu Asp Asn Glu Arg Phe Asp Asn Pro Glu Tyr Lys  
 65 70 75 80

Glu Ala Met Lys Lys Tyr Gln Gln Arg Phe Met Ala Glu Asp Glu Ala  
 85 90 95

Leu Lys Lys Phe Phe Ser Glu Glu Lys Lys Ile Lys Asn Gly Asn Thr  
 100 105 110

Asp Asn Leu Asp Tyr Leu Gly Leu Ser His Glu Arg Tyr Glu Ser Val  
 115 120 125

Phe Asn Thr Leu Lys Lys Gln Ser Glu Glu Phe Leu Lys Glu Ile Glu  
 130 135 140

Asp Ile Lys Lys Asp Asn Pro Glu Leu Lys Asp Phe Asn Glu Glu Glu  
 145 150 155 160

Gln Leu Lys Cys Asp Leu Glu Leu Asn Lys Leu Glu Asn Gln Ile Leu  
 165 170 175

Met Leu Gly Lys Thr Phe Tyr Gln Asn Tyr Arg Asp Asp Val Glu Ser  
 180 185 190

Leu Tyr Ser Lys Leu Asp Leu Ile Met Gly Tyr Lys Asp Glu Glu Arg  
 195 200 205

Ala Asn Lys Lys Ala Val Asn Lys Arg Met Leu Glu Asn Lys Lys Glu  
 210 215 220

Asp Leu Glu Thr Ile Ile Asp Glu Phe Phe Ser Asp Ile Asp Lys Thr  
 225 230 235 240

Arg Pro Asn Asn Ile Pro Val Leu Glu Asp Glu Lys Gln Glu Glu Lys  
 245 250 255

Asn His Lys Asn Met Ala Gln Leu Lys Ser Asp Thr Glu Ala Ala Lys  
 260 265 270

Ser Asp Glu Ser Lys Arg Ser Lys Arg Ser Lys Arg Ser Leu Asn Thr  
 275 280 285

Gln Asn His Lys Pro Ala Ser Gln Glu Val Ser Glu Gln Gln Lys Ala  
 290 295 300

Glu Tyr Asp Lys Arg Ala Glu Glu Arg Lys Ala Arg Phe Leu Asp Asn

-continued

---

```

305                310                315                320
Gln Lys Ile Lys Lys Thr Pro Val Val Ser Leu Glu Tyr Asp Phe Glu
      325                330                335
His Lys Gln Arg Ile Asp Asn Glu Asn Asp Lys Lys Leu Val Val Ser
      340                345                350
Ala Pro Thr Lys Lys Pro Thr Ser Pro Thr Thr Tyr Thr Glu Thr Thr
      355                360                365
Thr Gln Val Pro Met Pro Thr Val Glu Arg Gln Thr Gln Gln Gln Ile
      370                375                380
Ile Tyr Asn Ala Pro Lys Gln Leu Ala Gly Leu Asn Gly Glu Ser His
385                390                395                400
Asp Phe Thr Thr Thr His Gln Ser Pro Thr Thr Ser Asn His Thr His
      405                410                415
Asn Asn Val Val Glu Phe Glu Glu Thr Ser Ala Leu Pro Gly Arg Lys
      420                425                430
Ser Gly Ser Leu Val Gly Ile Ser Gln Ile Asp Ser Ser His Leu Thr
      435                440                445
Glu Arg Glu Lys Arg Val Ile Lys Arg Glu His Val Arg Glu Ala Gln
      450                455                460
Lys Leu Val Asp Asn Tyr Lys Asp Thr His Ser Tyr Lys Asp Arg Ile
465                470                475                480
Asn Ala Gln Gln Lys Val Asn Thr Leu Ser Glu Gly His Gln Lys Arg
      485                490                495
Phe Asn Lys Gln Ile Asn Lys Val Tyr Asn Gly Lys
      500                505

```

&lt;210&gt; SEQ ID NO 33

&lt;211&gt; LENGTH: 520

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 33

```

Met Leu Thr Leu Gln Ile His Thr Gly Gly Ile Asn Leu Lys Lys Lys
1                5                10                15
Asn Ile Tyr Ser Ile Arg Lys Leu Gly Val Gly Ile Ala Ser Val Thr
      20                25                30
Leu Gly Thr Leu Leu Ile Ser Gly Gly Val Thr Pro Ala Ala Asn Ala
      35                40                45
Ala Gln His Asp Glu Ala Gln Gln Asn Ala Phe Tyr Gln Val Leu Asn
      50                55                60
Met Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe Ile Gln Ser Leu
65                70                75                80
Lys Asp Asp Pro Ser Gln Ser Ala Asn Val Leu Gly Glu Ala Gln Lys
      85                90                95
Leu Asn Asp Ser Gln Ala Pro Lys Ala Asp Ala Gln Gln Asn Asn Phe
      100                105                110
Asn Lys Asp Gln Gln Ser Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn
      115                120                125
Leu Asn Glu Ala Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp
      130                135                140
Pro Ser Gln Ser Thr Asn Val Leu Gly Glu Ala Lys Lys Leu Asn Glu
145                150                155                160
Ser Gln Ala Pro Lys Ala Asp Asn Asn Phe Asn Lys Glu Gln Gln Asn
      165                170                175

```

-continued

---

Ala Phe Tyr Glu Ile Leu Asn Met Pro Asn Leu Asn Glu Glu Gln Arg  
180 185 190

Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn  
195 200 205

Leu Leu Ser Glu Ala Lys Lys Leu Asn Glu Ser Gln Ala Pro Lys Ala  
210 215 220

Asp Asn Lys Phe Asn Lys Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu  
225 230 235 240

His Leu Pro Asn Leu Asn Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser  
245 250 255

Leu Lys Asp Asp Pro Ser Gln Ser Ala Asn Leu Leu Ala Glu Ala Lys  
260 265 270

Lys Leu Asn Asp Ala Gln Ala Pro Lys Ala Asp Asn Lys Phe Asn Lys  
275 280 285

Glu Gln Gln Asn Ala Phe Tyr Glu Ile Leu His Leu Pro Asn Leu Thr  
290 295 300

Glu Glu Gln Arg Asn Gly Phe Ile Gln Ser Leu Lys Asp Asp Pro Ser  
305 310 315 320

Val Ser Lys Glu Ile Leu Ala Glu Ala Lys Lys Leu Asn Asp Ala Gln  
325 330 335

Ala Pro Lys Glu Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp Gly Asn  
340 345 350

Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp Asn Lys  
355 360 365

Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp Asn Asn  
370 375 380

Lys Pro Gly Lys Glu Asp Gly Asn Lys Pro Gly Lys Glu Asp Asn Lys  
385 390 395 400

Lys Pro Gly Lys Glu Asp Asn Asn Lys Pro Gly Lys Glu Asp Gly Asn  
405 410 415

Lys Pro Gly Lys Glu Asp Gly Asn Gly Val His Val Val Lys Pro Gly  
420 425 430

Asp Thr Val Asn Asp Ile Ala Lys Ala Asn Gly Thr Thr Ala Asp Lys  
435 440 445

Ile Ala Ala Asp Asn Lys Leu Ala Asp Lys Asn Met Ile Lys Pro Gly  
450 455 460

Gln Glu Leu Val Val Asp Lys Lys Gln Pro Ala Asn His Ala Asp Ala  
465 470 475 480

Asn Lys Ala Gln Ala Leu Pro Glu Thr Gly Glu Glu Asn Pro Phe Ile  
485 490 495

Gly Thr Thr Val Phe Gly Gly Leu Ser Leu Ala Leu Gly Ala Ala Leu  
500 505 510

Leu Ala Gly Arg Arg Arg Glu Leu  
515 520

&lt;210&gt; SEQ ID NO 34

&lt;211&gt; LENGTH: 291

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 34

Ala Gln His Asp Glu Ala Lys Lys Asn Ala Phe Tyr Gln Val Leu Asn  
1 5 10 15

Met Pro Asn Leu Asn Ala Asp Gln Arg Asn Gly Phe Ile Gln Ser Leu  
20 25 30



-continued

Met Gly Ser Ser His His His His His His Ser Ser Gly Leu Val Pro  
 1 5 10 15

Arg Gly Ser

<210> SEQ ID NO 38

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 38

aacatatggt caacaagat caacaagc

29

<210> SEQ ID NO 39

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 39

aaggatccag attcgtttaa ttttttagc

29

<210> SEQ ID NO 40

<211> LENGTH: 43

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 40

cttcattcaa agtcttaaag cgccecaag ccaaagcact aac

43

<210> SEQ ID NO 41

<211> LENGTH: 43

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 41

gtttagtgett tggettgggg cggetttaa gactttgaatg aag

43

<210> SEQ ID NO 42

<211> LENGTH: 42

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 42

catatgttca acaaagataa aaaaagcgcc ttctatgaaa tc

42

<210> SEQ ID NO 43

<211> LENGTH: 42

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 43

gatttcatag aaggcgcttt ttttatcttt gttgaacata tg

42

<210> SEQ ID NO 44

<211> LENGTH: 42

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus sp.

-continued

---

<400> SEQUENCE: 44  
catatgttca acaaagatgg aggaagcgcc ttctatgaaa tc 42

<210> SEQ ID NO 45  
<211> LENGTH: 42  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 45  
gatttcatag aaggcgcttc ctccatcttt gttgaacata tg 42

<210> SEQ ID NO 46  
<211> LENGTH: 52  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 46  
ggggacaagt ttgtacaaaa aagcaggctg atgactaagt tgaaaaaaga ag 52

<210> SEQ ID NO 47  
<211> LENGTH: 28  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 47  
aaggatcccc tccaaaatgt aattgccc 28

<210> SEQ ID NO 48  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 48  
aaggatccgt ttgtaactct atccaaagac 30

<210> SEQ ID NO 49  
<211> LENGTH: 49  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 49  
ggggaccact ttgtacaaga aagctgggtg acacctattg cacgattcg 49

<210> SEQ ID NO 50  
<211> LENGTH: 50  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 50  
ggggacaagt ttgtacaaaa aagcaggctc agatagcgat tcagattcag 50

<210> SEQ ID NO 51  
<211> LENGTH: 31  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 51  
aaggatccct gtattttctc cttaattttc c 31

<210> SEQ ID NO 52  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

-continued

---

<400> SEQUENCE: 52  
aaggatccca tggctgcaaa gcaaataatg 30

<210> SEQ ID NO 53  
<211> LENGTH: 51  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 53  
ggggaccact ttgtacaaga aagctgggtg ccttgggtga acaaatttat g 51

<210> SEQ ID NO 54  
<211> LENGTH: 37  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 54  
gaaggatccg tttattctag ttaatatata gttaatg 37

<210> SEQ ID NO 55  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 55  
gaactgcagc tgtatgtctt tggatagagt tac 33

<210> SEQ ID NO 56  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 56  
gaaggatccg gtggcttttt tacttggatt ttc 33

<210> SEQ ID NO 57  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 57  
gaactgcagc gacaaactca ttatttgctt tgc 33

<210> SEQ ID NO 58  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 58  
gaactcgagt ctagcttatt tacatgg 27

<210> SEQ ID NO 59  
<211> LENGTH: 45  
<212> TYPE: DNA  
<213> ORGANISM: Staphylococcus sp.

<400> SEQUENCE: 59  
gaactcgaga tagaaggcag aatagtaaca aaggattata gtggg 45

<210> SEQ ID NO 60  
<211> LENGTH: 27  
<212> TYPE: DNA

-continued

---

 <213> ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 60

gtaggatcct gggatagagt tacaaac

27

&lt;210&gt; SEQ ID NO 61

&lt;211&gt; LENGTH: 34

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 61

gaactcgagg cattatgtgt atcacaatt tggg

34

&lt;210&gt; SEQ ID NO 62

&lt;211&gt; LENGTH: 43

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 62

gaactcgaga tagaaggcag agtggtttct ggggagaaga atc

43

&lt;210&gt; SEQ ID NO 63

&lt;211&gt; LENGTH: 33

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 63

gaactcgagg cagccatgca ttaattattt gcc

33

&lt;210&gt; SEQ ID NO 64

&lt;211&gt; LENGTH: 677

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Staphylococcus sp.

&lt;400&gt; SEQUENCE: 64

Met	Lys	Ser	Asn	Leu	Arg	Tyr	Gly	Ile	Arg	Lys	His	Lys	Leu	Gly	Ala
1			5						10					15	
Ala	Ser	Val	Phe	Leu	Gly	Thr	Met	Ile	Val	Val	Gly	Met	Gly	Gln	Glu
			20				25						30		
Lys	Glu	Ala	Ala	Ala	Ser	Glu	Gln	Asn	Asn	Thr	Thr	Val	Glu	Glu	Ser
		35					40					45			
Gly	Ser	Ser	Ala	Thr	Glu	Ser	Lys	Ala	Ser	Glu	Thr	Gln	Thr	Thr	Thr
		50				55					60				
Asn	Asn	Val	Asn	Thr	Ile	Asp	Glu	Thr	Gln	Ser	Tyr	Ser	Ala	Thr	Ser
65				70						75				80	
Thr	Glu	Gln	Pro	Ser	Gln	Ser	Thr	Gln	Val	Thr	Thr	Glu	Glu	Ala	Pro
			85						90					95	
Lys	Thr	Val	Gln	Ala	Pro	Lys	Val	Glu	Thr	Ser	Arg	Val	Asp	Leu	Pro
		100						105					110		
Ser	Glu	Lys	Val	Ala	Asp	Lys	Glu	Thr	Thr	Gly	Thr	Gln	Val	Asp	Ile
		115					120					125			
Ala	Gln	Pro	Ser	Asn	Val	Ser	Glu	Ile	Lys	Pro	Arg	Met	Lys	Arg	Ser
		130				135					140				
Thr	Asp	Val	Thr	Ala	Val	Ala	Glu	Lys	Glu	Val	Val	Glu	Glu	Thr	Lys
145					150					155				160	
Ala	Thr	Gly	Thr	Asp	Val	Thr	Asn	Lys	Val	Glu	Val	Glu	Glu	Gly	Ser
				165					170					175	
Glu	Ile	Val	Gly	His	Lys	Gln	Asp	Thr	Asn	Val	Val	Asn	Pro	His	Asn
		180						185					190		

-continued

---

Ala Glu Arg Val Thr Leu Lys Tyr Lys Trp Lys Phe Gly Glu Gly Ile  
 195 200 205

Lys Ala Gly Asp Tyr Phe Asp Phe Thr Leu Ser Asp Asn Val Glu Thr  
 210 215 220

His Gly Ile Ser Thr Leu Arg Lys Val Pro Glu Ile Lys Ser Thr Asp  
 225 230 235 240

Gly Gln Val Met Ala Thr Gly Glu Ile Ile Gly Glu Arg Lys Val Arg  
 245 250 255

Tyr Thr Phe Lys Glu Tyr Val Gln Glu Lys Lys Asp Leu Thr Ala Glu  
 260 265 270

Leu Ser Leu Asn Leu Phe Ile Asp Pro Thr Thr Val Thr Gln Lys Gly  
 275 280 285

Asn Gln Asn Val Glu Val Lys Leu Gly Glu Thr Thr Val Ser Lys Ile  
 290 295 300

Phe Asn Ile Gln Tyr Leu Gly Gly Val Arg Asp Asn Trp Gly Val Thr  
 305 310 315 320

Ala Asn Gly Arg Ile Asp Thr Leu Asn Lys Val Asp Gly Lys Phe Ser  
 325 330 335

His Phe Ala Tyr Met Lys Pro Asn Asn Gln Ser Leu Ser Ser Val Thr  
 340 345 350

Val Thr Gly Gln Val Thr Lys Gly Asn Lys Pro Gly Val Asn Asn Pro  
 355 360 365

Thr Val Lys Val Tyr Lys His Ile Gly Ser Asp Asp Leu Ala Glu Ser  
 370 375 380

Val Tyr Ala Lys Leu Asp Asp Val Ser Lys Phe Glu Asp Val Thr Asp  
 385 390 395 400

Asn Met Ser Leu Asp Phe Asp Thr Asn Gly Gly Tyr Ser Leu Asn Phe  
 405 410 415

Asn Asn Leu Asp Gln Ser Lys Asn Tyr Val Ile Lys Tyr Glu Gly Tyr  
 420 425 430

Tyr Asp Ser Asn Ala Ser Asn Leu Glu Phe Gln Thr His Leu Phe Gly  
 435 440 445

Tyr Tyr Asn Tyr Tyr Tyr Thr Ser Asn Leu Thr Trp Lys Asn Gly Val  
 450 455 460

Ala Phe Tyr Ser Asn Asn Ala Gln Gly Asp Gly Lys Asp Lys Leu Lys  
 465 470 475 480

Glu Pro Ile Ile Glu His Ser Thr Pro Ile Glu Leu Glu Phe Lys Ser  
 485 490 495

Glu Pro Pro Val Glu Lys His Glu Leu Thr Gly Thr Ile Glu Glu Ser  
 500 505 510

Asn Asp Ser Lys Pro Ile Asp Phe Glu Tyr His Thr Ala Val Glu Gly  
 515 520 525

Ala Glu Gly His Ala Glu Gly Thr Ile Glu Thr Glu Glu Asp Ser Ile  
 530 535 540

His Val Asp Phe Glu Glu Ser Thr His Glu Asn Ser Lys His His Ala  
 545 550 555 560

Asp Val Val Glu Tyr Glu Glu Asp Thr Asn Pro Gly Gly Gly Gln Val  
 565 570 575

Thr Thr Glu Ser Asn Leu Val Glu Phe Asp Glu Asp Ser Thr Lys Gly  
 580 585 590

Ile Val Thr Gly Ala Val Ser Asp His Thr Thr Ile Glu Asp Thr Lys  
 595 600 605

-continued

---

Glu Tyr Thr Thr Glu Ser Asn Leu Ile Glu Leu Val Asp Glu Leu Pro  
 610 615 620

Glu Glu His Gly Gln Ala Gln Gly Pro Ile Glu Glu Ile Thr Glu Asn  
 625 630 635 640

Asn His His Ile Ser His Ser Gly Leu Gly Thr Glu Asn Gly His Gly  
 645 650 655

Asn Tyr Gly Val Ile Glu Glu Ile Glu Glu Asn Ser His Val Asp Ile  
 660 665 670

Lys Ser Glu Leu Gly  
 675

---

What is claimed is:

1. An isolated polypeptide comprising a variant *Staphylococcus aureus* Protein A (SpA) domain D segment having amino acid substitutions at amino acids corresponding to position 9, 10, 36 and 37 of SEQ ID NO: 2, wherein the substitutions at amino acids corresponding to positions 9 and 10 are with a glycine residue and the substitutions at amino acids corresponding to positions 36 and 37 are with a serine residue.
2. A pharmaceutical composition comprising the isolated polypeptide of claim 1 in a pharmaceutically acceptable formulation.
3. A method for eliciting an immune response against a *Staphylococcus aureus* bacterium in a subject comprising providing to the subject an effective amount of a composition comprising a variant *Staphylococcus aureus* Protein A (SpA) domain D segment having amino acid substitutions at amino acids corresponding to position 9, 10, 36 and 37 of SEQ ID NO: 2, wherein the substitutions at amino acids corresponding to positions 9 and 10 are with a glycine residue and the

substitutions at amino acids corresponding to positions 36 and 37 are with a serine residue.

4. The method of claim 3, where the subject is also administered an adjuvant.
5. The method of claim 3, wherein the staphylococcus bacterium is a *S. aureus* bacterium.
6. The method of claim 3, wherein the staphylococcus bacterium is resistant to one or more drug treatments.
7. The method of claim 6, wherein the bacterium is methicillin resistant.
8. The method of claim 3, wherein the composition is administered orally, parenterally, subcutaneously, intramuscularly, or intravenously.
9. The method of claim 3, further comprising administering to the subject a composition comprising a second staphylococcal antigen.
10. The method of claim 3, wherein the subject is human.
11. The method of claim 3, wherein the immune response is a protective immune response.

\* \* \* \* \*